

TYPES ARE WEAK ω -GROUPOIDS

BENNO VAN DEN BERG AND RICHARD GARNER

ABSTRACT. We define a notion of weak ω -category internal to a model of Martin-Löf type theory, and prove that each type bears a canonical weak ω -category structure obtained from the tower of iterated identity types over that type. We show that the ω -categories arising in this way are in fact ω -groupoids.

1. INTRODUCTION

It has long been understood that there is a close connection between algebraic topology and higher-dimensional category theory [10]. More recently, it has become apparent that both are in turn related to the intensional type theory of Martin-Löf [18]. Whilst attempts to make this link precise have only borne fruit in the past few years [1, 7, 9, 20], the basic idea dates back to an observation made by Hofmann and Streicher in [11]. Recall that in Martin-Löf type theory, we may construct from a type A and elements $a, b \in A$, a new type $\text{Id}(a, b)$ whose elements are to be thought of as proofs that a and b are propositionally equal. Hofmann and Streicher observe that the type-theoretic functions

$$1 \rightarrow \text{Id}(a, a), \quad \text{Id}(b, c) \times \text{Id}(a, b) \rightarrow \text{Id}(a, c) \quad \text{and} \quad \text{Id}(a, b) \rightarrow \text{Id}(b, a)$$

expressing the reflexivity, transitivity and symmetry of propositional equality allow us to view the type A as a *groupoid*—a category whose every morphism is invertible—wherein objects are elements $a \in A$ and morphisms $a \rightarrow b$ are elements $p \in \text{Id}(a, b)$. However, as is made clear in [11], this is not the end of the story. The groupoid axioms for A hold only “up to propositional equality”; which is to say that, for example, the associativity diagram

$$\begin{array}{ccc} \text{Id}(c, d) \times \text{Id}(b, c) \times \text{Id}(a, b) & \longrightarrow & \text{Id}(c, d) \times \text{Id}(a, c) \\ \downarrow & & \downarrow \\ \text{Id}(b, d) \times \text{Id}(a, b) & \longrightarrow & \text{Id}(a, d) \end{array}$$

does not commute on the nose, but only up to suitable terms

$$\alpha_{p,q,r} \in \text{Id}(r \circ (q \circ p), (r \circ q) \circ p) \quad (p \in \text{Id}(a, b), q \in \text{Id}(b, c), r \in \text{Id}(c, d)).$$

Thus, if we wish to view A as an honest groupoid, we must first quotient out the sets of elements $p \in \text{Id}(a, b)$ by propositional equality. A more familiar instance of the same phenomenon occurs in constructing the fundamental

Date: August 23, 2010.

2000 Mathematics Subject Classification. Primary: 03B15, 18D05; Secondary: 18D50.

groupoid of a space—where we must identify paths up to homotopy—and this suggests the following analogy: that types are like topological spaces, and that propositional equality is like the homotopy relation. Using the machinery of abstract homotopy theory, this analogy has been given a precise form in [1], which constructs type theoretic structures from homotopy theoretic ones, and in [7], which does the converse.

The connection with algebraic topology in turn suggests the one with higher-dimensional category theory. A more sophisticated construction of the fundamental groupoid of a space (suggested in [10] and made rigorous in [2]) does not quotient out paths by the homotopy relation; but instead incorporates these homotopies—and all higher homotopies between them—into an infinite-dimensional categorical structure known as a *weak ω -groupoid*, whose various identities, compositions and inverses satisfy coherence laws, not strictly, but “up to all higher homotopies”. This leads us to ask whether the construction of the type-theoretic “fundamental groupoid” admits a similar refinement, which constructs a weak ω -groupoid from a type by considering not just elements of the type, and proofs of their equality, but also proofs of equality between such proofs, and so on. The principal aim of this paper is to show this to be the case.

In order to give the proof, we must first choose an appropriate notion of weak ω -groupoid to work with; and since, in the literature, weak ω -groupoids are studied in the broader context of *weak ω -categories*—which are “weak ω -groupoids without the inverses”—this is tantamount to choosing an appropriate notion of weak ω -category. There are a number of definitions to pick from, and these differ from each other both in their general approach and in the details; see [14] for an overview. Of these, it is the definition of Batanin [2] which matches the type theory most closely; this for the following two reasons. Firstly, its basic cellular data are *globular*: which is to say that an n -cell $\alpha: x \rightarrow y$ can only exist between a pair of parallel $(n - 1)$ -cells $x, y: f \rightarrow g$. A corresponding property holds for proofs of equality in type theory: to know that $\alpha \in \text{Id}(x, y)$, we must first know that x and y inhabit the same type $\text{Id}(f, g)$. Secondly, Batanin’s definition is *algebraic*: which is to say that composition operations are explicitly specified, rather than merely asserted to exist. This accords with the constructivist notion—central to the spirit of intensional type theory—that to know something to exist is nothing less than to be provided with a witness to that fact. On these grounds, it is Batanin’s definition which we will adopt here; or rather, a mild reformulation of his definition given by Leinster in [14].

The paper is arranged as follows. In Section 2, we recall Batanin’s theory of weak ω -categories, the appropriate specialisation to weak ω -groupoids, and the necessary background from intensional type theory. Then in Section 3 we give the proof of our main result. We begin in Section 3.1 with an explicitly type-theoretic, but informal, account. When we come to make this precise, it turns out to be convenient to isolate just those categorical properties of the type theory which make the proof go through, and then to work in an axiomatic setting assuming only these. We describe this setting in Section 3.2, and then in Sections 3.3 and 3.4, use it to give a formal proof that every type is a weak ω -groupoid.

It seems appropriate to say a few words about the history of this paper. The main result was described by the first-named author in 2006 in a presentation at the workshop “Identity Types—Topological and Categorical Structure” held at Uppsala University [3]. The details of the proof were then worked out by both authors during a 2008 visit by the first author to Uppsala; and it was at this stage that the axiomatic approach was introduced. Whilst preparing this manuscript for publication, we become aware that, independently, Peter Lumsdaine had been considering the same question. His analysis may be found in [17]. Let us remark only that, where our argument is category-theoretic in nature, that given by Lumsdaine is essentially proof-theoretic. We gratefully acknowledge the support of Uppsala University’s Department of Mathematics, and extend our thanks to Erik Palmgren for organising the aforementioned workshop. The second-named author also acknowledges the support of a Research Fellowship of St John’s College, Cambridge and a Marie Curie Intra-European Fellowship, Project No. 040802.

2. PREPARATORY MATERIAL

In this section, we review the material necessary for our main result; firstly, from higher category theory, and secondly, from Martin-Löf type theory.

2.1. Weak ω -categories and weak ω -groupoids. As mentioned in the Introduction, the most appropriate definition of weak ω -category for our purposes is that of [2], which describes them as globular sets equipped with algebraic structure. A *globular set* is a diagram of sets and functions

$$X_0 \begin{array}{c} \xleftarrow{s} \\ \xleftarrow{t} \end{array} X_1 \begin{array}{c} \xleftarrow{s} \\ \xleftarrow{t} \end{array} X_2 \begin{array}{c} \xleftarrow{s} \\ \xleftarrow{t} \end{array} X_3 \begin{array}{c} \xleftarrow{s} \\ \xleftarrow{t} \end{array} \cdots$$

satisfying the globularity equations $ss = st$ and $ts = tt$. We refer to elements $x \in X_n$ as *n-cells* of X , and write them as $x: sx \rightarrow tx$. In this terminology, the globularity equations express that any $(n+2)$ -cell $f \rightarrow g$ must mediate between $(n+1)$ -cells f and g which are *parallel*, in the sense of having the same source and target. Globular sets also have a coinductive characterisation: to give a globular set X is to give a set $\text{ob } X$ of objects, and for each $x, y \in \text{ob } X$, a globular set $X(x, y)$.

The algebraic structure required to make a globular set into a weak ω -category is encoded by any one of a certain class of monads on the category of globular sets: those arising from *normalised, contractible, globular operads*. Informally, such monads are obtained by “deforming” the monad T whose algebras are strict ω -categories. To make this precise, we must first recall some details concerning strict ω -categories.

If \mathcal{V} is any category with finite products, then one can speak of categories enriched in \mathcal{V} , and of \mathcal{V} -enriched functors between them [13]. The category $\mathcal{V}\text{-Cat}$ of small \mathcal{V} -categories is then itself a category with finite products, so that we can iterate the process; and when we do so starting from $\mathcal{V} = 1$, we obtain the sequence 1, Set, Cat, 2-Cat, \dots , whose n th term is the category of small strict $(n-1)$ -categories. Now, because any finite-product preserving

functor $\mathcal{V} \rightarrow \mathcal{W}$ induces a finite-product preserving functor $\mathcal{V}\text{-Cat} \rightarrow \mathcal{W}\text{-Cat}$, we obtain, by iteration on the unique functor $\text{Set} \rightarrow 1$, a chain

$$\cdots \rightarrow n\text{-Cat} \cdots \rightarrow 2\text{-Cat} \rightarrow \text{Cat} \rightarrow \text{Set} \rightarrow 1 ;$$

and $\omega\text{-Cat}$, the category of small strict ω -categories, is the limit of this sequence. Unfolding this definition, we find that a strict ω -category is given by first, an underlying globular set; next, operations of identity and composition: so for each n -cell x , an $(n+1)$ -cell $\text{id}_x: x \rightarrow x$, and for each pair of n -cells f and g sharing a k -cell boundary (for $k < n$), a composite n -cell $g \circ_k f$; and finally, axioms which express that any two ways of composing a diagram of n -cells using the above operations yield the same result.

There is an evident forgetful functor $U: \omega\text{-Cat} \rightarrow \text{GSet}$, where GSet denotes the category of globular sets; and it is shown in [16, Appendix B] that this has a left adjoint and is finitarily monadic. The corresponding monad T on the category of globular sets may be described as follows. First we give an inductive characterisation of $T1$, its value at the globular set with one cell in every dimension. We have:

- $(T1)_0 = \{\star\}$; and
- $(T1)_{n+1} = \{(\pi_1, \dots, \pi_k) \mid k \in \mathbb{N}, \pi_1, \dots, \pi_k \in (T1)_n\}$.

The source and target maps $s, t: (T1)_{n+1} \rightarrow (T1)_n$ coincide and we follow [16] in writing ∂ for the common value. This too may be described inductively:

- $\partial(\pi) = \star$ for $\pi \in (T1)_1$; and
- $\partial(\pi_1, \dots, \pi_k) = (\partial(\pi_1), \dots, \partial(\pi_k))$ otherwise.

We regard elements of $(T1)_n$ as indexing possible shapes for pasting diagrams of n -cells. For example, $((*), (*, *)) \in (T1)_2$ corresponds to the shape

$$(1) \quad \bullet \begin{array}{c} \curvearrowright \\ \Downarrow \\ \curvearrowleft \end{array} \bullet \begin{array}{c} \curvearrowright \\ \Downarrow \\ \curvearrowleft \end{array} \bullet .$$

We can make this formal as follows. By induction, we associate to each element $\pi \in (T1)_n$ a globular set $\hat{\pi}$ which is the “shape indexed by π ”:

- If $\pi = \star$, then $\hat{\pi}$ is the globular set with $\text{ob } \hat{\pi} = \{\bullet\}$ and $\hat{\pi}(\bullet, \bullet) = \emptyset$.
- If $\pi = (\pi_1, \dots, \pi_k)$, then $\hat{\pi}$ is the globular set with $\text{ob } \hat{\pi} = \{0, \dots, k\}$, $\hat{\pi}(i-1, i) = \hat{\pi}_i$ (for $1 \leq i \leq k$), and $\hat{\pi}(i, j) = \emptyset$ otherwise.

By a further induction, we define source and target embeddings $\sigma, \tau: \widehat{\partial\pi} \rightarrow \hat{\pi}$:

- For $\pi \in (T1)_1$, the maps $\sigma, \tau: \hat{\star} \rightarrow \hat{\pi}$ send the unique object of $\hat{\star}$ to the smallest and largest elements of $\text{ob } \hat{\pi}$, respectively.
- Otherwise, for $\pi = (\pi_1, \dots, \pi_k)$ the morphisms σ and τ are the identity on objects and map $\widehat{\partial\pi}(i-1, i)$ into $\hat{\pi}(i-1, i)$ via $\sigma, \tau: \widehat{\partial\pi}_i \rightarrow \hat{\pi}_i$.

Taken together, these data—the globular set $T1$, the globular sets $\hat{\pi}$ and the maps σ and τ —completely determine the functor T ; this by virtue of it being *familially representable* in the sense of [15, Definition C.3.1] (though see also [5]). Explicitly, TX is the globular set whose cells are pasting diagrams

labelled with cells of X :

$$(TX)_n = \sum_{\pi \in (T1)_n} \text{GSet}(\hat{\pi}, X),$$

and whose source and target maps are induced in an obvious way by the maps σ and τ . The unit and multiplication of the monad T are *cartesian* natural transformations—which is to say that all of their naturality squares are pullbacks—from which it follows that these are in turn determined by the components $\eta_1: 1 \rightarrow T1$ and $\mu_1: TT1 \rightarrow T1$. The former map associates to the unique n -cell of 1 the pasting diagram $\iota_n := (\cdots(\star)\cdots) \in (T1)_n$, whilst the latter sends a typical element

$$(\pi \in (T1)_n, \phi: \hat{\pi} \rightarrow T1)$$

of $(TT1)_n$ to the element $\phi \circ \pi \in (T1)_n$ obtained by substituting into π the pasting diagrams which ϕ indexes (see [16, Section 4.2] for a pictorial account of this process).

A *globular operad* can now be defined rather succinctly: it is a monad P on GSet equipped with a cartesian monad morphism $\rho: P \Rightarrow T$. The cartesianness of ρ implies that the functor part of P is determined by its component at 1 together with the augmentation map $\rho_1: P1 \rightarrow T1$, and it will be convenient to have a description of P in these terms. Given $\pi \in (T1)_n$, we write P_π for the set of those $\theta \in (P1)_n$ which are mapped to π by ρ_1 , and write $s, t: P_\pi \rightarrow P_{\partial\pi}$ for the corresponding restriction of the source and target maps of $P1$. The value of P at an arbitrary globular set X is now given (up to isomorphism) by

$$(2) \quad (PX)_n = \sum_{\pi \in (T1)_n} P_\pi \times \text{GSet}(\hat{\pi}, X),$$

with the source and target maps determined in the obvious way. Thus, if we think of a T -algebra structure on X as providing a unique way of composing each X -labelled pasting diagram of shape π , then a P -algebra structure provides a set of possible ways of composing such diagrams, indexed by the elements of P_π .

It follows from the cartesianness of ρ that the unit and the multiplication of P are themselves cartesian natural transformations, and hence determined by their components $\eta_1: 1 \rightarrow P1$ and $\mu_1: PP1 \rightarrow P1$. The former sends the unique n -cell of 1 to an element $\iota_n \in P_{\iota_n}$, which we think of as the trivial composition operation of dimension n ; whilst the latter assigns to the element

$$(\pi \in (T1)_n, \theta \in P_\pi, \psi: \hat{\pi} \rightarrow P1)$$

of $(PP1)_n$ an element $\theta \circ \psi \in P_{\phi \circ \pi}$ (where ϕ is the composite $\rho_1 \psi: \hat{\pi} \rightarrow T1$), which we think of as the composition operation obtained by substituting into θ the collection of operations indexed by ψ .

Not every globular operad embodies a sensible theory of weak ω -categories—since, for example, the identity monad on GSet is a globular operad—but [2] provides two conditions which together distinguish those which do: *normalisation* and *contractibility*. Normalisation is straightforward; it asserts that the monad P is bijective on objects in the sense that $(PX)_0 \cong X_0$, naturally in

X , or equivalently, that the set P_\star is a singleton. The second condition is a little more subtle. A globular operad P is said to be *contractible* if:

- (a) Given $\pi \in (T1)_1$ and $\theta_1, \theta_2 \in P_\star$, there exists an element $\phi \in P_\pi$ with $s(\phi) = \theta_1$ and $t(\phi) = \theta_2$;
- (b) Given $\pi \in (T1)_n$ (for $n > 1$) and $\theta_1, \theta_2 \in P_{\partial\pi}$ satisfying $s(\theta_1) = s(\theta_2)$ and $t(\theta_1) = t(\theta_2)$, there exists an element $\phi \in P_\pi$ such that $s(\phi) = \theta_1$ and $t(\phi) = \theta_2$.

Contractibility expresses that that a globular operad has “enough” ways of composing to yield a theory of weak ω -categories. In homotopy-theoretic terms, a contractible globular operad is a “deformation” of the monad T ; an idea which can be made precise using the language of weak factorisation systems: see [8].

2.1.1. Definition. A *weak ω -category* is an algebra for a contractible, normalised, globular operad: more formally, it is a pair (P, X) , where P is a contractible, normalised, globular operad and X is an algebra for it.

2.1.2. Remark. Some consideration must be paid to the exact force of the term *contractible*, which has been used in different ways by different authors; our usage accords with that of [15, Definition 9.1.3]. In particular, the reader should carefully distinguish between the *property* of being contractible described above, and the corresponding *structure* of being equipped with a contraction.

We now turn from the definition of weak ω -category to that of weak ω -groupoid. For this we will require the coinductive notion of *equivalence* in a weak ω -category.

2.1.3. Definition. Let (P, X) be a weak ω -category. An *equivalence* $x \simeq y$ between parallel n -cells x, y is given by:

- $n + 1$ -cells $f: x \rightarrow y$ and $g: y \rightarrow x$;
- Equivalences $\eta: g \circ f \simeq \text{id}_x$ and $\epsilon: f \circ g \simeq \text{id}_y$.

We say that an $(n + 1)$ -cell $f: x \rightarrow y$ is *weakly invertible* if it participates in an equivalence (f, g, η, ϵ) .

In order for this definition to make sense, we must determine what is meant by the expressions “ id_x ”, “ id_y ”, “ $g \circ f$ ” and “ $f \circ g$ ” appearing in it, which we may do as follows. First, for each $n \geq 1$, we define the pasting diagrams 0_n and $2_n \in (T1)_n$ to be given by

$$0_n := \underbrace{(\cdots () \cdots)}_{n \text{ times}} \quad \text{and} \quad 2_n := \underbrace{(\cdots (\star, \star) \cdots)}_{n \text{ times}} .$$

Next, if P is a normalised, contractible globular operad, then we define a *system of compositions* for P to be a choice, for each $n \geq 1$, of operations $i_n \in P_{0_n}$ and $m_n \in P_{2_n}$. Note that the contractibility of P ensures that it will possess at least one system of compositions. Finally, if we are given a system of compositions and a P -algebra X , then we define the functions

$$\text{id}_{(-)}: X_{n-1} \rightarrow X_n \quad \text{and} \quad \circ: X_{n+1} \times_t X_{n+1} \rightarrow X_n$$

to be the interpretations of the operations i_n and m_n respectively. This allows us to give meaning to the undefined expressions appearing in Definition 2.1.3.

2.1.4. Definition. A weak ω -category (P, X) is a *weak ω -groupoid* if every cell of X is weakly invertible with respect to every system of compositions on P .

It will be convenient to give a more elementary reformulation of the notion of weak ω -groupoid due to Cheng [6]. This is given in terms of *duals*. If $f: x \rightarrow y$ is an n -cell (for $n \geq 1$) in a weak ω -category, then a *dual* for f is an n -cell $f^*: y \rightarrow x$ together with $(n+1)$ -cells $\eta: \text{id}_x \rightarrow f^* \circ f$ and $\epsilon: f \circ f^* \rightarrow \text{id}_y$, subject to no axioms. Again, this definition is to be interpreted with respect to some given system of compositions.

2.1.5. Proposition. *A weak ω -category is a weak ω -groupoid if and only if, with respect to every system of compositions, every cell has a dual.*

Proof. By coinduction. □

2.2. Martin-Löf type theory. By *intensional Martin-Löf type theory*, we mean the logical calculus set out in Part II of [18]. We now summarise this calculus. It has four basic forms of judgement: **A type** (“ A is a type”); **$a \in A$** (“ a is an element of the type A ”); **$A = B$ type** (“ A and B are definitionally equal types”); and **$a = b \in A$** (“ a and b are definitionally equal elements of the type A ”). These judgements may be made either absolutely, or relative to a context Γ of assumptions, in which case we write them as

$$(\Gamma) A \text{ type}, \quad (\Gamma) a \in A, \quad (\Gamma) A = B \text{ type} \quad \text{and} \quad (\Gamma) a = b \in A$$

respectively. Here, a *context* is a list $\Gamma = (x_1 \in A_1, x_2 \in A_2, \dots, x_n \in A_{n-1})$, wherein each A_i is a type relative to the context $(x_1 \in A_1, \dots, x_{i-1} \in A_{i-1})$. There are now some rather natural requirements for well-formed judgements: in order to assert that $a \in A$ we must first know that **A type**; to assert that **$A = B$ type** we must first know that **A type** and **B type**; and so on. We specify intensional Martin-Löf type theory as a collection of inference rules over these forms of judgement. Firstly we have the *equality rules*, which assert that the two judgement forms **$A = B$ type** and **$a = b \in A$** are congruences with respect to all the other operations of the theory; then we have the *structural rules*, which deal with weakening, contraction, exchange and substitution; and finally, the *logical rules*, which specify the type-formers of our theory, together with their introduction, elimination and computation rules. For the purposes of this paper, we require only the rules for the identity types, which we list in Table 1. We commit the usual abuse of notation in leaving implicit an ambient context Γ common to the premisses and conclusions of each rule, and omitting the rules expressing stability under substitution in this ambient context. Let us remark also that in the rules **Id-ELIM** and **Id-COMP** we allow the type C over which elimination is occurring to depend upon an additional contextual parameter Δ . Were we to add Π -types (dependent products) to our calculus, then these rules would be equivalent to the usual identity type rules. However, in the absence of Π -types, this extra parameter is essential to derive all but the most basic properties of the identity type.

$$\begin{array}{c}
\textit{Identity types} \\
\frac{A \text{ type} \quad a, b \in A}{\text{Id}_A(a, b) \text{ type}} \text{Id-FORM}; \quad \frac{A \text{ type} \quad a \in A}{r(a) \in \text{Id}_A(a, a)} \text{Id-INTRO}; \\
\frac{\begin{array}{c} (x, y \in A, p \in \text{Id}_A(x, y), \Delta(x, y, p)) C(x, y, p) \text{ type} \\ (x \in A, \Delta(x, x, r(x))) d(x) \in C(x, x, r(x)) \\ a, b \in A \quad p \in \text{Id}_A(a, b) \end{array}}{(\Delta(a, b, p)) J_d(a, b, p) \in C(a, b, p)} \text{Id-ELIM}; \\
\frac{\begin{array}{c} (x, y \in A, p \in \text{Id}_A(x, y), \Delta(x, y, p)) C(x, y, p) \text{ type} \\ (x \in A, \Delta(x, x, r(x))) d(x) \in C(x, x, r(x)) \quad a \in A \end{array}}{(\Delta(a, a, r(a))) J_a(a, a, r(a)) = d(a) \in C(a, a, r(a))} \text{Id-COMP}.
\end{array}$$

TABLE 1. Identity type rules

We now establish some further notational conventions. Where it improves clarity we may omit brackets in function applications, writing $hgfx$ in place of $h(g(f(x)))$, for example. We may drop the subscript A in an identity type $\text{Id}_A(a, b)$ where no confusion seems likely to occur. Given $a, b \in A$, we may say that a and b are *propositionally equal* to indicate that the type $\text{Id}(a, b)$ is inhabited. We will also make use of *vector notation* in the style of [4]. Given a context $\Gamma = (x_1 \in A_1, \dots, x_n \in A_n)$, we may abbreviate a series of judgements:

$$a_1 \in A_1, \quad a_2 \in A_2(a_1), \quad \dots \quad a_n \in A_n(a_1, \dots, a_{n-1}),$$

as $a \in \Gamma$, where $a := (a_1, \dots, a_n)$. We may also use this notation to abbreviate sequences of hypothetical elements; so, for example, we may specify a dependent type in context Γ as $(x \in \Gamma) A(x) \text{ type}$. We will also make use of [4]'s notion of *telescope*. Given Γ a context as before, this allows us to abbreviate the series of judgements

$$\begin{array}{c}
(x \in \Gamma) B_1(x) \text{ type}, \\
(x \in \Gamma, y_1 \in B_1) B_2(x, y_1) \text{ type}, \\
\vdots \\
(x \in \Gamma, y_1 \in B_1, \dots, y_{m-1} \in B_{m-1}) B_m(x, y_1, \dots, y_{m-1}) \text{ type}.
\end{array}$$

as $(x \in \Gamma) \Delta(x) \text{ ctxt}$, where $\Delta(x) := (y_1 \in B_1(x), y_2 \in B_2(x, y_1), \dots)$. We say that Δ is a *context dependent upon* Γ . Given such a dependent context, we may abbreviate the series of judgements

$$\begin{array}{c}
(x \in \Gamma) f_1(x) \in B_1(x) \\
\vdots \\
(x \in \Gamma) f_m(x) \in B_m(x, f_1(x), \dots, f_{m-1}(x)),
\end{array}$$

as $(x \in \Gamma) f(x) \in \Delta(x)$, and say that f is a *dependent element* of Δ . We can similarly assign a meaning to the judgements $(x \in \Gamma) \Delta(x) = \Theta(x) \text{ ctxt}$

and $(x \in \Gamma) f(x) = g(x) \in \Delta(x)$, expressing the definitional equality of two dependent contexts, and the definitional equality of two dependent elements of a dependent context.

Let us now recall some basic facts about categorical models of type theory. For a more detailed treatment we refer the reader to [12, 19], for example. If \mathbb{T} is a dependently typed calculus admitting each of the rules described above, then we may construct from it a category $\mathcal{C}_{\mathbb{T}}$ known as the *classifying category* of \mathbb{T} . Its objects are contexts Γ, Δ, \dots , in \mathbb{T} , considered modulo definitional equality (so we identify Γ and Δ whenever $\Gamma = \Delta$ ctxt is derivable); and its map $\Gamma \rightarrow \Delta$ are *context morphisms*, which are judgements $(x \in \Gamma) f(x) \in \Delta$ considered modulo definitional equality. The identity map on Γ is given by $(x \in \Gamma) x \in \Gamma$; whilst composition is given by substitution of terms. Now, for any judgement $(x \in \Gamma) A(x)$ type of \mathbb{T} , there is a distinguished context morphism

$$(x \in \Gamma, y \in A(x)) \rightarrow (x \in \Gamma)$$

which sends (x, y) to x . We call morphisms of $\mathcal{C}_{\mathbb{T}}$ of this form *basic dependent projections*. By a *dependent projection*, we mean any composite of zero or more basic dependent projections. An important property of dependent projections is that they are stable under pullback, in the sense that for every $(x \in \Gamma) A(x)$ type and context morphism $f: \Delta \rightarrow \Gamma$, we may show the square

$$\begin{array}{ccc} (w \in \Delta, y \in A(f(w))) & \longrightarrow & (x \in \Gamma, y \in A(x)) \\ p' \downarrow & & \downarrow p \\ \Delta & \xrightarrow{f} & \Gamma \end{array},$$

wherein the uppermost arrow sends (w, y) to (fw, y) , to be a pullback in $\mathcal{C}_{\mathbb{T}}$. Let us now recall from [7] a second class of maps in $\mathcal{C}_{\mathbb{T}}$ which will play an important role in this paper. A context morphism $f: \Gamma \rightarrow \Delta$ is said to be an *injective equivalence* if it validates type-theoretic rules:

$$\frac{(y \in \Delta) \Lambda(y) \text{ ctxt} \quad (x \in \Gamma) d(x) \in \Lambda(f(x)) \quad b \in \Delta}{E_d(b) \in \Lambda(b)}$$

and

$$\frac{(y \in \Delta) \Lambda(y) \text{ ctxt} \quad (x \in \Gamma) d(x) \in \Lambda(f(x)) \quad a \in \Gamma}{E_d(f(a)) = d(a) \in \Lambda(f(a))}.$$

The name is motivated by the groupoid model of type theory, wherein the injective equivalences are precisely the injective groupoid equivalences. Intuitively, a morphism $f: \Gamma \rightarrow \Delta$ is an injective equivalence just when every (dependent) function out of Δ is determined, up to propositional equality, by its restriction to Γ . The leading example of an injective equivalence is given by the context morphism $A \rightarrow (x, y \in A, p \in \text{Id}(x, y))$ sending x to (x, x, rx) . That this map is an injective equivalence is precisely the content of the Id-elimination and computation rules. Diagrammatically, a map f is an injective equivalence if for

every commutative square of the form

$$\begin{array}{ccc} \Gamma & \xrightarrow{d} & (\Delta, \Lambda) \\ h \downarrow & & \downarrow p \\ \Delta & \xlongequal{\quad} & \Delta \end{array}$$

with p a dependent projection, we may find a diagonal filler $E_d: \Delta \rightarrow (\Delta, \Lambda)$ making both induced triangles commute. By the stability of dependent projections under pullback, this is equivalent with the property that we should be able to find fillers for all commutative squares of the form

$$(3) \quad \begin{array}{ccc} \Gamma & \xrightarrow{d} & (\Phi, \Lambda) \\ h \downarrow & & \downarrow p \\ \Delta & \xrightarrow{k} & \Phi \end{array}$$

again with p a dependent projection. See [7, Section 5] for an elementary characterisation of the class of injective equivalences.

3. THE MAIN RESULT

3.1. An overview of the proof. We are now ready to begin the proof of our main result: that if \mathbb{T} is a dependently typed calculus admitting each of the rules described in Section 2.2, then each type A therein gives rise to a weak ω -groupoid whose objects are elements of A , and whose higher cells are elements of the iterated identity types on A . In fact, we will be able to prove a stronger result: that A provides the “type of objects” for a weak ω -groupoid which is, in a suitable sense, *internal* to \mathbb{T} .

As explained in the Introduction, we will give our proof twice: once informally, using a type-theoretic language, and once formally, using an axiomatic categorical framework which captures just those aspects of the type theory which allow the proof to go through. In this Section, we give the informal proof. We shall concentrate in the first instance on constructing a weak ω -category, and defer the question of whether or not it is a weak ω -groupoid until the formal proof.

We begin by defining what we mean by a weak ω -category internal to a type theory \mathbb{T} . More specifically, given some globular operad P , we define a notion of P -algebra internal to \mathbb{T} . The underlying data for such a P -algebra is a *globular context* $(\Delta) \Gamma \in \mathbb{T}$; which is a sequence of judgements

$$\begin{aligned} & (\Delta) \Gamma_0 \text{ ctxt} \\ & (\Delta, x, y \in \Gamma_0) \Gamma_1(x, y) \text{ ctxt} \\ & (\Delta, x, y \in \Gamma_0, p, q \in \Gamma_1(x, y)) \Gamma_2(x, y, p, q) \text{ ctxt} \\ & \vdots \end{aligned}$$

Just as globular sets have a coinductive characterisation, so too do globular contexts: to give a globular context $(\Delta) \Gamma$ is to give a context $(\Delta) \Gamma_0$ together

with a globular context $(\Delta, x, y : \Gamma_0) \Gamma_{+1}(x, y)$. In order to define the operations making a globular context Γ (where henceforth we simplify the notation by omitting the precontext Δ) into a P -algebra, we first define for each pasting diagram $\pi \in (T1)_n$ the context Γ^π consisting of “ π -indexed elements of Γ ”. This is done by induction on π :

- If $\pi = \star$, then $\Gamma^\pi := \Gamma_0$;
- If $\pi = (\pi_1, \dots, \pi_k)$, then Γ^π is the context

$$(x_0, \dots, x_k \in \Gamma_0, y_1 \in \Gamma_{+1}(x_0, x_1)^{\pi_1}, \dots, y_k \in \Gamma_{+1}(x_{k-1}, x_k)^{\pi_k}) .$$

For example, if π is the pasting diagram (1), then the context Γ^π is given by:

$$\begin{aligned} & (x_0, x_1, x_2 \in \Gamma_0, \\ & \quad s, t \in \Gamma_1(x_0, x_1), \alpha \in \Gamma_2(x_0, x_1, s, t), \\ & \quad u, v, w \in \Gamma_1(x_1, x_2), \beta \in \Gamma_2(x_1, x_2, u, v), \gamma \in \Gamma_2(x_1, x_2, v, w)) \end{aligned}$$

whilst if $\pi \in (T1)_n$ is the element $\iota_n = (\dots (\star) \dots)$, then Γ^{ι_n} is the context

$$(x_0, y_0 \in \Gamma_0, x_1, y_1 \in \Gamma_1(x_0, y_0), \dots, x_n \in \Gamma_n(x_0, y_0, \dots, x_{n-1}, y_{n-1}))$$

indexing the totality of the n -cells of Γ . Now to give a P -algebra structure on the globular context Γ will be to give, for every $\pi \in (T1)_n$ and $\theta \in P_\pi$, a context morphism

$$[\theta]: \Gamma^\pi \rightarrow \Gamma^{\iota_n}$$

interpreting the operation θ , subject to the following axioms. Firstly, the interpretations should be compatible with source and target, which is to say that diagrams of the form

$$\begin{array}{ccc} \Gamma^\pi & \xrightarrow{[\theta]} & \Gamma^{\iota_n} \\ \sigma \downarrow & & \downarrow \sigma \\ \Gamma^{\partial\pi} & \xrightarrow{[s\theta]} & \Gamma^{\iota_{n-1}} \end{array} \quad \text{and} \quad \begin{array}{ccc} \Gamma^\pi & \xrightarrow{[\theta]} & \Gamma^{\iota_n} \\ \tau \downarrow & & \downarrow \tau \\ \Gamma^{\partial\pi} & \xrightarrow{[t\theta]} & \Gamma^{\iota_{n-1}} \end{array}$$

should commute; here, $\sigma, \tau: \Gamma^\pi \rightarrow \Gamma^{\partial\pi}$ are source and target projections defined by a further straightforward induction over π . Secondly, the trivial pasting operations should have a trivial interpretation; which is to say that

$$[\iota_n] = \text{id}_{\Gamma^{\iota_n}} : \Gamma^{\iota_n} \rightarrow \Gamma^{\iota_n} .$$

Thirdly, the interpretation of a composite $[\theta \circ \psi]$ should be “given by the composite of $[\theta]$ with $[\psi]$ ”, in the sense that the following diagram commutes:

$$\begin{array}{ccc} \Gamma^{\pi \circ \phi} & \xrightarrow{[\psi]} & \Gamma^\pi \\ & \searrow [\theta \circ \psi] & \downarrow [\theta] \\ & & \Gamma^{\iota_n} \end{array} .$$

This is not yet entirely formal, because we have not indicated how the map $[\psi]: \Gamma^{\pi \circ \phi} \rightarrow \Gamma^\pi$ should be defined. Intuitively, it is the morphism which applies simultaneously the interpretations of the operations indexed by $\psi: \hat{\pi} \rightarrow P1$; but it is not immediately clear how to make this precise. We will do so in

Section 3.3 below, using Michael Batanin’s machinery of *monoidal globular categories* [2]. A general result from this theory allows us to associate to the globular context Γ a particular globular operad $[\Gamma, \Gamma]$ —the *endomorphism operad* of Γ —which is such that we may *define* P -algebra structures on Γ to be globular operad morphisms $P \rightarrow [\Gamma, \Gamma]$. This operad $[\Gamma, \Gamma]$ has as operations of shape π , all serially commutative diagrams

$$(4) \quad \begin{array}{ccc} \Gamma^\pi & \xrightarrow{f_n} & \Gamma^{\iota_n} \\ \sigma \downarrow \tau & & \sigma \downarrow \tau \\ \Gamma^{\partial\pi} & \xrightarrow[g_{n-1}]{} & \Gamma^{\iota_{n-1}} \\ \sigma \downarrow \tau & & \sigma \downarrow \tau \\ \vdots & & \vdots \\ \Gamma^* & \xrightarrow[g_0]{} & \Gamma^* \end{array}$$

of context morphisms. The source and target functions $[\Gamma, \Gamma]_\pi \rightarrow [\Gamma, \Gamma]_{\partial\pi}$ send such a diagram to its sub-diagram headed by f_{n-1} , respectively g_{n-1} ; the identity operation $\iota_n \in [\Gamma, \Gamma]_{\iota_n}$ has each f_i and g_i given by an identity map; whilst to describe substitution of operations in $[\Gamma, \Gamma]$ is precisely the problem that we encountered above, and that which Batanin’s machinery solves. It is easy to see that a map of globular operads $P \rightarrow [\Gamma, \Gamma]$ encodes exactly the structure of an internal P -algebra sketched above.

We may now give a precise statement of the main result. Given a type theory \mathbb{T} admitting the rules of Section 2.2 and a type $A \in \mathbb{T}$, we will construct a normalised, contractible, globular operad P such that the globular context \underline{A} given by

$$\begin{array}{l} A \text{ ctxt} \\ (x, y \in A) \text{ Id}_A(x, y) \text{ ctxt} \\ (x, y \in A, p, q \in \text{Id}_A(x, y)) \text{ Id}_{\text{Id}_A(x, y)}(p, q) \text{ ctxt} \\ \vdots \end{array}$$

admits an internal P -algebra structure. Now, it is straightforward to find an operad for which \underline{A} is an algebra—namely, the endomorphism operad $[\underline{A}, \underline{A}]$, with algebra structure given by the identity morphism $[\underline{A}, \underline{A}] \rightarrow [\underline{A}, \underline{A}]$ —but this does not help us, since there is no reason to expect this operad to be either normalised or contractible. However, it comes rather close to being contractible, in a sense which we will now explain. For $[\underline{A}, \underline{A}]$ to be contractible

would be for us to ask that, for every serially commutative diagram

$$(5) \quad \begin{array}{ccc} \underline{A}^\pi & & \underline{A}^{t_n} \\ \sigma \downarrow \tau & & \sigma \downarrow \tau \\ \underline{A}^{\partial\pi} & \xrightarrow[f_{n-1}]{g_{n-1}} & \underline{A}^{t_{n-1}} \\ \sigma \downarrow \tau & & \sigma \downarrow \tau \\ \vdots & & \vdots \\ \sigma \downarrow \tau & \xrightarrow[f_0]{g_0} & \underline{A}^* \\ \sigma \downarrow \tau & & \sigma \downarrow \tau \end{array}$$

of context morphisms, we could find a map $\underline{A}^\pi \rightarrow \underline{A}^{t_n}$ completing it to a diagram like (4). Let us consider in particular the case where π is the pasting diagram of (1). Here, to give the data of (5) is to give judgements

$$(6) \quad \begin{aligned} & (x \in A) \ f_0(x) \in A \\ & (x \in A) \ g_0(x) \in A \\ & (x, y, z \in A, p \in \text{Id}(x, y), q \in \text{Id}(y, z)) \ f_1(x, y, z, p, q) \in \text{Id}(f_0(x), g_0(z)) \\ & (x, y, z \in A, p \in \text{Id}(x, y), q \in \text{Id}(y, z)) \ g_1(x, y, z, p, q) \in \text{Id}(f_0(x), g_0(z)) \end{aligned}$$

whilst to give its completion $f_2: \underline{A}^\pi \rightarrow \underline{A}^{t_2}$ would be to give a judgement

$$\begin{aligned} & (x, y, z \in A, \\ & \quad s, t \in \text{Id}(x, y), \alpha \in \text{Id}(s, t), \\ & \quad u, v, w \in \text{Id}(y, z), \beta \in \text{Id}(u, v), \gamma \in \text{Id}(v, w)) \\ & \quad f_2(x, y, z, s, t, \alpha, u, v, w, \beta, \gamma) \in \text{Id}(f_1(x, y, z, s, u), g_1(x, y, z, t, w)) . \end{aligned}$$

We might attempt to obtain such a judgement by repeated application of the identity type elimination rule. Indeed, by Id-elimination on α it suffices to consider the case where $s = t$ and $\alpha = r(s)$; and by Id-elimination on γ and β , it suffices to consider the case where $u = v = w$ and $\gamma = \beta = r(u)$. Thus it suffices to find a term

$$\begin{aligned} & (x, y, z \in A, s \in \text{Id}(x, y), u \in \text{Id}(y, z)) \\ & \quad f_2'(x, y, z, s, u) \in \text{Id}(f_1(x, y, z, s, u), g_1(x, y, z, s, u)) . \end{aligned}$$

But now by Id-elimination on s and on u , it suffices to consider the case where $x = y = z$ and $s = u = r(x)$; so that it even suffices to find a term

$$(7) \quad (x \in A) \ f_2''(x) \in \text{Id}(f_1(x, x, x, rx, rx), g_1(x, x, x, rx, rx)) .$$

Yet here we encounter the problem that f_1 and g_1 , being arbitrarily defined, need not agree at (x, x, x, rx, rx) , so that there is in general no reason for a term like (7) to exist. However, there is a straightforward way of removing this obstruction: we restrict attention to those operations of shape π which, when applied to a term consisting solely of reflexivity proofs, yield another

reflexivity proof. We may formalise this as follows. For each $\pi \in (T1)_n$, we define, by induction on π , a pointing $r_\pi: A \rightarrow \underline{A}^\pi$:

- If $\pi = \star$, then $r_\star := \text{id}: A \rightarrow A$;
- If $\pi = (\pi_1, \dots, \pi_k)$, then r_π is the context morphism

$$(x \in A) \underbrace{(x, \dots, x, r_{\pi_1}(rx), \dots, r_{\pi_k}(rx))}_{k \text{ times}} \in \underline{A}^\pi .$$

In our example, if the judgements in (6) commuted with the A -pointings, then we would have that $f_1(x, x, x, rx, rx) = g_1(x, x, x, rx, rx) = r(x) \in \text{Id}(x, x)$, so that in (7) we could define

$$(x \in A) f_2''(x) := r(rx) \in \text{Id}(rx, rx)$$

and in this way obtain by repeated Id-elimination the desired completion $f_2: \underline{A}^\pi \rightarrow \underline{A}^{\iota_2}$. Motivated by this, we define the sub-operad $P \subset [\underline{A}, \underline{A}]$ to have as its operations of shape π , those diagrams of the form (4) in which each f_i and g_i commutes with the A -pointings just defined. Again, it is intuitively clear that this defines a sub-operad—which is to say that the operations with this property are closed under identities and substitution—but to prove this requires a further excursion into the theory of monoidal globular categories: one which for the purposes of the present section, we omit. However, we claim further that P is both normalised and contractible. This will then prove our main result, since the globular context \underline{A} is a P -algebra—as witnessed by the map of globular operads $P \hookrightarrow [\underline{A}, \underline{A}]$ —so that we will have shown the globular context \underline{A} to be an algebra for a normalised, contractible, globular operad P , and hence a weak ω -category.

Now, to show P normalised is trivial, since its operations of shape \star are those context morphisms $A \rightarrow A$ which commute with the pointing $\text{id}_A: A \rightarrow A$, and there is of course only one such. On the other hand, we see that it is contractible through a generalisation of the argument given in the example above. The only part requiring some thought is how to describe generically the process of repeatedly applying Id-elimination. The key to doing this is to prove by induction on π that each of the pointings $r_\pi: A \rightarrow \underline{A}^\pi$ is an injective equivalence in the sense defined in Section 2.2. The injective equivalence structure now encodes the process of repeated Id-elimination. Using this, we may show P contractible as follows. Suppose we are given a diagram like (5) where each f_i and g_i commutes with the A -pointings. We let $B\underline{A}^{\iota_n}$ denote the context obtained from \underline{A}^{ι_n} by removing its final variable, and let $p: \underline{A}^{\iota_n} \rightarrow B\underline{A}^{\iota_n}$ denote the corresponding dependent projection. Then we have a commutative square

$$\begin{array}{ccc} A & \xrightarrow{r_{\iota_n}} & \underline{A}^{\iota_n} \\ r_\pi \downarrow & & \downarrow p \\ \underline{A}^\pi & \xrightarrow{\quad} & B\underline{A}^{\iota_n} \end{array} ,$$

where the lower arrow is obtained by applying first the projection $\underline{A}^\pi \rightarrow B\underline{A}^\pi$, and then the maps f_{n-1} and g_{n-1} . Commutativity obtains by virtue of the fact that f_{n-1} and g_{n-1} commute with the pointings; and so, because r_π is

an injective equivalence and p a dependent projection, we can find a diagonal filler, which will be the required map $f_n: \underline{A}^\pi \rightarrow \underline{A}^{\iota_n}$.

3.2. An axiomatic framework. We now wish to make rigorous the above proof; and as we have already mentioned, we shall do so not in an explicitly type-theoretic manner, but rather within an axiomatic categorical framework. In this Section, we describe this framework and give the intended type-theoretic interpretation.

3.2.1. Definition. A category \mathcal{C} is an *identity type category* if it comes equipped with two classes of maps $\mathcal{I}, \mathcal{P} \subset \text{mor } \mathcal{C}$ satisfying the following axioms:

Empty: \mathcal{C} has a terminal object 1 , and for all $A \in \mathcal{C}$, the unique map $A \rightarrow 1$ is a \mathcal{P} -map.

Composition: The classes of \mathcal{P} -maps and \mathcal{I} -maps contain the identities and are closed under composition.

Stability: Pullbacks of \mathcal{P} -maps along arbitrary maps exist, and are again \mathcal{P} -maps.

Frobenius: The pullback of an \mathcal{I} -map along a \mathcal{P} -map is an \mathcal{I} -map.

Orthogonality: For every commutative square

$$(8) \quad \begin{array}{ccc} A & \xrightarrow{f} & C \\ i \downarrow & & \downarrow p \\ B & \xrightarrow{g} & D \end{array}$$

with $i \in \mathcal{I}$ and $p \in \mathcal{P}$, we can find a diagonal filler $j: B \rightarrow C$ such that $ji = f$ and $pj = g$.

Identities: For every \mathcal{P} -map $p: C \rightarrow D$, the diagonal map $\Delta: C \rightarrow C \times_D C$ has a factorisation

$$\Delta = C \xrightarrow{r} \text{Id}(C) \xrightarrow{e} C \times_D C$$

where $r \in \mathcal{I}$ and $e \in \mathcal{P}$.

We make two remarks concerning this definition. Firstly, by (Empty) and (Stability), any identity type category will have finite products, and product projections will be \mathcal{P} -maps. Secondly, in order to verify (Orthogonality), it suffices, by (Stability), to do so only in those cases where the map along the bottom of (8) is an identity.

3.2.2. Proposition. *Let \mathbb{T} be a dependent type theory admitting each of the inference rules described in Section 2.2. Then the classifying category $\mathcal{C}_{\mathbb{T}}$ is an identity type category, where we take \mathcal{P} to be the class of dependent projections and \mathcal{I} the class of injective equivalences.*

Proof. The empty context $()$ provides a terminal object of $\mathcal{C}_{\mathbb{T}}$. (Composition) is immediate from the definitions. (Stability) corresponds to the possibility of performing type-theoretic substitution. (Frobenius) is shown to hold in [7, Proposition 14]; it is a categorical correlate of the fact that we allow an extra contextual parameter Δ in the statement of the Id-elimination

rule. (Orthogonality) holds by the very definition of injective equivalence, together with the remark made above. Finally, (Identities) says something more than that identity types exist—it says that identity *contexts* exist: which is to say that, for every dependent context $(\Delta) \Gamma \text{ ctxt}$, we may find a context $(\Delta, x, y \in \Gamma) \text{ Id}_\Gamma(x, y) \text{ ctxt}$ such that the contextual analogues of the identity type rules are validated. That this is possible is proven in [9, Proposition 3.3.1]. \square

We will also require two stability properties of identity type categories.

3.2.3. Proposition. *Let \mathcal{C} be an identity type category, and $X \in \mathcal{C}$. Then the coslice category X/\mathcal{C} is also an identity type category, where we take the class of \mathcal{I} -maps (respectively, \mathcal{P} -maps) to consist of those morphisms which become \mathcal{I} -maps (respectively, \mathcal{P} -maps) upon application of the forgetful functor $X/\mathcal{C} \rightarrow \mathcal{C}$.*

3.2.4. Proposition. *Let \mathcal{C} be an identity type category, and $X \in \mathcal{C}$. Then the category \mathcal{C}_X , whose objects are \mathcal{P} -maps $A \rightarrow X$ and whose morphisms are commutative triangles, is also an identity type category, where we define the classes of \mathcal{I} -maps and \mathcal{P} -maps in a manner analogous to that of the previous Proposition.*

The proofs are trivial; the only point of note is that, in the second instance, we could not take \mathcal{C}_X to be the full slice category \mathcal{C}/X , as then (Empty) would not be satisfied.

3.3. Internal weak ω -groupoids. In this section, we describe the notion of weak ω -groupoid internal to an identity type category \mathcal{C} . We begin by defining internal \mathcal{P} -algebras for a globular operad \mathcal{P} .

3.3.1. Definition. A *pre-globular context* in \mathcal{C} is a diagram

$$\Gamma_0 \begin{array}{c} \xleftarrow{s} \\ \xrightarrow{t} \end{array} \Gamma_1 \begin{array}{c} \xleftarrow{s} \\ \xrightarrow{t} \end{array} \Gamma_2 \begin{array}{c} \xleftarrow{s} \\ \xrightarrow{t} \end{array} \Gamma_3 \begin{array}{c} \xleftarrow{s} \\ \xrightarrow{t} \end{array} \cdots$$

satisfying the globularity equations $ss = st$ and $ts = tt$. A pre-globular context is a *globular context* if, for each $n \geq 1$, the map

$$(9) \quad (s, t): \Gamma_n \rightarrow B_n \Gamma$$

is a \mathcal{P} -map, where $B_n \Gamma$ is defined as follows. We have $B_1 \Gamma := \Gamma_0 \times \Gamma_0$, and have $B_{n+1} \Gamma$ given by the pullback


$$(10) \quad \begin{array}{ccc} B_{n+1} \Gamma & \longrightarrow & \Gamma_n \\ \downarrow & & \downarrow (s,t) \\ \Gamma_n & \xrightarrow{(s,t)} & B_n \Gamma \end{array} .$$

Observe that requiring (9) to be a \mathcal{P} -map for $n = 1$ ensures the existence of the pullback (10) defining $B_2 \Gamma$; which in turn allows us to require that (9) should be a \mathcal{P} -map for $n = 2$, and so on. Once again, we have a coinductive characterisation of globular contexts: to give a globular context $\Gamma \in \mathcal{C}$ is to give an object Γ_0 together with a globular context $\Gamma_{+1} \in \mathcal{C}_{\Gamma_0 \times \Gamma_0}$.

The first step in defining \mathcal{P} -algebra structure on a globular context Γ is to describe the object Γ^π of “ π -indexed elements of Γ ”.

3.3.2. Definition. Let Γ be a globular context in \mathcal{C} , and let $\pi \in (T1)_n$. We define the object $\Gamma^\pi \in \mathcal{C}$ by the following induction:


- If $\pi = \star$ then $\Gamma^\pi := \Gamma_0$.
- If $\pi = (\pi_1, \dots, \pi_k)$, then we first form the objects $(\Gamma_{+1})^{\pi_1}, \dots, (\Gamma_{+1})^{\pi_k}$ of $\mathcal{C}_{\Gamma_0 \times \Gamma_0}$. This yields a diagram



in \mathcal{C} . Note that each s and t is a \mathcal{P} -map so that this diagram has a limit, which we define to be Γ^π .


We define maps $\sigma, \tau: \Gamma^\pi \rightarrow \Gamma^{\partial\pi}$ by a further induction:

- For $\pi \in (T1)_1$, we have Γ^π given by the limit of a diagram



and so we may take $\sigma, \tau: \Gamma^\pi \rightarrow \Gamma^\star = \Gamma_0$ to be given by the projections from this limit into the leftmost, respectively rightmost, copy of Γ_0 .

- Otherwise, given $\pi = (\pi_1, \dots, \pi_k)$, we first construct the morphisms $\sigma, \tau: (\Gamma_{+1})^{\pi_i} \rightarrow (\Gamma_{+1})^{\partial\pi_i}$. These give rise to a diagram



and correspondingly for τ . We now take $\sigma, \tau: \Gamma^\pi \rightarrow \Gamma^{\partial\pi}$ to be the induced maps from the limit of the upper subdiagram (which is Γ^π) to the limit of the lower one (which is $\Gamma^{\partial\pi}$).

3.3.3. Proposition. *Let $\Gamma \in \mathcal{C}$ be a globular context. Then there is a globular operad $[\Gamma, \Gamma]$ whose set of operations of shape π comprises all serially commutative diagrams of the form (4).*

We will prove this Proposition using Michael Batanin’s theory of *monoidal globular categories* [2]. The notion of monoidal globular category bears the same relationship to that of strict ω -category as the notion of monoidal category does to that of monoid; in both cases, the former notion is obtained from the latter by replacing everywhere sets with categories, functions with functors, and equalities with coherent natural isomorphisms.

3.3.4. Definition. A *monoidal globular category* \mathcal{E} is given by a sequence of categories and functors

$$\mathcal{E}_0 \begin{array}{c} \xleftarrow{S} \\ \xrightarrow{T} \end{array} \mathcal{E}_1 \begin{array}{c} \xleftarrow{S} \\ \xrightarrow{T} \end{array} \mathcal{E}_2 \begin{array}{c} \xleftarrow{S} \\ \xrightarrow{T} \end{array} \mathcal{E}_3 \begin{array}{c} \xleftarrow{S} \\ \xrightarrow{T} \end{array} \cdots$$

satisfying the globularity equations $SS = ST$ and $TS = TT$, together with, for each natural number n , an identities functor

$$Z: \mathcal{E}_n \rightarrow \mathcal{E}_{n+1}$$

and for each pair of natural numbers $0 \leq k < n$, a composition functor

$$\otimes_k: \mathcal{E}_n \times_k \mathcal{E}_n \rightarrow \mathcal{E}_n$$

where $\mathcal{E}_n \times_k \mathcal{E}_n$ denotes the pullback

$$\begin{array}{ccc} \mathcal{E}_n \times_k \mathcal{E}_n & \longrightarrow & \mathcal{E}_n \\ \downarrow & & \downarrow S^{n-k} \\ \mathcal{E}_n & \xrightarrow{T^{n-k}} & \mathcal{E}_k \end{array}$$

In addition, there are given invertible natural transformations witnessing:

- Associativity:

$$\alpha_{n,k}: A \otimes_k (B \otimes_k C) \cong (A \otimes_k B) \otimes_k C$$

- Unitality:

$$\lambda_n: Z^{n-k} T^{n-k} A \otimes_k A \cong A \quad \text{and} \quad \rho_n: A \otimes_k Z^{n-k} S^{n-k} A \cong A$$

- Interchange:

$$\chi_{n,k,l}: (A \otimes_k B) \otimes_l (C \otimes_k D) \cong (A \otimes_l C) \otimes_k (B \otimes_l D) \quad (\text{for } k < l).$$

These data are required to satisfy a number of coherence axioms, which the reader may find in [2, Definition 2.3].

Just as monoidal categories provide a general environment within which we can speak of monoids, so monoidal globular categories provide a general environment within which we can speak of algebras for a globular operad. The underlying data for an algebra in this general setting is given as follows:

3.3.5. Definition. A *globular object* X in a monoidal globular category \mathcal{E} is given by a sequence of objects $X_i \in \mathcal{E}_i$, one for each natural number i , such that $S(X_{i+1}) = T(X_{i+1}) = X_i$ for all i .

To describe the additional structure required to make a globular object into a P -algebra, we employ one of the central constructions of [2]. This associates to each globular object $X \in \mathcal{E}$ an endomorphism operad $[X, X]$; which allows us to define a P -algebra in \mathcal{E} to be a globular object X together with a globular operad morphism $P \rightarrow [X, X]$. We now describe the construction of $[X, X]$. First observe that if \mathcal{E} is a monoidal globular category, then so too is \mathcal{E}_{+1} , where $(\mathcal{E}_{+1})_n = \mathcal{E}_{n+1}$ and the remaining data is defined in the obvious way. Moreover, if X is a globular object in \mathcal{E} , then X_{+1} is a globular object in \mathcal{E}_{+1} , where again we define $(X_{+1})_n = X_{n+1}$. Now, given a globular object

$X \in \mathcal{E}$ and a pasting diagram $\pi \in (T1)_n$, we define by induction on π an object $X^{\otimes\pi} \in \mathcal{E}_n$:

- If $\pi = \star$, then $X^{\otimes\pi} := X_0 \in \mathcal{E}_0$;
- If $\pi = (\pi_1, \dots, \pi_k)$, then $X^{\otimes\pi} := (X_{+1})^{\otimes\pi_1} \otimes_0 \dots \otimes_0 (X_{+1})^{\otimes\pi_k}$.

3.3.6. Proposition. *Let \mathcal{E} be a monoidal globular category and $X \in \mathcal{E}$ a globular object. Then there is a globular operad $[X, X]$ with*

$$[X, X]_\pi := \mathcal{E}_n(X^{\otimes\pi}, X_n) \quad \text{for all } \pi \in (T1)_n.$$


Proof. This is Proposition 7.2 of [2]. □

We now use this result to prove Proposition 3.3.3. The first step is to construct, from our identity type category \mathcal{C} , a monoidal globular category $\mathcal{E}(\mathcal{C})$.


3.3.7. Definition. Let \mathbb{G} denote the category

$$0 \xleftarrow[\tau]{\sigma} 1 \xleftarrow[\tau]{\sigma} 2 \xleftarrow[\tau]{\sigma} \dots$$


The *generic n -span* \mathbb{S}_n is defined to be the coslice category n/\mathbb{G} . In low dimensions, we have that:



The monoidal globular category $\mathcal{E}(\mathcal{C})$ is defined by taking $\mathcal{E}(\mathcal{C})_n$ to be the full subcategory of the functor category $\mathcal{C}^{\mathbb{S}_n}$ on those functors which send every morphism of \mathbb{S}_n to a \mathcal{P} -map. The remaining structure of $\mathcal{E}(\mathcal{C})$ may be found described in [2, Definition 3.2]. As a representative sample, we describe on objects the functor $Z: \mathcal{E}(\mathcal{C})_1 \rightarrow \mathcal{E}(\mathcal{C})_2$, which is given by



and the functor $\otimes_0: \mathcal{E}(\mathcal{C})_2 \times_0 \mathcal{E}(\mathcal{C})_2 \rightarrow \mathcal{E}(\mathcal{C})_2$, which sends the object



of $\mathcal{E}(\mathcal{C})_2 \times_0 \mathcal{E}(\mathcal{C})_2$ to the object

$$\begin{array}{ccc}
 & H \times_B K & \\
 m \times_B u \swarrow & & \searrow n \times_B v \\
 D \times_B F & & E \times_B G \\
 f \pi_1 \downarrow & q \pi_2 \swarrow & \searrow h \pi_1 \\
 A & & C \\
 & & t \pi_2 \downarrow
 \end{array}$$

of $\mathcal{E}(\mathcal{C})_2$. Note that the requisite pullbacks exist by virtue of the requirement that every arrow in the above diagrams should be a \mathcal{P} -map.

We next observe that, if Γ is a globular context in \mathcal{C} , then there is an associated globular object $X_\Gamma \in \mathcal{E}(\mathcal{C})$ where $(X_\Gamma)_n$ is the n -span

$$\begin{array}{ccc}
 & \Gamma_n & \\
 s \swarrow & & \searrow t \\
 \Gamma_{n-1} & & \Gamma_{n-1} \\
 s \downarrow & t \swarrow & \searrow s \\
 \vdots & & \vdots \\
 s \downarrow & t \swarrow & \searrow s \\
 \Gamma_0 & & \Gamma_0
 \end{array}
 .$$

By a straightforward induction on π , we may now prove that for any $\pi \in (T1)_n$, $(X_\Gamma)^{\otimes \pi} \in \mathcal{E}_n$ is given by the n -span

$$\begin{array}{ccc}
 & \Gamma^\pi & \\
 \sigma \swarrow & & \searrow \tau \\
 \Gamma^{\partial \pi} & & \Gamma^{\partial \pi} \\
 \sigma \downarrow & \tau \swarrow & \searrow \sigma \\
 \vdots & & \vdots \\
 \sigma \downarrow & \tau \swarrow & \searrow \sigma \\
 \Gamma^\star & & \Gamma^\star
 \end{array}
 ;$$

from which it follows that the hom-set $\mathcal{E}(\mathcal{C})_n((X_\Gamma)^{\otimes \pi}, (X_\Gamma)_n)$ is precisely the set of commutative diagrams of the form (4). This allows us to complete the proof of Proposition 3.3.3: indeed, we may take the globular operad $[\Gamma, \Gamma]$ whose existence is asserted there to be the globular operad $[X_\Gamma, X_\Gamma]$ whose existence is assured by Proposition 3.3.6.

3.3.8. Definition. Let \mathcal{C} be an identity type category. An *internal P -algebra* for a globular operad P is a pair (Γ, f) , where Γ is a globular context in \mathcal{C} and $f: P \rightarrow [\Gamma, \Gamma]$ a map of globular operads. By a *weak ω -category* in \mathcal{C} , we

mean a triple (P, Γ, f) , where P is a normalised, contractible globular operad and (Γ, f) an internal algebra for it.

It remains to extend this definition to one of weak ω -groupoid in \mathcal{C} . To do this, we exploit the characterisation of weak ω -groupoids given by Proposition 2.1.5.

3.3.9. Definition. Let $f: P \rightarrow [\Gamma, \Gamma]$ be a weak ω -category in the identity type category \mathcal{C} . Now a *choice of duals* for Γ , with respect to some system of compositions (i_n, m_n) on P , is given by maps

$$\begin{aligned} (-)^*: \Gamma_n &\rightarrow \Gamma_n \\ \eta: \Gamma_n &\rightarrow \Gamma_{n+1} \\ \epsilon: \Gamma_n &\rightarrow \Gamma_{n+1} \end{aligned}$$

for each $n \geq 1$, making the following diagrams commute:

$$(11) \quad \begin{array}{ccc} \Gamma_n & \xrightarrow{(-)^*} & \Gamma_n \\ & \searrow (s,t) & \swarrow (t,s) \\ & \Gamma_{n-1} \times \Gamma_{n-1} & \end{array}$$

$$\begin{array}{ccc} \Gamma_n & \xrightarrow{\eta} & \Gamma_{n+1} \\ s \downarrow & & \downarrow s \\ \Gamma_{n-1} & \xrightarrow{[i_n]} & \Gamma_n \end{array} \quad \begin{array}{ccc} \Gamma_n & \xrightarrow{\eta} & \Gamma_{n+1} \\ ((-)^*, \text{id}) \downarrow & & \downarrow t \\ \Gamma_n & \xrightarrow{s \times t} & \Gamma_n \\ \text{id} \downarrow & & \downarrow s \\ \Gamma_{n-1} & \xrightarrow{[i_n]} & \Gamma_n \end{array}$$

$$\begin{array}{ccc} \Gamma_n & \xrightarrow{\epsilon} & \Gamma_{n+1} \\ t \downarrow & & \downarrow t \\ \Gamma_{n-1} & \xrightarrow{[i_n]} & \Gamma_n \end{array} \quad \begin{array}{ccc} \Gamma_n & \xrightarrow{\eta} & \Gamma_{n+1} \\ (\text{id}, (-)^*) \downarrow & & \downarrow s \\ \Gamma_n & \xrightarrow{s \times t} & \Gamma_n \\ \text{id} \downarrow & & \downarrow s \\ \Gamma_{n-1} & \xrightarrow{[i_n]} & \Gamma_n \end{array} .$$

We say that (Γ, f) is a *weak ω -groupoid* if it has a choice of duals with respect to every system of compositions on P .

3.4. Types are weak ω -groupoids. We are now ready to prove our main theorem. It will follow from a general result that shows a particular class of globular contexts to admit a weak ω -groupoid structure.


3.4.1. Definition. Let \mathcal{C} be an identity type category. A globular context Γ is said to be *reflexive* if it comes equipped with morphisms

$$\Gamma_0 \begin{array}{c} \xleftarrow[r_0]{s} \\ \xleftarrow[t]{t} \end{array} \Gamma_1 \begin{array}{c} \xleftarrow[r_1]{s} \\ \xleftarrow[t]{t} \end{array} \Gamma_2 \begin{array}{c} \xleftarrow[r_2]{s} \\ \xleftarrow[t]{t} \end{array} \cdots$$


where each r_i is an \mathcal{I} -map satisfying $sr_i = tr_i = \text{id}_{\Gamma_i}$.

3.4.2. Theorem. *Every reflexive globular context (Γ, r_i) admits a structure of weak ω -groupoid.*


and the inductive hypothesis, we see that $(\Gamma_*)^\pi$ is given by the unique map $\Gamma_0 \rightarrow \Gamma^\pi$ induced by the following cone:



Thus, it suffices to show that this cone coincides with (13); which is to show that, for each $1 \leq i \leq k$, we have $r_{\pi_i} = r'_{\pi_i} \circ r_0$. Now, observe that r_{π_i} is obtained as $((\Gamma_*)_{+1})^{\pi_i}$, where $(\Gamma_*)_{+1}$ is the globular context



in $\Gamma_0/\mathcal{C}_{\Gamma_0 \times \Gamma_0}$. On the other hand, by a further application of the inductive hypothesis, r'_{π_i} is obtained as the map $((\Gamma_{+1})_*)^{\pi_i}$, where $(\Gamma_{+1})_*$ is the globular context



in $\Gamma_1/\mathcal{C}_{\Gamma_0 \times \Gamma_0}$. But the functor $(r_0)! : \Gamma_1/\mathcal{C}_{\Gamma_0 \times \Gamma_0} \rightarrow \Gamma_0/\mathcal{C}_{\Gamma_0 \times \Gamma_0}$ given by pre-composition with the map $r_0 : \Gamma_0 \rightarrow \Gamma_1$ of $\mathcal{C}_{\Gamma_0 \times \Gamma_0}$ sends the latter of these globular contexts to the former; and thus, because $(r_0)!$ preserves limits, it must also send $((\Gamma_{+1})_*)^{\pi_i}$ to $((\Gamma_*)_{+1})^{\pi_i}$: which is to say that $r_{\pi_i} = r'_{\pi_i} \circ r_0$ as required. \square

Thus, for a reflexive globular context (Γ, r_i) , we have now defined the suboperad $P \subset [\Gamma, \Gamma]$ required for the proof of Theorem 3.4.2. It remains only to show that P is normalised and contractible. To do this, we will need:

3.4.5. Proposition. *Let (Γ, r_i) be a reflexive globular context in \mathcal{C} . Then the maps $r_\pi : \Gamma_0 \rightarrow \Gamma^\pi$ of Definition 3.4.3 are all \mathcal{I} -maps.*

Proof. We proceed by induction on π . When $\pi = \star$, we have r_π an identity map, and hence an \mathcal{I} -map. So suppose now that $\pi = (\pi_1, \dots, \pi_k)$. We consider the diagram (13) defining the map $r_\pi : \Gamma_0 \rightarrow \Gamma^\pi$. In it, each of the maps r'_{π_i} is an \mathcal{I} -map by induction, and so because r_0 is an \mathcal{I} -map by assumption, and \mathcal{I} -maps are closed under composition, $r'_{\pi_i} \circ r_0$ is also an \mathcal{I} -map. Repeated application of the following lemma now completes the proof. \square

3.4.6. **Lemma.** *Suppose that*

$$\begin{array}{ccc}
 & A & \\
 i \swarrow & & \searrow j \\
 B & \text{id}_A & C \\
 p \searrow & \downarrow & \swarrow q \\
 & A &
 \end{array}$$

is a commutative diagram in an identity type category \mathcal{C} . Suppose further that i and j are \mathcal{I} -maps, and p and q are \mathcal{P} -maps. Then the induced map $(i, j): A \rightarrow B \times_A C$ is also an \mathcal{I} -map.

Proof. We first form the pullback square

$$\begin{array}{ccc}
 B \times_A C & \xrightarrow{q'} & B \\
 p' \downarrow & & \downarrow p \\
 C & \xrightarrow{q} & A
 \end{array} .$$

Now the universal property of this pullback induces a factorisation of the commutative square

$$\begin{array}{ccc}
 B & \xrightarrow{\text{id}_B} & B \\
 jp \downarrow & & \downarrow p \\
 C & \xrightarrow{q} & A
 \end{array}$$

as

$$\begin{array}{ccccc}
 B & \xrightarrow{j'} & B \times_A C & \xrightarrow{q'} & B \\
 p \downarrow & & \downarrow p' & & \downarrow p \\
 A & \xrightarrow{j} & C & \xrightarrow{q} & A
 \end{array} .$$

Since the outer rectangle has identities along both horizontal edges, it is a pullback. But the right-hand square is a pullback, and so we deduce that the left-hand square is too. Now p' is a \mathcal{P} -map by (Stability), and j is an \mathcal{I} -map by assumption, and so by (Frobenius), j' is also an \mathcal{I} -map. It follows, by (Composition) and the fact that i is an \mathcal{I} -map, that

$$A \xrightarrow{i} B \xrightarrow{j'} B \times_A C$$

is also an \mathcal{I} -map. But this map is the induced map $(i, j): A \rightarrow B \times_A C$, since it has i as its projection onto B , and $jp'i = j$ as its projection onto C . \square

3.4.7. **Proposition.** *Let (Γ, r_i) be a reflexive globular context in \mathcal{C} . Then the suboperad $P \subset [\Gamma, \Gamma]$ of Proposition 3.4.4 is both normalised and contractible.*

Proof. Note first that the set P_\star is the set of all morphisms $f_0: \Gamma_0 \rightarrow \Gamma_0$ for which $f_0 \circ \text{id}_{\Gamma_0} = \text{id}_{\Gamma_0}$ and hence a singleton, which proves P is normalised. To

show it contractible, we must show that, given a serially commutative diagram of the form

$$(14) \quad \begin{array}{ccc} \Gamma^\pi & & \Gamma_n \\ \sigma \downarrow \tau & & s \downarrow t \\ \Gamma^{\partial\pi} & \xrightarrow[f_{n-1}]{g_{n-1}} & \Gamma_{n-1} \\ \sigma \downarrow \tau & & s \downarrow t \\ \vdots & & \vdots \\ \Gamma_0 & \xrightarrow[f_0]{g_0} & \Gamma_0 \end{array}$$

wherein each f_i and g_i commutes with the pointings, we can find a map $f_n: \Gamma^\pi \rightarrow \Gamma_n$ completing the diagram (and commuting with the pointings). First we note that the diagram

$$\begin{array}{ccc} \Gamma^\pi & \xrightarrow{f_{n-1} \circ \sigma} & \Gamma_{n-1} \\ g_{n-1} \circ \tau \downarrow & & \downarrow (s,t) \\ \Gamma_{n-1} & \xrightarrow{(s,t)} & B_{n-1}\Gamma \end{array}$$

commutes, as may be seen by postcomposing it with the two projections $B_{n-1}\Gamma \rightrightarrows \Gamma_{n-2}$, and observing the resultant diagrams commutative. Thus we induce a map $k: \Gamma^\pi \rightarrow B_n\Gamma$. We now consider the diagram

$$\begin{array}{ccc} \Gamma_0 & \xrightarrow{r_{n-1} \cdots r_0} & \Gamma_n \\ r_\pi \downarrow & & \downarrow (s,t) \\ \Gamma^\pi & \xrightarrow{k} & B_n\Gamma \end{array} .$$

That this is commutative once again follows from the fact that it is so upon postcomposition with the two projections $B_n\Gamma \rightrightarrows \Gamma_{n-1}$. Moreover, r_π is an \mathcal{I} -map by Proposition 3.4.5, and (s, t) a \mathcal{P} -map by the definition of globular context; so that by (Orthogonality), we can find a map $f_n: \Gamma^\pi \rightarrow \Gamma_n$ making both induced triangles commute. That the lower triangle commutes says that f_n renders the diagram (14) serially commutative; whilst that the upper triangle commutes says that f_n commutes with the pointings. \square

Thus we have shown the operad $P \subset [\Gamma, \Gamma]$ to be normalised and contractible, from which it follows that the inclusion $P \hookrightarrow [\Gamma, \Gamma]$ exhibits Γ as a weak ω -category. It remains to show that this weak ω -category is a weak ω -groupoid.

3.4.8. Proposition. *Let (Γ, r_i) be a reflexive globular context in \mathcal{C} , and let $P \subset [\Gamma, \Gamma]$ be the operad defined above. Then the inclusion $P \hookrightarrow [\Gamma, \Gamma]$ exhibits Γ as a weak ω -groupoid.*

Proof. According to Definition 3.3.9, we must show that for any given system of compositions (i_n, m_n) for P , there is a corresponding choice of duals for Γ . Now, for each $n \geq 1$ we have a commutative diagram

$$\begin{array}{ccc} \Gamma_0 & \xrightarrow{r_{n-1} \cdots r_0} & \Gamma_n \\ r_{n-1} \cdots r_0 \downarrow & & \downarrow (t, s) \\ \Gamma_n & \xrightarrow{(s, t)} & B_n \Gamma \end{array} .$$

The left-hand morphism is an \mathcal{I} -map, and the right-hand one a \mathcal{P} -map; and so by (Orthogonality) we have a diagonal filler $(-)^*: \Gamma_n \rightarrow \Gamma_n$. Commutativity of the lower triangle implies the commutativity of the first diagram in (11). We induce η and ϵ similarly, by considering the commutative squares

$$\begin{array}{ccc} \Gamma_0 & \xrightarrow{r_n r_{n-1} \cdots r_0} & \Gamma_{n+1} \\ r_{n-1} \cdots r_0 \downarrow & & \downarrow (s, t) \\ \Gamma_n & \xrightarrow{([i_n]s, [m_n] \circ (-)^*, \text{id})} & B_{n+1} \Gamma \end{array}$$

$$\begin{array}{ccc} \Gamma_0 & \xrightarrow{r_n r_{n-1} \cdots r_0} & \Gamma_{n+1} \\ r_{n-1} \cdots r_0 \downarrow & & \downarrow (s, t) \\ \Gamma_n & \xrightarrow{([m_n] \circ (\text{id}, (-)^*), [i_n]t)} & B_{n+1} \Gamma \end{array} .$$

Again, commutativity of the lower triangles entails the commutativity of the remaining four diagrams in (11). \square

We have thus shown that every reflexive globular context in an identity type category \mathcal{C} bears a structure of weak ω -groupoid. Note that in giving this proof, we have nowhere used the axiom (Identities). In fact, the only reason we need it is to show that from an object of \mathcal{C} we can construct a reflexive globular context corresponding to its tower of identity types.

3.4.9. Definition. Let \mathcal{C} be an identity type category and $A \in \mathcal{C}$. We define a reflexive globular context $\underline{A} \in \mathcal{C}$ by the following induction. For the base case, we take $A_0 = A$. For the inductive step, suppose we have defined A_0, \dots, A_n . Then we may form the n -dimensional boundary $B_n \underline{A}$ of A , and by induction the map $(s, t): A_n \rightarrow B_n \underline{A}$ is a \mathcal{P} -map. So by (Identities), we may factorise the diagonal morphism $A_n \rightarrow A_n \times_{B_n \underline{A}} A_n$ as

$$A_n \xrightarrow{r_{n+1}} \text{Id}(A_n) \xrightarrow{e_{n+1}} A_n \times_{B_n \underline{A}} A_n ,$$

with r_{n+1} an \mathcal{I} -map and e_{n+1} a \mathcal{P} -map. We now define A_{n+1} to be $\text{Id}(A_n)$, and $s, t: A_{n+1} \rightarrow A_n$ to be the composites of e_{n+1} with the two projection morphisms $A_n \times_{B_n \underline{A}} A_n \rightarrow A_n$. It remains to show that the induced map $(s, t): A_{n+1} \rightarrow B_{n+1} \underline{A}$ is a \mathcal{P} -map. But we recall that $B_{n+1} \underline{A}$ was defined by the pullback diagram (10), so that $B_{n+1} \underline{A} = A_n \times_{B_n \underline{A}} A_n$, and the induced map (s, t) is precisely e_{n+1} , which is, by assumption, a \mathcal{P} -map.

This definition, together with Theorem 3.4.2 now immediately imply:

3.4.10. Theorem. *Let \mathcal{C} be an identity type category and $A \in \mathcal{C}$. Then the globular context \underline{A} is a weak ω -groupoid in \mathcal{C} .*

In particular, taking \mathcal{C} to be the identity type category $\mathcal{C}_{\mathbb{T}}$ associated with some dependent type theory \mathbb{T} , we obtain:

3.4.11. Theorem. *Let \mathbb{T} be a dependent type theory admitting each of the rules described in Section 2.2. Then for each type A of \mathbb{T} , the tower of identity types over A is a weak ω -groupoid.*

REFERENCES

- [1] Steve Awodey and Michael Warren. Homotopy theoretic models of identity types. *Mathematical Proceedings of the Cambridge Philosophical Society*, 146(1):45–55, 2009.
- [2] Michael Batanin. Monoidal globular categories as a natural environment for the theory of weak n -categories. *Advances in Mathematics*, 136(1):39–103, 1998.
- [3] Benno van den Berg. Types as weak ω -groupoids. Workshop on “Identity Types—Topological and Categorical Structure”, Uppsala University, 2006.
- [4] Nicolaas de Bruijn. Telescopic mappings in typed lambda calculus. *Information and Computation*, 91(2):189–204, 1991.
- [5] Aurelio Carboni and Peter Johnstone. Connected limits, familial representability and Artin glueing. *Mathematical Structures in Computer Science*, 5(4):441–459, 1995.
- [6] Eugenia Cheng. An ω -category with all duals is an ω -groupoid. *Applied Categorical Structures*, 15(4):439–453, 2007.
- [7] Nicola Gambino and Richard Garner. The identity type weak factorisation system. *Theoretical Computer Science*, 409:94–109, 2008.
- [8] Richard Garner. A homotopy-theoretic universal property of Leinster’s operad for weak ω -categories. *Mathematical Proceedings of the Cambridge Philosophical Society*, 147(3):615–628, 2009.
- [9] Richard Garner. Two-dimensional models of type theory. *Mathematical Structures in Computer Science*, 19(4):687–736, 2009.
- [10] Alexander Grothendieck. Pursuing stacks, 1983. Letter to D. Quillen.
- [11] Martin Hofmann and Thomas Streicher. The groupoid interpretation of type theory. In *Twenty-five years of constructive type theory (Venice, 1995)*, volume 36 of *Oxford Logic Guides*, pages 83–111. Oxford University Press, 1998.
- [12] Bart Jacobs. *Categorical logic and type theory*, volume 141 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Company, 1999.
- [13] G. Maxwell Kelly. *Basic concepts of enriched category theory*, volume 64 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, 1982.
- [14] Tom Leinster. A survey of definitions of n -category. *Theory and Applications of Categories*, 10:1–70, 2002.
- [15] Tom Leinster. *Higher operads, higher categories*, volume 298 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, 2004.
- [16] Tom Leinster. Operads in higher-dimensional category theory. *Theory and Applications of Categories*, 12(3):73–194, 2004.
- [17] Peter Lumsdaine. Weak ω -categories from intensional type theory. In *Typed Lambda Calculi and Applications (Brasilia, 2009)*, Logical Methods in Computer Science, In press.
- [18] Bengt Nordström, Kent Petersson, and Jan M. Smith. *Programming in Martin-Löf’s Type Theory*, volume 7 of *International Series of Monographs on Computer Science*. Oxford University Press, 1990.
- [19] Andrew M. Pitts. Categorical logic. In S. Abramsky, D. M. Gabbay, and T. S. E. Maibaum, editors, *Handbook of Logic in Computer Science, Volume 5. Algebraic and Logical Structures*, pages 39–128. Oxford University Press, 2000.

- [20] Michael Warren. *Homotopy theoretic aspects of constructive type theory*. PhD thesis, Carnegie Mellon University, 2008.

TECHNISCHE UNIVERSITÄT DARMSTADT, FACHBEREICH MATHEMATIK, SCHLOSSGARTEN-
STRASSE 7, 64289 DARMSTADT, GERMANY

E-mail address: `berg@mathematik.tu-darmstadt.de`

DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS, UNIVERSITY OF
CAMBRIDGE, CAMBRIDGE CB3 0WB, UK

E-mail address: `rhgg2@cam.ac.uk`