

December 4, 2009

INTERMEDIATE LOGICS AND THE DE JONGH PROPERTY

DICK DE JONGH, RINEKE VERBRUGGE, AND ALBERT VISSER

ABSTRACT. We prove that all extensions of Heyting Arithmetic with a logic that has the finite frame property possess the de Jongh property.

Dedicated to Petr Hájek, on the occasion of his 70th Birthday

1. PREFACE

The three authors of this paper have enjoyed Petr Hájek's acquaintance since the late eighties, when a lively community interested in the metamathematics of arithmetic shared ideas and traveled among the beautiful cities of Prague, Moscow, Amsterdam, Utrecht, Siena, Oxford and Manchester. At that time, Petr Hájek and Pavel Pudlák were writing their landmark book *Metamathematics of First-Order Arithmetic* [HP91], which Petr Hájek tried out on a small group of eager graduate students in Siena in the months of February and March 1989.

Since then, Petr Hájek has been a role model to us in many ways. First of all, we have always been impressed by Petr's meticulous and clear use of correct notation, witness all his different types of dots and corners, for example in the Tarskian 'snowing'-snowing lemmas [HP91]. But also as a human being, Petr has been a role model by his example of *living in truth*, even in averse circumstances [Hav89]. The tragic story of the Logic Colloquium 1980, which was planned to be held in Prague and of which Petr Hájek was the driving force, springs to mind [DvDLS82]. Finally, we were moved by Petr's open-mindedness when coming to terms with a situation that turned out to look disconcertingly unlike the 'standard model' ¹.

Therefore, in this paper, we would like to pay homage to Petr Hájek. Unfortunately, we cannot hope to emulate his correct use of dots and corners. Instead, we do our best to provide some pleasing non-standard models and non-classical arithmetics.

2. INTRODUCTION

Consider a theory T in constructive predicate logic. A propositional formula φ is T -valid iff, for all substitutions σ of formulas of the language of T for propositional variables, we have $T \vdash \sigma(\varphi)$. The set of T -valid formulas is the propositional logic of T . We will call this logic Λ_T . The de Jongh property for T is the statement that the propositional logic Λ_T of T is precisely Intuitionistic Propositional Logic, in other words, $\Lambda_T = \text{IPC}$.

2000 *Mathematics Subject Classification.* 03F25, 03F30, 03-02, 03B20, 03F50, 03F40.

Key words and phrases. Intuitionistic Logic, Heyting's Arithmetic.

¹Rineke would also like to take the opportunity to thank the whole Hájek family – Petr, Maria, Maria, Jonáš and Jonáš – for their great hospitality extended to her during her long-time research stays in Prague in 1991, 1992 and 1993. The typical Sunday lunches at the Hájeks' flat are especially fondly remembered; these occasions could move from *svíčková na smetaně* with *knedlíky*, wonderfully cooked by Maria, through lively conversations over coffee, to an impressive private cello concert by the then eight-year old *střední* Jonáš.

The original theorem of de Jongh was that Heyting's Arithmetic HA has the de Jongh property. In the past many generalizations have been proved; we will give an enumeration below. In these generalizations the objective was mostly to prove (or in rare cases disprove) the de Jongh property for extensions of HA with some properties like e.g. Church's Thesis. In this paper we will go in a different direction. The idea is to strengthen the logic from the intuitionistic logic to an intermediate one. Intermediate logics are all those logics of strength between intuitionistic and classical logics. Define the de Jongh property for T with respect to an intermediate logic Λ as the statement that $\Lambda_T = \Lambda$. Our conjecture is the following.

Conjecture 2.1. Let $\text{HA}(\Lambda)$ be the result of extending HA with Λ for all formulas. Then $\Lambda_{\text{HA}(\Lambda)} = \Lambda$. □

In this paper we will prove this conjecture for logics Λ with the *Finite Frame Property*, namely,

there exists a class of finite frames \mathfrak{F} such that Λ is precisely the logic valid on all models on frames in \mathfrak{F} . For this class \mathfrak{F} we then have:

$$\Lambda = \{\varphi \mid \mathfrak{F} \models \varphi\} = \{\varphi \mid \text{for all } \mathcal{M} \text{ on } \mathfrak{F}, \mathcal{M} \models \varphi\}$$

For intermediate logics, as for normal modal logics, the finite frame property Λ is in fact equivalent to the ostensibly weaker finite model property (FMP), which expresses that there is a class of finite models \mathfrak{M} such that Λ is precisely the logic valid on all models in \mathfrak{M} . (See section 4 for more background.)

For logics Λ with the finite frame property, indeed $\text{HA}(\Lambda)$ has the de Jongh property with respect to Λ . As we will discuss in the conclusion (section 7), there seems to be little chance of generalizing the methods of this paper to a more extensive class of intermediate logics.

In our proof we will only employ substitutions of Π_2^0 -sentences. From this it follows by quite general reasoning, that, assuming that our class of frames is recursive, we have a uniform version of the de Jongh property. This means that, for Λ with the finite frame property, there is a single substitution σ^* such that $\text{HA}(\Lambda) \vdash \sigma^*(\varphi)$ iff $\Lambda \vdash \varphi$. Or, in a different formulation, there is an embedding of the Lindenbaum Heyting algebra of Λ into the Lindenbaum Heyting algebra of $\text{HA}(\Lambda)$.

3. A BRIEF HISTORY OF DE JONGH'S THEOREM

The following brief overview of the history of de Jongh's Theorem for propositional logic is adapted from [Vis99].

1969: Dick de Jongh proves in an unpublished paper his original theorem that $\Lambda_{\text{HA}} = \text{IPC}$. He uses substitutions of formulas of a complicated form, namely $\forall x(\alpha(x) \vee \neg \alpha(x))$ with α almost negative. As a reminder, a formula is almost negative if it does not contain \vee , and \exists only in front of an equation between terms (see [Tro73]). In fact he proves a much stronger result, namely that the logic of relative interpretations in HA is Intuitionistic Predicate Logic. See the extended abstract [Jon70]. de Jongh's argument uses an ingenious combination of Kripke models and realizability.

1973: Harvey Friedman in his paper [Fri73] gives another proof of de Jongh's theorem for HA. He provides a *single* substitution σ mapping the propositional variables to Π_2^0 -sentences such that $\text{HA} \vdash \sigma(\varphi) \Leftrightarrow \text{IPC} \vdash \varphi$. Thus,

Friedman shows that IPC is *uniformly complete* for Π_2^0 -substitutions in HA. Friedman employs slash-theoretic methods as introduced by Kleene [Kle62].

1973: Craig Smoryński strengthens and extends de Jongh’s work in a number of respects in his very readable paper [Smo73]. To state his results we need a few definitions. We write $D(\Pi_1)$ for the set of disjunctions of Π_1^0 -sentences, $\text{Prop}(\Sigma_1)$ for propositional combinations of Σ_1^0 -sentences. Let us remind the reader of some relevant principles (see [Tro73] for extensive discussions). MP is *Markov’s Principle* MP:

$$\forall x(A \vee \neg A) \wedge \neg \neg \exists x A \rightarrow \exists x A.$$

RFN_{HA} is the formalized uniform reflection principle for HA, where $\forall y Ay$ is closed:

$$\text{Proof}_{\text{HA}}(x, \ulcorner A\bar{y}\urcorner) \rightarrow Ay$$

$\text{TI}(\prec)$ is the transfinite induction scheme for a primitive recursive well-ordering \prec :

$$\forall x((\forall y \prec x) Ay \rightarrow Ax) \rightarrow \forall y Ay.$$

We have de Jongh’s Theorem for the following theories T :

HA, HA+RFN(HA), HA+TI(\prec), and HA+MP.

For the first three theories we can take the range of our substitutions either Σ_1^0 or $D(\Pi_1)$. For HA+MP we can take the range of our substitutions $\text{Prop}(\Sigma_1)$. Smoryński uses Kripke models in combination with the Gödel-Rosser-Mostowski-Kripke-Myhill theorem to prove his results.

1975: Daniel Leivant in his PhD Thesis [Lei79] shows that the predicate logic of interpretations of predicate logic in HA is precisely intuitionistic predicate logic. Leivant’s method is proof-theoretical. In fact Leivant shows that one can use as interpretation a fixed sequence of Π_2^0 -predicates. Leivant’s results yield another proof of Friedman’s results described above.

1976: de Jongh and Smoryński in their paper [JS76] show de Jongh’s Theorem for HAS. They also show uniform completeness for HAS with respect to a substitution with range among the Π_2^0 -sentences.

1981: Yu.V. Gavrilenko in [Gav81] proves de Jongh’s Theorem for HA+ECT₀, i.e., the theory of provable realizability over HA. The principle ECT₀ is Extended Church’s Thesis (see [Tro73]):

$$\forall x(A \rightarrow \exists y By) \rightarrow \exists u \forall x(A \rightarrow \exists v(Tuxv \wedge B(Uv))),$$

where A is almost negative and u does not occur free in A and B and v not in B . In the formula ECT₀ above, T stands for Kleene’s T -predicate and U for the corresponding result-extracting function [Kle43]. Gavrilenko proves this result as a corollary of the similar result of Smoryński for HA.

1981: As a reminder, the principle DNS stands for ‘double negation shift’ (see [Tro73]):

$$\forall x \neg \neg Ax \rightarrow \neg \neg \forall x Ax$$

Albert Visser in his Ph.D. thesis [Vis81] provides an alternative proof of de Jongh’s theorem for HA, HA+DNS, and HA+ECT₀ for Σ_1^0 -substitutions adapting the method of Solovay’s proof of the arithmetical completeness of Löb’s logic for substitutions in PA [Sol76]. In fact, his proof extends to the same theories extended with appropriate reflection principles or transfinite induction over primitive recursive well-orderings.

- 1985:** In his [Vis85], Albert Visser provides an alternative proof of de Jongh’s uniform completeness theorem employing a single Σ_1^0 -substitution. The proof is verifiable in $\text{HA}+\text{con}(\text{HA})$. Here, $\text{con}(\text{HA})$ formalizes the consistency of Heyting Arithmetic. Note that de Jongh’s theorem *implies* $\text{con}(\text{HA})$, so the result is, in a sense, optimal. Visser’s proof uses the NNIL-algorithm, an algorithm that is used to characterize the admissible rules for Σ_1^0 -substitutions. See also [Vis02].
- 1991:** Jaap van Oosten in his paper [Oos91b] provides a more perspicuous version of de Jongh’s semantical proof of de Jongh’s theorem for (non-relativized) interpretations of predicate logic. Van Oosten uses Beth models and realizability. See also [Oos91a].
- 1996:** Using the methods developed by Visser in [Vis82] and by de Jongh and Visser in [JV96], one can prove uniform completeness with respect to Σ_1^0 -substitutions for $\text{HA}+\text{ECT}_0$, $\text{HA}+\text{ECT}_0+\text{RFN}(\text{HA}+\text{ECT}_0)$, and $\text{HA}+\text{TI}(\prec)+\text{ECT}_0$.

It is well known that the de Jongh property does not hold for $\text{HA} + \text{MP} + \text{ECT}_0$. Consider the formulas χ and ρ , which are defined as follows.

- $\chi := (\neg p \vee \neg q)$,
- $\rho := [(\neg\neg\chi \rightarrow \chi) \rightarrow (\chi \vee \neg\chi)] \rightarrow (\neg\neg\chi \vee \neg\chi)$

Clearly, ρ is not provable in IPC. We use \mathbf{r} for *Kleene realizability*. In his classical paper [Ros53], G.F. Rose showed that: $\exists e \forall \sigma \in \text{sub}_{\text{HA}} \mathbb{N} \models e \mathbf{r} \sigma(\rho)$. Here sub_{HA} is the set of substitutions from the propositional variables to sentences of the arithmetical language. Thus, Rose refuted a conjecture of Kleene that a propositional formula is IPC-provable if all its arithmetical instances are (truly and classically) realizable. Note the remarkable fact that one and the same realizer realizes all instances! Inspecting the proof, one sees that only a small part of classical logic is involved in the verification of realizability: Markov’s Principle. See David McCarthy’s paper [McC91] for a detailed analysis. Thus we obtain:

$$\exists e \forall \sigma \in \text{sub}_{\text{HA}} \text{HA} + \text{MP} \vdash e \mathbf{r} \sigma(\rho).$$

Hence, a fortiori, $\rho \in \Lambda_{\text{HA}+\text{MP}+\text{ECT}_0}$. See [Pli09] for an interesting recent survey of propositional realizability logic.

4. INTERMEDIATE LOGICS AND THE FINITE MODEL PROPERTY

Intermediate logics (also called superintuitionistic logics) are the logics between IPC and CPC, classical propositional logic, i.e. the sets of formulas closed under IPC-deduction and uniform substitution. Most anything one needs to know about these logics can be found in [CZ97]. We will enumerate the best known of these logics and mention a few basic facts, mainly concerning the finite model property. The well-known theorem for normal modal logics that the finite frame property and the finite model property for a model logic L coincide (see e.g. [BdRV02]), applies to intermediate logics as well [CZ97]. This means that every sentence not provable from the logic cannot only be refuted in a finite model of the logic, it can also be refuted in a frame validating the logic. So, it is appropriate to use the terminology that a logic has the finite model property (FMP) if there is a class of finite frames for which it is complete. As in the case of modal logics, not all intermediate logics are complete with respect to a class of frames, and not all those which are complete

for such a class have the FMP. But all except one of the well-known intermediate logics which we will now discuss are known to be complete with respect to a class of finite frames.

LC, Dummett's logic, axiomatized by, e.g. $(\varphi \rightarrow \psi) \vee (\psi \rightarrow \varphi)$, is complete with respect to the finite linear frames.

KC, Jankov's logic (also called the logic of weak decidability, whereas real intuitionists might prefer, if anything, the logic of testability), axiomatized by $\neg\varphi \vee \neg\neg\varphi$, is complete with respect to the finite frames with a unique endpoint.

KP, the logic of Kreisel and Putnam, axiomatized by

$$(\neg\varphi \rightarrow \psi \vee \chi) \rightarrow (\neg\varphi \rightarrow \psi) \vee (\neg\varphi \rightarrow \chi).$$

It is complete with respect to the finite partial orderings satisfying the property: For each u and each set X of points succeeding u , there exists a v accessible from u such that all points of X are accessible from v and every endpoint above v is also above some point from X . KP was the first logic shown to have the *disjunction property*: If $\vdash \varphi \vee \psi$, then $\vdash \varphi$ or $\vdash \psi$, where \vdash stands for provability in the given logic (here KP) [KP57].

T_n , the Gabbay-deJongh logics, which are complete with respect to the finite trees which have splittings of exactly n , i.e. each node has exactly n immediate successors. T_n is axiomatized by

$$\bigwedge_{k \leq n+1} ((\varphi_k \rightarrow \bigvee_{j \neq k} \varphi_j) \rightarrow \bigvee_{j \neq k} \varphi_j) \rightarrow \bigvee_{k \leq n+1} \varphi_k$$

T_1 coincides with LC. T_n -frames for $n > 1$ do not have a first order definition. Moreover, T_n for $n > 1$ is not canonical ([CZ97]). However, it is first-order definable on finite frames: it is characterized by the condition that every point has at most n immediate successors. Finally, T_n for $n > 1$ has the disjunction property [dJG74].

BD $_n$, the depth n logics are complete with respect to the finite partial orderings (but also to the trees, or to the splitting trees) of depth n . BD_1 is classical logic, and is axiomatized by Peirce's Law, $((\varphi \rightarrow \psi) \rightarrow \varphi) \rightarrow \varphi$. An axiomatization of BD_n for $n > 1$ is obtained by iteratedly substituting Peirce's Law into itself, e.g. BD_2 is axiomatized by $((\varphi \rightarrow ((\psi \rightarrow \chi) \rightarrow \psi) \rightarrow \psi)) \rightarrow \varphi \rightarrow \varphi$.

Sc, Scott's logic, axiomatized by $((\neg\neg\varphi \rightarrow \varphi) \rightarrow (\varphi \vee \neg\varphi)) \rightarrow \neg\varphi \vee \neg\neg\varphi$. It is complete with respect to the finite partial orderings satisfying the property: In each generated subframe all the endpoints are connected by an R, R^{-1} -chain containing only points of depth 0 or 1 (or equivalently the finite frames which do not have a p-morphism onto the asymmetric four-element tree of depth 3). This is not equivalent to a first-order definition. Like T_n and KP, Scott's logic also has the disjunction property [KP57, CZ97].

ML, Medvedev's logic of finite problems [Med66] has the finite frame property (it is complete with respect to the finite boolean algebras without their top element), but it has no known axiomatization, is and is not known to be decidable.

It contains Scott's logic and KP and is contained in Jankov's logic. It is not finitely axiomatizable and its infinite axiom systems will need an infinity of atoms [MSS79]. ML coincides with the set of formulas all whose essentially negative substitution instances are provable in KP, or stated in another form, it is the logic of the valid schemata obtained by adding $\neg\neg p \rightarrow p$ for atoms only to KP (due to [Lev69], see also [Cia09]).

The **Propositional Logic of Realizability**. Plisko [Pli09] discusses several variants out of which we choose the logic of the effectively realizable formulas, although for most purposes it makes little difference. The logic has no known axiomatization, it is not known whether it has the finite model property or is decidable. It does have the disjunction property. From our earlier remarks it follows that it contains Scott's logic with $\neg\psi \vee \neg\theta$ substituted for the sole variable in Scott's axiom. Just like ML, the Propositional Logic of Realizability is contained in KC, but it has been shown to neither contain nor be contained in ML.

All logics using only $\rightarrow, \wedge, \neg$ in their axioms have the finite model property, but none but IPC itself have the disjunction property (see [CZ97]). The same is true for logics with axioms using only *NNIL*-formulas (No Nesting of Implications on the Left); in fact, these logics are the same as the ones using only \rightarrow (see [Yan08]).

5. ARITHMETICS AND FINITE FRAMES

In this section, we prove our main result. We follow Smoryński's classical paper [Smo73]. In that paper Smoryński gives two proofs. We will use the second more complicated one, which is more flexible for applications. But first we will sketch in a few lines the first proof which gives us a partial result, and indicate why it does not generalize. The idea (of both proofs) is, given a propositional Kripke-model, to construct a Kripke model for HA on the same frame, and then to exhibit arithmetical sentences with exactly the same forcing behavior as the propositional variables on the propositional model.

First, one should realize that any node k in a model \mathcal{K} is attached to a classical structure \mathcal{M}_k (not necessarily a model of HA [Bus93]). The simple proof now starts with the basic lemma:

Lemma 5.1. *Given a (non-rooted) model of HA (or equivalently a set of models for HA), one can obtain a new model of HA by equipping this non-rooted model with a root to which the standard model of the natural numbers is attached.*

Proof. Assume $\mathcal{K}, \mathcal{K}', r$ as in the statement of the lemma. Assume also that $r \Vdash A(0) \wedge \forall x (A(x) \rightarrow A(x+1))$. It is sufficient to show that $r \Vdash \forall x A(x)$, since the \mathcal{K} -part of the model already satisfies (internal) induction. We have that $r \Vdash A(0) \wedge (A(n) \rightarrow A(n+1))$ for each natural number n . This enables us, by applying external induction, to conclude that the new root forces $A(n)$ for each n , from which $r \Vdash \forall x A(x)$ follows. \square

The simple proof now continues using the fact that IPC is complete with respect to the finite trees that are everywhere splitting (each node has at least two immediate successors). One builds a propositional Kripke-model on the same tree by starting with non-standard models for PA (and thus for HA!) on the endpoints, and




FIGURE 1. Scheme of lemma 5.1: A new model of HA is constructed by equipping a set of models of HA with a new root, and attaching the standard model ω to it.

attaching the standard model to each of the other nodes. By the basic lemma 5.1, using induction on the depth of the nodes, this is a model of HA.

The non-standard models at the endpoints are chosen to be incomparable in the following sense: each of them has its own Σ_1^0 -sentence, which, in PA, contradicts all the others. Then, because one has a finite everywhere splitting tree, each node is completely determined by the endpoints which are accessible from it. This is sufficient to create for each node a sentence true on exactly that node (and its successors). Finally, disjunctions of those sentences will characterize arbitrary upward closed subsets of the model, in particular those determined by the valuations of the propositional variables in the propositional model.

To those intermediate logics that are complete with respect to a set of finite splitting trees, this method will immediately apply. The point is that we do not only get a model of HA but, because of the form of the frame, also of Λ . This is immediately clear for $\varphi(\alpha_1, \dots, \alpha_n)$ if $\varphi(p_1, \dots, p_n)$ is a member of Λ and $\alpha_1, \dots, \alpha_n$ are arithmetic sentences with constants for the elements of the appropriate domains of the models. If in such a case $\alpha_1, \dots, \alpha_n$ have free variables, then the truth value of the universally quantified version of $\varphi(\alpha_1, \dots, \alpha_n)$ depends only on the value of sentences arising by substitution in $\varphi(p_1, \dots, p_n)$, and, since we declared these to be true already, the universally quantified forms have to be true as well. In particular, one obtains in this manner the de Jongh property for HA with respect to the logics T_n (for $n > 1$) and BD_n , but not for KC, KP, and ML. Also not for $T_1 = LC$, because linear frames do not split everywhere.

Especially, the need to characterize each node exactly by the endpoints accessible from it seems essential in this setup of the proof. So, for the general case we will need a different way to proceed. One basic idea remains the same. We want to fit a root to a number of models. Smoryński's second method does this in a much more sophisticated manner than the first, and with this method one can under certain strict conditions have a root with a non-standard model attached to it. To set the stage, we need a sequence of lemmas. We start with a few well-known lemmas on

Kripke models of HA. For definitions of forcing in such Kripke models, see [Smo73] and [TvD88].

Lemma 5.2. *Suppose \mathcal{K} is a Kripke model of HA. Let $A\vec{x}$ be a Σ_1^0 -formula. Then, for any node k and any \vec{d} in D_k , we have $k \Vdash A\vec{d}$ iff $\mathcal{M}_k \models A\vec{d}$.*

Proof. We first prove, by induction on the complexity of A , that, for any Δ_0^0 -formula $A\vec{x}$, and any node k and any \vec{d} in D_k , we have $k \Vdash A\vec{d}$ iff $\mathcal{M}_k \models A\vec{d}$. The proof uses the fact that HA proves the decidability of Δ_1^0 -formulas. We treat the case of restricted universal quantification.

Suppose we have the desired property for $By\vec{x}$ in Δ_0^0 . If $k \Vdash \forall y < e By\vec{d}$, then evidently, for all $d' <_{\mathcal{M}_k} e$, $k \Vdash Bd'\vec{d}$. Hence, by the Induction Hypothesis, for all $d' <_{\mathcal{M}_k} e$, $\mathcal{M}_k \models Bd'\vec{d}$. Hence, $\mathcal{M}_k \models \forall y < e By\vec{d}$.

Conversely, suppose (a) $\mathcal{M}_k \models \forall y < e By\vec{d}$. In order to derive a contradiction, suppose (b) $k \Vdash \exists y < e \neg By\vec{d}$. It follows by the Induction Hypothesis and writing out the negation and restricted existential quantification, that $\mathcal{M}_k \models \exists y < e \neg By\vec{d}$. Contradicting (a). So, canceling (b), we conclude $k \not\Vdash \exists y < e \neg By\vec{d}$. Then, by decidability, $k \Vdash \neg \exists y < e \neg By\vec{d}$. So, by intuitionistic predicate logic, $k \Vdash \forall y < e \neg \neg By\vec{d}$. By decidability again, $k \Vdash \forall y < e By\vec{d}$.

The step from Δ_0^0 to Σ_1^0 is easy. \square

The following lemma is due to Wim Ruitenburg.

Lemma 5.3. *Let \mathcal{K} be a model of HA. Let $A\vec{x}$ be a Π_2^0 -formula. Then, for any node k and for any \vec{d} in D_k , we have: $k \Vdash A\vec{d}$ iff, for all $k' \geq k$, $\mathcal{M}_{k'} \models A\vec{d}$.*

Proof. Suppose $A\vec{x}$ is $\forall \vec{y} S\vec{y}\vec{x}$, where S is Σ_1^0 . We have:

$$\begin{aligned} k \Vdash \forall \vec{y} S\vec{y}\vec{d} &\Leftrightarrow \forall k' \geq k \forall \vec{e} \in D_{k'} k' \Vdash S\vec{e}\vec{d} \\ &\Leftrightarrow \forall k' \geq k \forall \vec{e} \in D_{k'} \mathcal{M}_{k'} \models S\vec{e}\vec{d} \\ &\Leftrightarrow \forall k' \geq k \mathcal{M}_{k'} \models \forall \vec{y} S\vec{y}\vec{d} \end{aligned}$$

This gives us the desired property. \square

Next, we need some basic insights from the theory of interpretations.

Lemma 5.4. *Let T be any RE theory. Then $\mathbf{Q} + \text{con}(T)$ interprets T .*

Here \mathbf{Q} is Robinson's Arithmetic, a very weak arithmetical theory introduced by Tarski, Mostowski and Robinson in their book [TMR53].

This is a general version of a fundamental theorem due to Hilbert & Bernays, which was worked out by Wang. The proof was simplified by Feferman. The methods that lead to the general version are due to among others Solovay, Friedman, and Pudlák. See [Vis09], for context, explanation and references. A very crude explanation of the result is as follows. The Model Existence Lemma tells us that, if a theory T is consistent, there is a model \mathcal{N} of T . A moment's reflection shows that the construction of this model, via the Henkin construction, is syntactical in nature. By fine-tuning the argument, we can see that inside a theory that contains the consistency statement of T we can construct an interpretation of T . Here the heuristic is that interpretations are something like uniformly defined internal models.

Thus, inside any model \mathcal{M} of $\mathbf{Q} + \text{con}(T)$ we can construct a model \mathcal{M}' of T . Moreover, if \mathcal{M} is a model of PA , and if T is an arithmetical theory extending \mathbf{Q} , then we can easily prove, using an argument that is essentially due to Dedekind, that \mathcal{M}' is an *end-extension* of \mathcal{M} modulo a definable embedding. The idea is that inside \mathcal{M} we can define the obvious function that maps the numbers of the theory into the numbers of the internal model \mathcal{M}' by recursion. We can prove by induction that the numbers of \mathcal{M} are mapped to an initial segment of \mathcal{M}' . We formulate this in a lemma. For precise discussions of this and similar lemmas, see [HP91].

Lemma 5.5. *Suppose T is an arithmetical theory extending \mathbf{Q} . Then every model \mathcal{M} of $\text{PA} + \text{con}(T)$ contains an internally definable end-extension (modulo a definable embedding) satisfying T .*

Finally, we formulate Smoryński's fundamental lemma. Consider a non-rooted Kripke model \mathcal{K} of HA and a model \mathcal{M} of PA . Suppose \mathcal{K} is definable in \mathcal{M} . This means that the set of nodes K of \mathcal{K} is definable, that the ordering on the nodes is definable, that there is a formula $\delta(k, d)$ giving the domain elements of the node k , and that there are arithmetical formulas $A_P(k, \vec{d})$ representing $k \Vdash P\vec{d}$. Moreover, we ask that \mathcal{M} verifies basic properties like the fact that the relation between the nodes is a partial ordering. We have:

Lemma 5.6. *There is a definable embedding of \mathcal{M} as an initial segment of the model associated with each node of \mathcal{K} . This embedding is unique in the strong sense that there can be only one definable embedding that commutes with 0 and successor. Thus we can form a rooted model \mathcal{K}^+ by adding \mathcal{M} as a new root to \mathcal{K} . We have: \mathcal{M}^+ is a model of HA .*

The proof of this result is in Smoryński's classical paper [Smo73]. The idea is a simple extension of the idea of the basic lemma 5.1. Given a nonrooted model of HA , we cannot just add an arbitrary non-standard model as the root, since such a model does have induction for its own language but not for the enriched language in which we can also talk about \mathcal{K} . However, if \mathcal{K} is internally definable this problem disappears and we can add the non-standard model as a root. One could say that the non-standard model internally thinks it is standard and that as soon as it can talk about \mathcal{K} the earlier external argument can be internalized.

We are now ready for the main construction, which diverges from Smoryński's proof [Smo73].

Theorem 5.1. *Suppose that Λ is the logic of a given class of finite frames \mathfrak{F} . Let $\text{HA}(\Lambda)$ be the result of extending HA with Λ for all formulas. Then*

$$\Lambda_{\text{HA}(\Lambda)} = \Lambda,$$

i.e. the propositional logic of $\text{HA}(\Lambda)$ is Λ . Our result works both when we consider logics of substitutions of formulas and when we consider the logics of substitutions of sentences.

Proof. Let Λ be the logic of a given class of finite frames \mathfrak{F} . Suppose $\Lambda \not\vdash \varphi$. There is a finite model \mathcal{K} with frame in \mathfrak{F} , such that $\mathcal{K} \not\vdash \varphi$. Let the ordering of our model be \preceq . We can arrange it so that the nodes of \mathcal{K} are $0, \dots, n-1$ and, if $i \preceq j$, then $i \leq j$. We define:

- $\text{incon}^0(\text{PA}) := \perp$, $\text{incon}^{k+1}(\text{PA}) := \text{prov}_{\text{PA}}(\text{incon}^k(\text{PA}))$,

- $C_k := \neg \text{incon}^k(\text{PA}) \wedge \text{incon}^{k+1}(\text{PA})$,
- $\overline{C}_0(x) := \text{incon}(\text{PA})$,
- $\overline{C}_{k+1}(x) := \neg \text{proof}_{\text{PA}}(x, \text{incon}^k(\text{PA})) \wedge \text{incon}^{k+2}(\text{PA})$.

We note that the C_i are mutually exclusive. We define T_i , for $0 \leq i < n$, by $T_i := \text{PA} + C_{n-i}$. By Löb's Theorem and Σ_1^0 -soundness, each of the T_i is consistent. Moreover, again by Löb's Theorem, for $i < n - 1$, we have $T_i \vdash \text{con}(T_{i+1})$. Let \mathcal{N}_0 be any model of T_0 . By Lemma 5.4, we can construct an internal model, say \mathcal{N}_1 of T_1 , in \mathcal{N}_0 . The model \mathcal{N}_1 will be an end-extension of \mathcal{N}_0 . We iterate this construction, obtaining an internal model \mathcal{N}_{i+1} of T_{i+1} in \mathcal{N}_i . Since 'being an internal model of' and 'being an end-extension of' are transitive relations, we find that if $i \leq j \leq n - 1$, then \mathcal{N}_j is an internally defined end-extension of \mathcal{N}_i .

We now construct a Kripke model \mathcal{S} by making \mathcal{M}_k the model associated to the node k and taking over the ordering of \mathcal{K} . By Smoryński's Lemma 5.6, we find that \mathcal{S} is a model of HA. Since the frame of our model is in \mathfrak{F} , we see that \mathcal{S} satisfies $\text{HA}(\Lambda)$.

Consider any proposition p in \mathcal{K} . We want to find an arithmetical formula $\sigma(p)$ such that, for all $k < n$, $\mathcal{S}, k \Vdash \sigma(p)$ iff $\mathcal{K}, k \Vdash p$. A first choice to consider would be $\sigma(p) := \bigvee_{\mathcal{K}, k \Vdash p} C_k$. However, it is easy to see this won't wash. Consider, for example, $i \prec j$ and suppose $\mathcal{S}, i \Vdash \sigma(p)$. Then $\mathcal{S}, i \Vdash C_m$, for some m with $\mathcal{K}, m \Vdash p$. It follows that $\mathcal{N}_i \models C_m$, and hence that $m = i$. By persistence, we find that $\mathcal{S}, j \Vdash C_i$, but then $\mathcal{N}_j \models C_i$. A contradiction.

So this does not work. In the light of Ruitenburg's lemma 5.3, we can diagnose the problem as follows: the sentence $\sigma(p)$ is not *constructively* equivalent to a Π_2^0 -sentence or some other kind of sentence that gives us a transfer from classical satisfaction and constructive forcing. However, our $\sigma(p)$ is *classically* equivalent to a Π_2^0 -sentence, and therein lies the simple solution. Suppose $\{k \mid \mathcal{K}, k \Vdash p\} = \{k_0, \dots, k_{m-1}\}$. We define:

$$\bullet \sigma^*(p) := \forall x_0, \dots, x_{m-1} (\overline{C}_{k_0}(x_0) \vee \dots \vee \overline{C}_{k_{m-1}}(x_{m-1})).$$

It is easy to see that *constructively* $\sigma^*(p)$ is equivalent to a Π_2^0 -sentence and that *classically* $\sigma^*(p)$ is equivalent to $\sigma(p)$.

By Ruitenburg's Lemma 5.3, we find:

$$\mathcal{S}, i \Vdash \sigma^*(p) \Leftrightarrow \mathcal{K}, i \Vdash p.$$

Hence, by induction, we find that, for any ψ :

$$\mathcal{S}, i \Vdash \sigma^*(\psi) \Leftrightarrow \mathcal{K}, i \Vdash \psi.$$

So, $\mathcal{S}, 0 \not\Vdash \sigma^*(\varphi)$. □

Remark 5.2. A remarkable aspect of the above proof is that, where Smoryński used Rosser-style self-reference, it only employs Gödelean self-reference —since it uses Löb's Theorem. □

Remark 5.3. We note that we can employ a fixed series of models in our main argument that can be chosen independently of the finite frames. This uses an argument originally due to Harvey Friedman [Fri78]. We define by Carnap's version of the Fixed Point Lemma a formula $A(x)$ such that:

$$\text{PA} \vdash \forall x (A(x) \leftrightarrow (\text{con}(\text{PA} + A(x+1)) \vee \exists y \leq x \text{proof}_{\text{PA}}(y, \neg A(0)))).$$

Let $T_i := \text{PA} + A(i) + \text{incon}(\text{PA} + A(i))$. Let $C_i := \text{con}(T_{i+1}) \wedge \text{incon}(T_i)$. We easily see that the T_i are consistent and that T_i is equivalent to $\text{PA} + C_i$. Moreover, the C_i

are incompatible. We note that, by Löb's Theorem, $T_i \vdash \text{con}(T_{i+1})$. We now start with a model \mathcal{N}_0 of T_0 in which we construct an internally definable end-extension \mathcal{N}_1 that is a model of T_1 , etc. Thus we obtain a sequence of models \mathcal{N}_i , where each next model is a definable end-extension of the previous one. Moreover, $\mathcal{N}_i \models C_i$. \square

Can we extend the above proof to other arithmetical theories besides HA as basis? We note that we can extend it to weaker theories like $i\text{-}I\Sigma_1$, the intuitionistic version of $I\Sigma_1$, since the construction of an end-extension from a consistency statement already works in the classical theory $I\Sigma_1$ [HP91]. Secondly, the proof also works for stronger theories like HA plus uniform reflection, or HA extended with an RE set of negations that are Π_2^0 -conservative over PA.

We have one somewhat more interesting extension of our result, for intermediate logics with what we will call the endpoint replacement property:

Definition 5.4. Consider an intermediate logic Λ .

- (1) Let \mathfrak{F} be a class of frames for the logic. We say that Λ has the *endpoint replacement property with respect to \mathfrak{F}* if the following holds. Take any \mathcal{F} in \mathfrak{F} , a set of endpoints $\{u_i \mid i \in I\}$ of \mathcal{F} , and a set $\{\mathcal{M}_i \mid i \in I\}$ of (possibly infinite) models of Λ with roots $\{r_i \mid i \in I\}$. Let \mathcal{G} be the frame that is the result of replacing the endpoints $\{u_i \mid i \in I\}$ by the frames of $\{\mathcal{M}_i \mid i \in I\}$. Then any model \mathcal{M} on \mathcal{G} which is such that its submodels generated by $\{r_i \mid i \in I\}$ are exactly the $\{\mathcal{M}_i \mid i \in I\}$ will validate Λ .
- (2) Λ has the *endpoint replacement property* if Λ has the endpoint replacement property with respect to its class of frames.

\square

Of the logics we mentioned, the following have the endpoint replacement property: IPC, KC, LC, and T_n . The logics KP, Sc, ML, and BD_n do not have the property with respect to a class of frames for which they are complete. On the negative side: for KP and Sc this is obvious from the fact that the simple 3-element fork which is a frame for both is no longer a frame for either when attached to an endpoint of itself. On the positive side, we will just show the most complicated case: the T_n .

Proposition 5.5. T_n has the endpoint replacement property. \square

Proof. Let \mathcal{F} be a frame for T_n , let $\{u_i \mid i \in I\}$ be a set of endpoints of \mathcal{F} , and let $\{\mathcal{M}_i \mid i \in I\}$ be a set of models of T_n with roots $\{r_i \mid i \in I\}$. Let \mathcal{M} be a model on the frame \mathcal{G} that is the result of replacing the endpoints $\{u_i \mid i \in I\}$ by the frames of $\{\mathcal{M}_i \mid i \in I\}$ such that the models generated by the r_i are exactly the \mathcal{M}_i . We have to show that the assumption that $T_n(\varphi_1, \dots, \varphi_{n+1})$ is falsified in \mathcal{M} will lead to a contradiction. We can then assume without loss of generality that the root satisfies the antecedent of T_n for $\varphi_1, \dots, \varphi_{n+1}$ and falsifies each of $\varphi_1, \dots, \varphi_{n+1}$. Let us define N as the set of nodes in \mathcal{M} that falsify each of $\varphi_1, \dots, \varphi_{n+1}$. This will be a downward closed set containing the root, falsifying $T_n(\varphi_1, \dots, \varphi_{n+1})$ everywhere. Since for each $k \leq n+1$, $\bigvee_{j \neq k} \varphi_j$ is falsified at each $v \in N$, also $\varphi_k \rightarrow \bigvee_{j \neq k} \varphi_j$ is false at v for some $k \leq n+1$. This means that for each $v \in N$, there is a $k \leq n+1$ and a w_k with $v \leq w_k$ such that w_k makes φ_k true but no φ_j for $j \neq k$. Note that such nodes w_k and w_j for different k and j are always incomparable. Note also that $x \Vdash T_n$ iff $x \in N$.

Let us first consider the case that one of the r_i is in N . That is impossible, since then $r_i \not\vdash \top_n(\varphi_1, \dots, \varphi_{n+1})$ whereas the model \mathcal{M}_i is supposed to be a model for \top_n . The second possibility is that none of the r_i are in N . Note that then, if \mathcal{M}_i contains any w_k -node as described above, its root r_i will have to be a w_k -node, and there cannot be w_j -nodes for any $j \neq k$ in \mathcal{M}_i . If we now restore the endpoints $\{u_i \mid i \in I\}$ in \mathcal{M} in place of the models $\{\mathcal{M}_i \mid i \in I\}$ and define

$$V'(p_m) = \{w \in \mathcal{F} - \{u_i \mid i \in I\} \mid w \Vdash \varphi_m\} \cup \{u_i \mid r_i \Vdash \varphi_m\},$$

then it is obvious that for this new valuation, for each $v \in N$ (and in particular for the root of \mathcal{F}), $v \not\vdash \top_n(p_1, \dots, p_{n+1})$, since one of the $p_k \rightarrow \bigvee_{j \neq k} p_j$ will be falsified at the appropriate w_k . Thus, the resulting model will falsify $\top_n(p_1, \dots, p_{n+1})$. This is again a contradiction, because \mathcal{F} is a frame for \top_n . \square

Theorem 5.6. *Suppose that Λ has both the finite model property and the endpoint replacement property with respect to a class \mathfrak{F} of finite frames. Suppose that U is a consistent extension of $\text{HA}(\Lambda)$ with an RE set of negations N . Then U has the de Jongh property for Λ .*

Before we turn to the proof of our theorem, we remind the reader of some basic facts concerning Rosser sentences. Let V be any consistent RE extension of $i\text{-EA}$, the intuitionistic version of Elementary Arithmetic. Using Craig's trick, we can arrange that the axiom set of V is given by an elementary predicate α . This predicate can be taken to be a $\Delta_1^0(i\text{-EA})$ -formula. This means that both αx and $\neg \alpha x$ are $i\text{-EA}$ -provably equivalent to a Σ_1^0 -formula and that αx is $i\text{-EA}$ -provably decidable. Using this predicate, we can find a reasonable arithmetization proof_V , the proof-predicate for V , such that $\text{proof}_V(x, y)$ is $\Delta_1^0(i\text{-EA})$. Let $\text{prov}_V(x)$ be $\exists y \text{proof}_V(y, x)$.

Consider formulas A and B . Suppose $A = \exists x A_0 x$ and $B = \exists y B_0 y$. We define:

- $A \leq B := \leftrightarrow \exists x (A_0 x \wedge \forall y < x \neg B_0 y)$,
- $A < B := \leftrightarrow \exists x (A_0 x \wedge \forall y \leq x \neg B_0 y)$,
- $(A < B)^\perp := \leftrightarrow B \leq A$,
- $(A \leq B)^\perp := \leftrightarrow B < A$.

Using Gödel's Fixed Point Lemma, we can find a Rosser sentence R for V such that $i\text{-EA} \vdash R \leftrightarrow \text{prov}_V(\neg R) \leq \text{prov}_V(R)$. We clearly have $i\text{-EA} \vdash \neg(R \wedge R^\perp)$. Using Rosser's argument, we can show that both $V + R$ and $V + \neg R$ are consistent. We note that both R and R^\perp can be rewritten over $i\text{-EA}$ plus Σ_1^0 -collection to the strict Σ_1^0 form. Since we work in extensions of HA , we may assume that R and R^\perp are Σ_1^0 .

We turn to the proof of theorem 5.6.

Proof. Let Λ and U be as required for theorem 5.6. Consider any φ such that $\Lambda \not\vdash \varphi$. Let \mathcal{K} be a counter-model in \mathfrak{F} . Suppose the non-endpoints of \mathcal{K} are a_0, \dots, a_{n-1} and that we have chosen our enumeration in such a way that $i \leq j$ implies $a_i \preceq a_j$. We repeat our construction from the proof of theorem 5.1 of a sequence of models with \mathcal{N}_i for $0, \dots, n-1$, with the modification that we take as our base theory $W := \text{PA} + \text{con}(U)$. So \mathcal{N}_{n-1} will satisfy $\text{con}(U)$ and $\text{incon}(\text{PA} + \text{con}(U))$.

We enumerate the endpoints as b_0, \dots, b_{r-1} . We construct a sequence of Σ_1^0 -sentences S_0, \dots, S_{r-1} , with the following properties, for $i < j \leq r-1$:

$$W \vdash \text{con}(U + S_i) \text{ and } U \vdash \neg(S_i \wedge S_j).$$

The easiest way to construct such a sequence is by induction on r . If $r = 1$, we take $S_0 := \top$. Suppose we have constructed S'_0, \dots, S'_{s-1} with the desired properties. Let R and R^\perp be the pair consisting of the Σ_1^0 Rosser sentence for $U + S'_{s-1}$ and its Σ_1^0 opposite. We take $S_i := S'_i$, for $i < r - 1$, $S_{r-1} := (S'_{r-1} \wedge R)$, and $S_r := (S'_{r-1} \wedge R^\perp)$. We now construct in \mathcal{N}_{n-1} inner Kripke models of $U + S_i$. The construction is analogous to the classical case. It is essential that the base theory in which we do the construction is classical! See Appendix A of [Vis98] for a description of how to do an internal Kripke model construction. This gives us models \mathcal{M}_j of $U + S_j$, internally definable in each \mathcal{N}_i .

We build a Kripke model for arithmetic by associating \mathcal{N}_i with a_i and \mathcal{M}_j with b_j . This will be a model of $\text{HA} + N$ by the internal version of Smoryński's Lemma 5.6, noting that the additional negations in N are always downwards preserved. It will be a model of Λ by the endpoint replacement property. We associate to each propositional atom p in \mathcal{K} the sentence:

$$E_p := \bigvee_{a_i \Vdash p} (C_i \wedge \text{con}(U)) \vee \bigvee_{b_j \Vdash p} S_j.$$

As before in the proof of theorem 5.1, we transform E_p to a classically equivalent Π_2^0 -sentence, say E_p^+ . Let σ be the substitution $p \mapsto E_p^+$. By Ruitenburg's Lemma 5.3, we find that our new model does not force $\sigma(\varphi)$. \square

So, for example, for $\Lambda = \text{IPC}, \text{LC}, \text{KC}, \text{T}_n$, we do have the de Jongh property for $\text{HA}(\Lambda)$ plus the negation of the sentence expressing the Primitive Recursive Markov's Principle, and for $\text{HA}(\Lambda)$ plus the negation of the sentence expressing the decidability of the Halting Problem. (See [Tro73] for definitions of these properties.)

6. A UNIFORMIZATION RESULT

In this section we show how to prove uniformization using recursion-theoretic arguments. This style of result is originally due to Franco Montagna [Mon79] and, independently, Albert Visser [Vis81].

Theorem 6.1. *Consider any theory T with elementary axiom set and any interpretation N , such that N interprets $i\text{-EA}$, the constructive version of EA (which is also known as $I\Delta_0 + \text{exp}$), in T . Let A_i be an elementary decidable sequence of sentences in the language of T .*

Suppose, for every i , $T \not\vdash A_i$. Then there is a Σ_1^0 -formula $R(x)$ such that:

- a. $i\text{-EA} \vdash (R(x) \wedge R(y)) \rightarrow x = y$,
- b. for any i , $T \not\vdash R^N(i) \rightarrow A_i$.

Proof. Let $R(x)$ be such that:

$$i\text{-EA} \vdash R(x) \leftrightarrow \exists p (\text{proof}_T(p, R^N(x) \rightarrow A_x) \wedge \forall q \leq p \forall y \leq q \neg \text{proof}_T(q, R^N(y) \rightarrow A_y)).$$

We work with the reasonable assumption that the code of the numeral of y is larger than y , that the code of a formula in which a numeral occurs is larger than the code of that numeral, and that the code of a proof is larger than the code of any formula occurring in it. Also we assume that proofs have single conclusions. (See [HP91] for discussions of such reasonable assumptions.)

We verify the uniqueness clause (a). Reason in i -EA. Suppose $R(x)$ and $R(y)$ with witnesses p and q , respectively. In case $p = q$, we are done. Suppose $q < p$. From $R(x)$, we have: $\forall y' \leq q \neg \text{proof}_T(q, R^N(y') \rightarrow A_{y'})$. Since, by our assumptions on coding, we must have $y < q$, we find $\neg \text{proof}_T(q, R^N(y) \rightarrow A_y)$. This contradicts $R(y)$.

We verify (b). Suppose, in order to derive a contradiction, that $T \vdash R^N(i) \rightarrow A_i$. Let π be a proof that witnesses this fact. Via a finite search among proofs with Gödel number smaller or equal to the Gödel number of π , we may obtain the T -proof π^* with smallest Gödel number with a conclusion of the form $R^N(j) \rightarrow A_j$. Say we have: $\pi^* : T \vdash R^N(i^*) \rightarrow A_{i^*}$. By the definition of R , it follows that $R(i^*)$ is true. So, by Σ_1^0 -completeness, $T \vdash R^N(i^*)$. Hence, $T \vdash A_{i^*}$, contradicting the assumption. We may conclude that $T_i \not\vdash R(i) \rightarrow A_i$. \square

Let Λ be any intermediate logic with the finite frame property. Suppose the set of finite frames \mathfrak{F} corresponding to Λ is decidable. We can easily find an elementary enumeration $(\varphi_k)_{k \in \omega}$ of all Λ -underivable formulas, such that we can find a counter-model \mathcal{K}_k with frame in \mathfrak{F} of φ_k in an elementary way from k . In our main theorem, we have shown that we can transform \mathcal{K}_k in an elementary way into a Π_2^0 -substitution σ_k such that $\text{HA}(\Lambda) \not\vdash \sigma_k(\varphi_k)$.

Theorem 6.2. *Under the circumstances described above, we have the uniform de Jongh property for $\text{HA}(\Lambda)$ with respect to Λ .*

Proof. Let $A_i := \sigma_i(\varphi_i)$. Applying theorem 6.1, with $\text{HA}(\Lambda)$ in the role of T , we find $R(x)$ with the promised properties. Take $\tau(p) := \exists y (R(y) \wedge \text{true}_{\Pi_2^0}(\ulcorner \sigma_y(p) \urcorner))$. Here $\text{true}_{\Pi_2^0}$ is the Π_2^0 -truth predicate and $\ulcorner \sigma_y(p) \urcorner$ is an arithmetization that sends y to the Gödel number of $\sigma_y(p)$.

Suppose $\text{HA}(\Lambda) \vdash \tau(\varphi_j)$. It follows that $\text{HA}(\Lambda) \vdash R(j) \rightarrow \sigma_j(\varphi_j)$, contradicting the assumption. \square

7. CAUTIONARY AFTERWORD

We have proved our results with pleasant, pedestrian methods. The reason for this luxurious situation is the simplicity and power of the two basic ideas on which everything rests: Smoryński's idea of the preservation of HA under adding ω as a root and Smoryński's extension of the idea to non-standard models, provided that the rest of the Kripke model is definable in the new root. These ideas allow us to construct Kripke models step-by-step with a lot of control over their properties.

However, as Johan Cruijff said *elk voordeel heb z'n nadeel*, i.e., *every advantage comes with a disadvantage*. The disadvantage is that the essential dependence on these ideas makes the results not easily extendible.

A first possible direction of extension would be *more frame classes*. We note that there is not much hope to extend our results to frame classes with infinite models—at least not with the methods at hand. The reason is that the preservation of HA under adding a suitable root is essentially a tool for constructing finite objects. Of course, we may extend the construction to *non-standardly finite models*, but these are rather artificial from the point of view of frame classes of intermediate logics. The good news here is that almost all known natural logics do have the finite frame property. A possible exception is the Propositional Logic of Realizability. In this

connection it is worth pointing out that de Jongh's original proof could handle infinite frames but that it did need trees.

The second possible extension would be *more arithmetical theories*. There seems to be no hope to extend our results to extensions of HA that lack the Smoryński property. For example, we do know that $\text{HA} + \text{ECT}_0$ has the de Jongh property of IPC, but the extension of $\text{HA} + \text{ECT}_0$ with intermediate logics is a complete *terra incognita*. In this case it could be interesting to look for *counterexamples* to the extension of our results.

REFERENCES

- [BdRV02] P. Blackburn, M. de Rijke, and Y. Venema. *Modal Logic*, volume 53 of *Cambridge Tracts in Theoretical Computer Science*. Cambridge University Press, 2002.
- [Bus93] S. Buss. Intuitionistic validity in T -normal Kripke structures. *Annals of Pure and Applied Logic*, 59:159–173, 1993.
- [Cia09] I.A. Ciardelli. Inquisitive semantics and intermediate logics. Technical report, ILLC, University of Amsterdam, Amsterdam, 2009.
- [CZ97] A. Chagrov and M. Zakharyashev. *Modal Logic*. Oxford Logic Guides. Oxford University Press, Oxford, 1997.
- [dJG74] D.H.J. de Jongh and D. Gabbay. A sequence of decidable finitely axiomatizable intermediate logics with the disjunction property. *The Journal of Symbolic Logic*, 39:67–78, 1974.
- [DvDLS82] D D. van Dalen, D. Lascar, and T.J. Smiley, editors. *Logic Colloquium '80: Papers Intended for the European Summer Meeting of the Association for Symbolic Logic*, volume 108 of *Studies in Logic and the Foundations of Mathematics*, Amsterdam, 1982. North-Holland.
- [Fri73] H. Friedman. Some applications of Kleene's methods for intuitionistic systems. In A.R.D. Mathias and H. Rogers, editors, *Cambridge Summer @S@chool in Mathematical Logic*, pages 113–170. Springer Verlag, Berlin, Heidelberg, New York, 1973.
- [Fri78] H. Friedman. Classically and intuitionistically provably recursive functions. In G.H. Müller and D.S. Scott, editors, *Higher Set Theory*, pages 21–27. Springer Verlag, Berlin, Heidelberg, New York, 1978.
- [Gav81] Yu. V. Gavrilenko. Recursive realizability from the intuitionistic point of view. *Soviet Mathematical Dokl.*, 23:9–14, 1981.
- [Hav89] V. Havel. *Living in Truth*. Faber and Faber, London, 1989.
- [HHS^T96] W. Hodges, M. Hyland, C. Steinhorn, and J. Truss, editors. *Logic: From Foundations to Applications*. Clarendon Press, Oxford, 1996.
- [HP91] P. Hájek and P. Pudlák. *Metamathematics of First-Order Arithmetic*. Perspectives in Mathematical Logic. Springer, Berlin, 1991.
- [Jon70] D.H.J. de Jongh. The maximality of the intuitionistic predicate calculus with respect to Heyting's Arithmetic. *The Journal of Symbolic Logic*, 36:606, 1970.
- [JS76] D.H.J. de Jongh and C. Smoryński. Kripke models and the intuitionistic theory of species. *Annals of Mathematical Logic*, 9:157–186, 1976.
- [JV96] D.H.J. de Jongh and A. Visser. Embeddings of Heyting algebras. In [?], pages 187–213, 1996.
- [Kle43] S.C. Kleene. Recursive predicates and quantifiers. *Transactions of the AMS*, 53:41–73, 1943.
- [Kle62] S.C. Kleene. Disjunction and existence under implication in elementary intuitionistic formalisms. *Journal of Symbolic Logic*, 27:11–18, 1962.
- [KP57] G. Kreisel and H. Putnam. Eine Unableitbarkeitsbeweismethode für den Intuitionistischen Aussagenkalkül. *Zeitschrift für Mathematische Logik and Grundlagen der Mathematik*, 3:74–78, 1957.
- [Lei79] D. Leivant. *Absoluteness in Intuitionistic Logic*, volume 73. Mathematical Centre Tract, Amsterdam, 1979. The Thesis was originally published in 1975.
- [Lev69] V.A. Levin. Some syntactic theorems on the calculus of finite problems of Yu.T. Medvedev. *Soviet Mathematics Doklady*, 10:288–290, 1969.

- [McC91] D.C. McCarty. Incompleteness in intuitionistic metamathematics. *The Notre Dame Journal of Formal Logic*, 32:323–358, 1991.
- [Med66] Yu. Medvedev. Interpretation of logical formulas by means of finite problems. *Soviet Mathematics Doklady*, 7:180–183, 1966.
- [Mon79] F. Montagna. On the diagonalizable algebra of Peano Arithmetic. *Bollettino Unione Matematica Italiana*, 16-B:795–813, 1979.
- [MSS79] L. Maksimova, V. Shehtman, and D. Skvortsov. The impossibility of a finite axiomatization of Medvedev’s logic of finitary problems. *Soviet Mathematics Doklady*, 20:394–398, 1979.
- [Oos91a] J. van Oosten. *Exercises in Realizability*. Ph. D. Thesis. Department of Mathematics and Computer Science, University of Amsterdam, Amsterdam, 1991.
- [Oos91b] J. van Oosten. A semantical proof of de Jongh’s theorem. *Archive for Mathematical Logic*, 31:105–114, 1991.
- [Pli09] V. E. Plisko. A survey of propositional realizability logic. *the Bulletin of Symbolic Logic*, 15:1–42, 2009.
- [Ros53] G.F. Rose. Propositional calculus and realizability. *Transactions of the American Mathematical Society*, 75:1–19, 1953.
- [Smo73] C. Smoryński. Applications of Kripke models. In A.S. Troelstra, editor, *Metamathematical Investigations of Intuitionistic Arithmetic and Analysis*, Springer Lecture Notes 344, pages 324–391. Springer, Berlin, 1973.
- [Sol76] R.M. Solovay. Provability interpretations of modal logic. *Israel Journal of Mathematics*, 25:287–304, 1976.
- [TMR53] A. Tarski, A. Mostowski, and R.M. Robinson. *Undecidable theories*. North-Holland, Amsterdam, 1953.
- [Tro73] A.S. Troelstra, editor. *Metamathematical Investigations of Intuitionistic Arithmetic and Analysis*. Springer Lecture Notes 344. Springer, Berlin, 1973.
- [TvD88] A.S. Troelstra and D. van Dalen. *Constructivism in Mathematics, vol 2*. Studies in Logic and the Foundations of Mathematics, vol. 123. North Holland, Amsterdam, 1988.
- [Vis81] A. Visser. *Aspects of Diagonalization and Provability*. Ph.D. Thesis, Department of Philosophy, Utrecht University, 1981.
- [Vis82] A. Visser. On the completeness principle. *Annals of Mathematical Logic*, 22:263–295, 1982.
- [Vis85] A. Visser. Evaluation, provably deductive equivalence in Heyting’s Arithmetic of substitution instances of propositional formulas. Logic Group Preprint Series 4, Department of Philosophy, Utrecht University, Heidelberglaan 8, 3584 CS Utrecht, 1985.
- [Vis98] A. Visser. Interpretations over Heyting’s Arithmetic. In E. Orłowska, editor, *Logic at Work*, Studies in Fuzziness and Soft Computing, pages 255–284. Physica-Verlag, Heidelberg/New York, 1998.
- [Vis99] A. Visser. Rules and arithmetics. *The Notre Dame Journal of Formal Logic*, 40(1):116–140, 1999.
- [Vis02] A. Visser. Substitutions of Σ_1^0 -sentences: Explorations between intuitionistic propositional logic and intuitionistic arithmetic. *Annals of Pure and Applied Logic*, 114:227–271, 2002.
- [Vis09] A. Visser. Can we make the Second Incompleteness Theorem coordinate free? *Journal of Logic and Computation*, 2009. doi: 10.1093/logcom/exp048.
- [Yan08] F. Yang. Intuitionistic subframe formulas, nnil-formulas and n-universal models. Technical report, ILLC, University of Amsterdam, Amsterdam, 2008.

ILLC, UNIVERSITY OF AMSTERDAM, ROOM C3.130 B, BUILDING C, SCIENCE PARK 904, 1098
XH AMSTERDAM, THE NETHERLANDS

E-mail address: `d.h.j.dejongh@uva.nl`

ARTIFICIAL INTELLIGENCE, UNIVERSITY OF GRONINGEN, PO BOX 407, 9700 AK GRONINGEN,
THE NETHERLANDS

E-mail address: `rineke@ai.rug.nl`

DEPARTMENT OF PHILOSOPHY, UTRECHT UNIVERSITY, HEIDELBERGLAAN 8, 3584 CS UTRECHT,
THE NETHERLANDS

E-mail address: `Albert.Visser@phil.uu.nl`