

## STABILITY OF A FAKE TOPOLOGICAL HILBERT SPACE

JAN J. DIJKSTRA AND JAN VAN MILL

**ABSTRACT.** The space under consideration is the basic fake Hilbert space  $Y$  of Anderson, Curtis and van Mill. It is shown that the product of an arbitrary space  $A$  with  $Y$  is homeomorphic to  $Y$  if and only if  $A$  is a compact absolute retract. Furthermore, we prove that the complement of  $Y \times Y$  is a capset in  $Q \times Q$ , which implies the known result that  $Y \times Y$  is homeomorphic to Hilbert space.

**1. Introduction.** We are interested in the basic fake Hilbert space  $Y$  that was constructed by Anderson, Curtis and van Mill [1]. The space  $Y$  is the complement of a  $\sigma Z$ -set in the Hilbert cube  $Q$  and, hence, a complete AR. The following properties can be found in [1] and illustrate the closeness of  $Y$  to the Hilbert space  $\ell^2$ : (a)  $Y$  is homogeneous, (b)  $Y \times Y$  is homeomorphic to  $\ell^2$ , and (c)  $Y$  has the weak discrete approximation property. The space has proved to be a very useful basis for the construction of other peculiar spaces and counterexamples as is witnessed by the papers of Anderson et al. [1], Dijkstra and van Mill [8], Dijkstra [7], and Bowers [3]. More information on  $Y$  can be found in Dijkstra [6, Chapters 4 and 5]. The most important results here are the Unknotting Theorem (homeomorphisms between compacta in  $Y$  can be extended with control) and the Negligibility Theorem (the negligible compacta in  $Y$  are precisely the compacta with the shape of a finite set).

In this article we investigate the stability of  $Y$  under multiplication. The result  $Y \times Y \approx \ell^2$  can be improved by showing that the complement of  $Y \times Y$  in  $Q \times Q$  is a capset. We are mainly interested, however, in determining for which spaces  $A$  the product  $Y \times A$  is homeomorphic to  $Y$ . We show that this is the case precisely if  $A$  is a compact absolute retract.

---

1980 Mathematics Subject Classification. Primary 57N20.

*Key words and phrases.* Fake topological Hilbert space, capset, Hilbert cube, absolute retract, continuum-component.

The research of the second author was in part supported by NSF grant DMS-86 06113.

Received by the editors on October 9, 1986.

All spaces in this paper are Tychonoff. Absolute retracts are assumed to be separable metric. For undefined terms from infinite-dimensional topology, see, e.g., Bessaga and Pełczyński [2].

**2. Preliminaries.** In this section we introduce the fake Hilbert space  $Y$  and we state several facts that will be used in the next sections.

If  $\varepsilon > 0$ , then  $Q(\varepsilon)$  denotes the space  $\prod_{i=1}^{\infty} [-\varepsilon, \varepsilon]_i$  equipped with the product topology. The standard representation of the Hilbert cube  $Q$  is  $Q(1) = J^{\mathbf{N}}$ , where  $J = [-1, 1]$ . Let  $\mathcal{R}$  be the set of all sequences  $p_1, p_2, p_3, \dots$  in the interval  $(0, 1)$  such that  $\lim_{i \rightarrow \infty} p_i = 1$  and let  $\mathcal{R}^{\uparrow}$  be the subset consisting of all strictly increasing sequences. If  $(p_i)_{i=1}^{\infty}$  and  $(q_i)_{i=1}^{\infty}$  are elements of  $\mathcal{R}$ , then  $(p_i)_{i=1}^{\infty} < (q_i)_{i=1}^{\infty}$  means that  $p_i < q_i$  for every  $i \in \mathbf{N}$ . Select a  $p = (p_i)_{i=1}^{\infty}$  in  $\mathcal{R}$ . For every natural number  $n$  we define the *shrunkened endface in the  $n$ -coordinate direction* by

$$W_n = [-p_n, p_n]_1 \times \cdots \times [-p_n, p_n]_{n-1} \times \{1\}_n \times [-p_n, p_n]_{n+1} \\ \times [-p_n, p_n]_{n+2} \times \cdots \subset Q.$$

Note that  $W_n$  is itself a Hilbert cube and that it is a  $Z$ -set in  $Q$ . Observe, furthermore, that the  $W_n$ 's are pairwise disjoint. Let  $A$  be an infinite subset of  $\mathbf{N}$ . We define

$$Y(A) = Q \setminus \bigcup_{n \in A} W_n.$$

Since  $\lim_{n \rightarrow \infty} p_n = 1$  there exists a sequence of maps  $\alpha_n : Q \rightarrow W_n$  such that  $\lim_{n \rightarrow \infty} \alpha_n = 1_Q$ , where  $1_Q$  denotes the identity mapping on  $Q$ . This implies that the complement of  $Y(A)$  in  $Q$  is both dense and connected. Moreover, it follows that every compact subset of  $Y(A)$  is a  $Z$ -set in  $Q$ . The fake Hilbert space  $Y$  is represented by  $Y(\mathbf{N})$ .

**Definition.** If  $X' \subset X$  and  $Z' \subset Z$  then we say that the pair  $(X', X)$  is homeomorphic to the pair  $(Z', Z)$ , notation  $(X', X) \approx (Z', Z)$ , if there is a homeomorphism  $h : X \rightarrow Z$  such that  $h(X') = Z'$ .

It is shown in Dijkstra [6, 4.4.3] that every  $p \in \mathcal{R}$  leads to the same topological type  $(Y, Q)$ . If  $X$  is a space, then  $\mathcal{H}(X)$  denotes the group of autohomeomorphisms of  $X$ . The following theorem was taken from Dijkstra [6, 4.3.6].

**The Unknotting Theorem.** *Let  $\mathcal{U}$  be an open covering of  $Q$ , let  $C$  be a compact metric space and let  $F : C \times [0, 1] \rightarrow Q$  be a homotopy that is limited by  $\mathcal{U}$  (i.e., the paths  $F(\{c\} \times [0, 1])$  are contained in elements of  $\mathcal{U}$ ). If  $F_0$  and  $F_1$  are imbeddings of  $C$  in  $Y$ , then there is an  $h \in \mathcal{H}(Q)$  such that  $h \circ F_0 = F_1$ ,  $h$  is  $\mathcal{U}$ -close to  $1_Q$  and  $h(W_n) = W_n$  for every  $n$ .*

The following theorem will be used several times in Section 3.

**The Sierpiński Theorem [12].** *No continuum can be partitioned into countably many pairwise disjoint nonempty closed sets.*

A space is called *continuum-connected* if for every two points of the space there is a continuum that contains them both. The Sierpiński Theorem is also valid for continuum-connected spaces. The *continuum-components* of a space are maximally continuum-connected subspaces.

**3. Stability of  $Y$ .** In this section we show that  $Y$  is stable under multiplication with compact absolute retracts only. We also consider a few other fake Hilbert spaces.

**Lemma 1.** *If  $A$  is an infinite subset of  $\mathbf{N}$ , then  $(Y(A), Q) \approx (Y, Q)$ .*

*Proof.* It is shown in Dijkstra [6, 4.4.4] that individual shrunken endfaces can be deleted. This proves that  $(Y(A), Q) \approx (Y, Q)$  if the complement of  $A$  is finite.

Consider now the case that  $A$  has an infinite complement. Precisely as for  $(Y, Q)$ , we have that every choice of  $p \in \mathcal{R}$  leads to the same topological type  $(Y(A), Q)$ . Moreover, if  $B$  is another set with infinite complement, then a simple permutation of coordinates shows that  $(Y(B), Q) \approx (Y(A), Q)$ . So it suffices to show that  $(Y, Q)$  is homeomorphic to, for instance,  $(Y(\mathbf{N}_{\text{even}}), Q)$ , or, separating even and odd coordinates, that  $(Y, Q)$  is homeomorphic to the pair  $(X, Q \times Q)$  where

$$X = (Q \times Q) \setminus \bigcup_{i=1}^{\infty} (W_i \times Q(p_i)).$$

Consider the pair  $(Y, Q)$ . According to Dijkstra [6, 4.4.4] there exists a  $\chi \in \mathcal{H}(Q)$  such that  $\chi(W_{2i-1}) = W_{2i-1}$  and

$$\chi(W_{2i}) = W_{2i-1}^- = [-p_{2i}, p_{2i}]_1 \times \cdots \times [-p_{2i}, p_{2i}]_{2i-2} \\ \times \{-1\}_{2i-1} \times [-p_{2i}, p_{2i}]_{2i} \times [-p_{2i}, p_{2i}]_{2i+1} \times \cdots$$

for every  $i \in \mathbf{N}$ . This means that in the pair  $(\chi(Y), Q)$  the even coordinates are interchangeable. Doubling the even coordinates we see that  $(\chi(Y), Q)$  is homeomorphic to  $(Z, Q \times Q)$  where

$$Z = (Q \times Q) \setminus \bigcup_{i=1}^{\infty} ((W_{2i-1}^- \times Q(p_{2i})) \cup (W_{2i-1} \times Q(p_{2i-1}))).$$

Observe now that  $\chi^{-1} \times 1_Q$  is a homeomorphism from  $(Z, Q \times Q)$  onto  $(X, Q \times Q)$ . This proves the Lemma.  $\square$

**Lemma 2.**  $(Y, Q) \approx (Y \times Q, Q \times Q)$ .

*Proof.* This proof uses Anderson's Convergence Criterion which states that if every element of a sequence  $(h_i)_{i=1}^{\infty}$  in  $\mathcal{H}(Q)$  can be chosen arbitrarily close to  $1_Q$ , then there is a selection possible such that  $\lim_{n \rightarrow \infty} h_n \circ \cdots \circ h_1 \in \mathcal{H}(Q)$  (see Dijkstra [6, 1.1.2] for a precise formulation).

Consider the Unknotting Theorem. Observe that the shrunken end-faces are the continuum-components of  $Q \setminus Y$  (Sierpiński) and, hence, the Theorem is a topological statement about the pair  $(Y, Q)$ . This means that the Theorem is valid for any pair  $(C, D)$  that is homeomorphic to  $(Y, Q)$ . Specifically, consider a pair  $(Z, Q \times Q)$  that is homeomorphic to  $(Y, Q)$  and let  $\rho$  be a standard convex metric on  $Q \times Q$ . Using a straight line homotopy the Unknotting Theorem leads to: if  $\varepsilon > 0$ ,  $C$  and  $D$  are compact subsets of  $Z$  and  $h$  is a homeomorphism from  $C$  onto  $D$  with  $\rho(h, 1_{Q \times Q}) < \varepsilon$ , then there is a  $\bar{h} \in \mathcal{H}(Q \times Q)$  which extends  $h$  and which has the properties  $\rho(\bar{h}, 1_{Q \times Q}) < \varepsilon$  and  $\bar{h}(W) = W$  for every continuum-component  $W$  of  $(Q \times Q) \setminus Z$ .

Consider now  $(X, Q \times Q)$  where

$$X = Q \times Q \setminus \bigcup_{i=1}^{\infty} W_i \times Q(p_i).$$

Let  $n_1$  be a natural number. Lemma 1 implies that  $(Q \times Q \setminus \cup_{i>n_1} W_i \times Q(p_i), Q \times Q) \approx (Y, Q)$ , and, hence, there exists an  $h_1 \in \mathcal{H}(Q \times Q)$  with  $h_1(W_i \times Q(p_i)) = W_i \times Q(p_i)$  if  $i > n_1$  and  $h_1(W_{n_1} \times Q(p_{n_1})) = W_{n_1} \times Q$ . By choosing  $n_1$  large we can get  $p_{n_1}$  as close to 1 as we want, which means that we can make the distance of  $h_1$  towards  $1_{Q \times Q}$  arbitrarily small. As step 2 of the induction select an  $n_2 > n_1$  and an  $h_2 \in \mathcal{H}(Q \times Q)$  such that  $h_2$  fixes  $W_{n_1} \times Q$ ,  $h_2(W_{n_2} \times Q(p_{n_2})) = W_{n_2} \times Q$  and  $h_2(W_i \times Q(p_i)) = W_i \times Q(p_i)$  if  $i > n_2$ . Continue this process. Since every  $h_n$  can be chosen arbitrarily close to  $1_{Q \times Q}$  we may assume that  $h = \lim_{n \rightarrow \infty} h_n \circ \dots \circ h_1 \in \mathcal{H}(Q \times Q)$ . Note that  $h$  has the property  $h(W_{n_i} \times Q(p_{n_i})) = W_{n_i} \times Q$  for  $i \in \mathbf{N}$ . Consequently, we have  $(Q \times Q \setminus \cup_{i=1}^{\infty} W_{n_i} \times Q(p_{n_i}), Q \times Q) \approx (Y(A) \times Q, Q \times Q)$  where  $A = \{n_i \mid i \in \mathbf{N}\}$ . According to Lemma 1, the first pair is homeomorphic to  $(Y, Q)$  and the second to  $(Y \times Q, Q \times Q)$ .

**Lemma 3.** *If a connected space is a product of two noncompact spaces, then it has a continuum-connected remainder in any compactification.*

*Proof.* Consider two connected spaces  $X_1$  and  $X_2$ , neither of them compact. Let  $C$  be a compactification of  $X_1 \times X_2$  and put  $R = C \setminus (X_1 \times X_2)$ . Select for  $i = 1, 2$  a filter  $\mathcal{F}_i$  in  $X_i$  without cluster points. Let  $p$  be a cluster point in  $C$  of the filter  $\mathcal{F}$  that is generated by  $\{F_1 \times F_2 \mid F_i \in \mathcal{F}_i\}$  and note that  $p \in R$ . We shall show that for any  $q \in R$  there is a continuum in  $R$  that contains both  $p$  and  $q$ .

Let  $q$  be an arbitrary point of  $R$  and select an ultrafilter  $\mathcal{G}$  in  $C$  that converges to  $q$  and contains the set  $X_1 \times X_2$ . Let for  $i = 1, 2$  the ultrafilter  $\mathcal{G}_i$  be given by

$$\mathcal{G}_i = \{\pi_i(G) \mid G \in \mathcal{G} \text{ and } G \subset X_1 \times X_2\},$$

where  $\pi_i : X_1 \times X_2 \rightarrow X_i$  is the projection. Observe that since  $\mathcal{G}_1$  and  $\mathcal{G}_2$  are ultrafilters and  $\mathcal{G}$  has no cluster points in  $X_1 \times X_2$  we find that  $\mathcal{G}_1$  or  $\mathcal{G}_2$  has no cluster points in  $X_1$ , respectively,  $X_2$ . We may assume without loss of generality that  $\mathcal{G}_2$  has no cluster points. Consider the collection

$$\mathcal{K} = \{\text{cl}_C((X_1 \times G) \cup (F \times X_2)) \mid F \in \mathcal{F}_1 \text{ and } G \in \mathcal{G}_2\}.$$

Note that since  $X_1$  and  $X_2$  are connected and  $F$  and  $G$  are nonempty, the set  $(X_1 \times G) \cup (F \times X_2)$  is connected and, hence,  $\mathcal{K}$  is a collection of continua. Observe that the inclusion relation makes an inverse system of  $\mathcal{K}$  and, hence, the limit  $\cap \mathcal{K}$  is a continuum, see Engelking [9, 6.1.18]. Since  $\mathcal{K} \subset \mathcal{F} \cap \mathcal{G}$ , we have  $\{p, q\} \subset \cap \mathcal{K}$ . It remains to verify that  $\cap \mathcal{K}$  is contained in  $R$ . Let  $x$  be an arbitrary point of  $X_1 \times X_2$ . Since  $\mathcal{F}_1$  and  $\mathcal{G}_2$  have no cluster points there exists a neighborhood  $U \times V$  of  $x$  in  $X_1 \times X_2$  such that  $X_1 \setminus U \in \mathcal{F}_1$  and  $X_2 \setminus V \in \mathcal{G}_2$ . Consequently, the set

$$\text{cl}_C((X_1 \times (X_2 \setminus V)) \cup ((X_1 \setminus U) \times X_2))$$

is an element of  $\mathcal{K}$  that does not contain  $x$ . So we may conclude that  $\cap \mathcal{K} \cap (X_1 \times X_2) = \emptyset$ .

**Stability Theorem.** *The product  $Y \times A$  is homeomorphic to  $Y$  if and only if  $A$  is a compact absolute retract.*

*Proof.* Let  $A$  be a compact AR. According to Edwards (see Chapman [4, §44]) we have  $A \times Q \approx Q$ . This leads to

$$Y \approx Y \times Q \approx Y \times Q \times A \approx Y \times A.$$

Now assume that  $Y \times A \approx Y$ . The space  $A$  is obviously a complete AR. Since  $Q \setminus Y$  is a countable disjoint union of Hilbert cubes, we have according to Sierpiński that it is not continuum-connected. With Lemma 3 we find that  $A$  is compact.  $\square$

A more ambitious task is to find all factors of  $Y$ . We make the following

**Conjecture.** *If  $Y \cong A \times B$ , then  $Y \cong A$  and  $B$  is a compact AR or vice versa.*

Note that if  $Y \approx A \times B$  then according to Lemma 3 one of the factors, say  $B$ , is compact and, hence,  $A \times Q \approx A \times B \times Q \approx Y \times Q \approx Y$ . So the conjecture is equivalent with the statement: if  $A \times Q \approx Y$ , then  $A \approx Y$ . Note that this statement is true for the Hilbert space  $\ell^2$  instead of  $Y$ , Mogilski [11]. Unfortunately, Mogilski's proof leans heavily on

the unknotting theorem for noncompact  $Z$ -sets in  $\ell^2$ . A similar theorem does not exist for  $Y$  (see Dijkstra [6, 4.3.10]).

We shall now consider two derived fake Hilbert spaces. The first one appears in Anderson et al. [1] and is obtained by deleting a countable dense subset  $D$  from  $Y$ . The second one can be found in Dijkstra and van Mill [9] and is obtained by deleting a so-called 0-dimensional capset  $A_0$  from  $Y$  ( $A_0$  is a dense copy of the product of the Cantor set and the set of rational numbers). Both spaces are complete AR's.

**Proposition 1.** *If  $X$  is either  $Y \setminus D$  or  $Y \setminus A_0$ , then  $X \times A \approx X$  only if  $A$  is a singleton.*

*Proof.* Assume that  $X \times A \approx X$  and note that  $Q \setminus X$  is not continuum-connected. In fact, the Sierpiński Theorem implies that the continuum-components of  $Q \setminus X$  are the shrunken endfaces plus the singletons of  $D$ , respectively  $A_0$ . So, according to Lemma 3 the factor  $A$  is compact.

Let  $f$  be the homeomorphism from  $X$  onto  $X \times A$ . Since  $Q \times A$  is a Hilbert cube (Edwards) we may apply Lemma 3.6 of Anderson et al. [1] to find a compact space  $M$  and monotone maps  $\alpha : M \rightarrow Q$  and  $\beta : M \rightarrow Q \times A$  such that  $\alpha^{-1}(X) = \beta^{-1}(X \times A)$  and  $f \circ \alpha \mid \alpha^{-1}(X) = \beta \mid \alpha^{-1}(X)$ . Recall that a monotone map is a continuous, closed surjection with the property that the preimage of each connected set is also connected. This means that the preimage under  $\alpha$  (or  $\beta$ ) of a continuum-component of  $Q \setminus X$  (or  $(Q \setminus X) \times A$ ) is a continuum-component of  $M \setminus \alpha^{-1}(X)$ . Since the singletons of  $D$ , respectively  $A_0$ , are continuum-components in  $Q \setminus X$ , it is possible to find a sequence  $(C_i)_{i=1}^{\infty}$  of continuum-components of  $Q \setminus X$  that has only one cluster point in  $Q$ , namely, some point  $x$  in  $X$ . So we have that every  $\alpha^{-1}(C_i)$  is a continuum-component of  $M \setminus \alpha^{-1}(X)$  and that every  $\beta(\alpha^{-1}(C_i))$  is a continuum-component of  $(Q \setminus X) \times A$ . Since  $A$  is a continuum this implies that each  $\beta(\alpha^{-1}(C_i))$  has the form  $F_i \times A$ . This means that the set of cluster points of  $\beta(\alpha^{-1}(C_i))_{i=1}^{\infty}$  in  $Q \times A$  also has the form  $F \times A$ . On the other hand,  $F \times A$  is contained in  $\beta(\alpha^{-1}(\{x\})) = \{f(x)\}$ . Since  $F \times A$  is nonempty we may conclude that  $A$  is a singleton.

**4. The pair  $(Y \times Y, Q \times Q)$ .** In Anderson et al. [1] it is shown that  $Y \times Y$  is homeomorphic to  $\ell^2$ . The aim of this section is to prove

a slightly stronger result, namely that  $(Y \times Y, Q \times Q)$  is homeomorphic to  $(s, Q)$ , where  $s$  is the pseudo-interior of  $Q$ . This means that we have to show that  $Q \times Q \setminus Y \times Y$  is a capset in  $Q \times Q$ . We shall use the following theorem.

**The Capset Characterization Theorem** (Curtis [5, Corollary 4.9]). *Let  $(B_i)_{i=1}^\infty$  be a sequence of subsets of  $Q$ . If  $(B_i)_{i=1}^\infty$  satisfies the properties*

- (1) *each  $B_i$  is a Z-set in  $Q$ ,*
  - (2) *each  $B_i$  is homeomorphic to  $Q$ ,*
  - (3) *each  $B_i$  is a Z-set in  $B_{i+1}$ , and*
  - (4) *there is a homotopy  $H : Q \times [0, 1] \rightarrow Q$  such that  $H_0 = 1_Q$  and, for every  $t \in (0, 1]$ , there is an  $n \in \mathbf{N}$  such that  $H(Q \times [t, 1]) \subset B_n$ ,*
- then  $(B_i)_{i=1}^\infty$  is a capset.*

**Proposition 2.**  $(Y \times Y, Q \times Q) \approx (s, Q)$ .

*Proof.* In order to show that  $Q \times Q \setminus Y \times Y$  is a capset in  $Q \times Q$  it suffices to prove that it contains a capset, see Bessaga and Pełczyński [2, Theorem IV.4.2]. We introduce some notations. If  $p \in \mathcal{R}^\dagger$ , then the shrunken endfaces it determines are denoted by  $W_n(p)$ . Furthermore, let  $C_n(p)$  stand for the set

$$\begin{aligned} &[-p_{n+1}, p_{n+1}]_1 \times \cdots \times [-p_{n+1}, p_{n+1}]_{n-1} \times J_n \times J_{n+1} \\ &\quad \times [-p_{n+1}, p_{n+1}]_{n+2} \times [-p_{n+1}, p_{n+1}]_{n+3} \times \cdots \end{aligned}$$

Note that  $C_n(p)$  is just as  $W_n(p)$  a convex Z-set in  $Q$  that is homeomorphic to  $Q$ . Moreover,  $C_n(p)$  contains  $W_n(p)$  and  $W_{n+1}(p)$  as Z-sets but does not meet any of the other shrunken endfaces. Let  $B_n(p)$  be given by

$$\begin{aligned} B_n(p) = & (W_1(p) \times C_1(p)) \cup (C_1(p) \times W_2(p)) \cup (W_2(p) \times C_2(p)) \\ & \cup (C_2(p) \times W_3(p)) \cup \cdots \cup (W_n(p) \times C_n(p)) \cup (C_n(p) \times W_{n+1}(p)). \end{aligned}$$

Note that in this union only adjacent terms have a nonempty intersection. We need the following fact: if  $X$  is a space that is the union of



two Hilbert cubes  $X_1$  and  $X_2$  such that  $X_1 \cap X_2 = X_0$  is also a Hilbert cube and, moreover, a Z-set in both  $X_1$  and  $X_2$ , then  $X$  is a Hilbert cube. This result can easily be obtained by observing that each pair  $(X_0, X_1)$  and  $(X_0, X_2)$  is homeomorphic to  $(F, Q)$  where  $F$  is an end-face of  $Q$ , or it can be seen as a special case of a much stronger theorem by Handel [10]. Moreover, if  $Z$  is a subset of  $X$  such that  $Z \cap X_i$  is a Z-set in  $X_i$  for  $i = 0, 1, 2$ , then one easily verifies that  $Z$  is a Z-set in  $X$ . Note that  $(W_i(p) \times C_i(p)) \cap (C_i(p) \times W_{i+1}(p)) = W_i(p) \times W_{i+1}(p)$  and that  $(C_i(p) \times W_{i+1}(p)) \cap (W_{i+1}(p) \times C_i(p)) = W_{i+1}(p) \times W_{i+1}(p)$  and, hence, every set  $B_n(p)$  is a Hilbert cube. If  $p < q \in \mathcal{R}^\dagger$ , then  $W_n(p)$  is a Z-set in  $W_n(q)$  and  $C_n(p)$  is a Z-set in  $C_n(q)$ . A tedious but straightforward argument involving the second part of the aforementioned fact now yields that  $B_n(p)$  is a Z-set in  $B_n(q)$ .

Select a sequence  $p^1 < p^2 < p^3 < \dots$  in  $\mathcal{R}^\dagger$  that has an upper bound  $q \in \mathcal{R}^\dagger$ . The sequence that satisfies the Capset Characterization Theorem is  $(B_n(p^n))_{n=1}^\infty$ . It is obvious that every  $B_n(p^n)$  is contained in

$$\bigcup_{i=1}^{\infty} (W_i(q) \times Q) \cup (Q \times W_i(q)) = Q \times Q \setminus Y \times Y$$

and, hence, every  $B_n(p^n)$  is a Z-set in  $Q \times Q$ . Every  $B_n(p^n)$  is a Hilbert cube and  $B_n(p^n)$  is contained in  $B_{n+1}(p^n)$  which is a Z-set in  $B_{n+1}(p^{n+1})$ . It remains to show that  $(B_n(p^n))_{n=1}^\infty$  satisfies property (4). Obviously, it suffices to show that  $(B_n(p^1))_{n=1}^\infty$  has this property. Select maps  $\alpha_i : Q \rightarrow W_i(p^1)$  such that  $\lim_{i \rightarrow \infty} \alpha_i = 1_Q$ . Consider the following sequence of maps from  $Q \times Q$  into  $Q \times Q$

$$\alpha_1 \times \alpha_1, \alpha_1 \times \alpha_2, \alpha_2 \times \alpha_2, \alpha_2 \times \alpha_3, \alpha_3 \times \alpha_3, \dots$$

Connect adjacent maps in this sequence by straight line homotopies. Since  $C_n(p^1)$  is a convex set which contains  $W_n(p^1)$  and  $W_{n+1}(p^1)$  we have that the image of the homotopy connecting  $\alpha_n \times \alpha_n$  with  $\alpha_n \times \alpha_{n+1}$  is contained in  $W_n(p^1) \times C_n(p^1) \subset B_n(p^1)$  and that the image of the homotopy that connects  $\alpha_n \times \alpha_{n+1}$  with  $\alpha_{n+1} \times \alpha_{n+1}$  is contained in  $C_n(p^1) \times W_{n+1}(p^1) \subset B^n(p^1)$ . "Glued" together these homotopies form an  $H$  as in property (4) of the Capset Characterization Theorem. This proves the proposition.

## REFERENCES

1. R.D. Anderson, D.W. Curtis and J. van Mill, *A fake topological Hilbert space*, Trans. Amer. Math. Soc. **272** (1982), 311–321.
2. C. Bessaga and A. Pełczyński, *Selected topics in infinite-dimensional topology*, PWN, Warsaw, 1975.
3. P.L. Bowers, *Fake boundary sets in the Hilbert cube*, Proc. Amer. Math. Soc. **93** (1985), 121–127.
4. T.A. Chapman, *Lectures on Hilbert cube manifolds*, CMBS Regional Conf. Ser. in Math., no. 28, Amer. Math. Soc., Providence, RI, 1976.
5. D.W. Curtis, *Boundary sets in the Hilbert cube*, Topology Appl. **20** (1985), 201–221.
6. J.J. Dijkstra, *Fake topological Hilbert spaces and characterizations of dimension in terms of negligibility*, Dissertation, Univ. of Amsterdam, 1983; publ. as CWI Tracts 2, Centre for Math. and Comp. Sci., Amsterdam, 1984.
7. ———, *A rigid space whose square is the Hilbert space*, Proc. Amer. Math. Soc. **93** (1985), 118–120.
8. J.J. Dijkstra and J. van Mill, *Fake topological Hilbert spaces and characterizations of dimension in terms of negligibility*, Fund. Math. **125** (1985), 143–153.
9. R. Engelking, *General topology*, PWN, Warsaw, 1977.
10. M. Handel, *On certain sums of Hilbert cubes*, General Topology and Appl. **9** (1978), 19–28.
11. J. Mogilski, *CE-decomposition of  $\ell_2$ -manifolds*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astr. Phys. **27** (1979), 309–314.
12. W. Sierpiński, *Un théorème sur les continus*, Tôhoku Math. J. **13** (1918), 300–303.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WASHINGTON, SEATTLE, WA 98195

*Current address:* Department of Mathematics, University of Alabama, Tuscaloosa, AL 35487

SUBFACULTEIT WISKUNDE, VRIJE UNIVERSITEIT AND MATHEMATISCH INSTITUUT, UNIVERSITEIT VAN AMSTERDAM, AMSTERDAM, THE NETHERLANDS.