

AN AR-MAP WHOSE RANGE IS MORE INFINITE-DIMENSIONAL THAN ITS DOMAIN

J. J. DIJKSTRA, J. VAN MILL, AND J. MOGILSKI

(Communicated by James E. West)

ABSTRACT. We construct an example of an AR-map $f : X \rightarrow Y$, where X is a strongly countable dimensional compact AR and Y is a countable dimensional AR which is not strongly countable dimensional. Using this map we find a shrinkable decomposition of the pre-Hilbert space l_f^2 whose quotient map does not stabilize to a near homeomorphism. We also present a partial result concerning the question whether cell-like maps preserve countable dimensionality.

1. INTRODUCTION

A space X is (*strongly*) *countable dimensional* if it is a countable union of (closed) finite-dimensional subspaces. The aim of this note is to construct an example of an AR-map $f : X \rightarrow Y$, where X is a strongly countable dimensional compact AR and Y is a countable dimensional compact AR which is not strongly countable dimensional. Observe that the range of our map is “more infinite-dimensional than its domain.” The simplicity of our example shows that this phenomenon has nothing to do with dimension raising cell-like maps (Dranišnikov [4]). This example can be used to find a shrinkable AR-map whose domain is the pre-Hilbert space

$$l_f^2 = \{(x_i)_i \in l^2 : x_i = 0 \text{ for all but finitely many } i\},$$

but that does not stabilize to a near homeomorphism. This answers several questions of J. P. Henderson (see West [13]). In the final section we discuss the problem whether cell-like maps between compact ANRs preserve countable dimensionality.

All spaces are assumed to be separable metric. As usual, I denotes the interval $[0, 1]$.

2. THE BASIC EXAMPLE

Proposition 2.1. *There exists an AR-map f from a strongly countable dimensional compact AR X onto a countable dimensional but not strongly countable dimensional compact AR Y .*

Received by the editors June 19, 1990.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 54G20, 57N20.

©1992 American Mathematical Society
0002-9939/92 \$1.00 + \$.25 per page

Proof. The construction is inspired by the classic examples of nonstrongly countable dimensional spaces, see [6, Example 1.12]. First, we construct the range of the map f . Let D denote a disk. Pick a null-sequence $\{J_k\}_{k=1}^\infty$ consisting of pairwise disjoint arcs in D the union of which is dense. For every k , let $f_k : J_k \rightarrow I^k$ be a continuous surjection with finite fibres. Standard examples of space-filling curves have this property. We recall the construction of one such map. Define the homeomorphism h between the Cantor sets $3^\mathbb{N} = \prod_{i=1}^\infty \{0, 1, 2\}$ and $3^\mathbb{N} \times 3^\mathbb{N}$ by $h(\varepsilon) = (\alpha, \beta)$, where

$$\alpha_i = \begin{cases} \varepsilon_{2i-1}, & \text{for } \sum_{j<i} \beta_j \text{ even,} \\ 2 - \varepsilon_{2i-1}, & \text{for } \sum_{j<i} \beta_j \text{ odd,} \end{cases}$$

and

$$\beta_i = \begin{cases} \varepsilon_{2i}, & \text{for } \sum_{j \leq i} \alpha_j \text{ even,} \\ 2 - \varepsilon_{2i}, & \text{for } \sum_{j \leq i} \alpha_j \text{ odd.} \end{cases}$$

Let $q : 3^\mathbb{N} \rightarrow I$ stand for the usual quotient map $q(\varepsilon) = \sum_{i=1}^\infty \varepsilon_i 3^{-i}$. Since the fibres of q contain at most two points, the fibres of $(q \times q) \circ h$ contain at most four points. One readily verifies that $q(\varepsilon) = q(\varepsilon')$ implies $(q \times q) \circ h(\varepsilon) = (q \times q) \circ h(\varepsilon')$. So there exists a continuous map $p : I \rightarrow I \times I$ with $(q \times q) \circ h = p \circ q$. Then p is obviously surjective and its fibres consist of at most four points. The functions f_k are obtained by iterating p .

Put

$$\mathcal{G} = \{ \{y : y = x \text{ or } f_k(y) = f_k(x) \text{ for some } k \in \mathbb{N}\} : x \in D \}.$$

Then \mathcal{G} is an upper semicontinuous decomposition of D , so we can consider the quotient space $Y = D/\mathcal{G}$ with corresponding quotient map $\pi : D \rightarrow Y$. Observe that Y is obtained from D by replacing each J_k by I^k . It clearly follows that Y is countable dimensional. We claim that Y is not strongly countable dimensional. This will follow by a standard argument. Let $U \subset Y$ be nonempty and open. Then $\pi^{-1}(U)$ is open in D and nonempty so it contains infinitely many J_k 's. This implies that U contains cells of arbitrarily large dimension, i.e., $\dim U = \infty$. Now assume that $Y = \bigcup_{i=1}^\infty A_i$, with each A_i closed. Then by the Baire Category Theorem there exists an i such that A_i has nonempty interior and hence we have $\dim A_i = \infty$. We conclude that Y is not strongly countable dimensional. By a direct argument, it is easily seen that Y is an AR. Alternatively, apply Kozłowski [10, Theorem 15]: the image of an ANR under a map f that is determined on a null-sequence of ARs A_i such that each $f(A_i)$ is an AR, is again an ANR.

We now look at the domain. Define

$$Z = (D \times \{0\}) \cup \bigcup_{k=1}^\infty J_k \times [0, 1/k].$$

It is easy to see that Z is a closed subspace of $D \times I$. Observe that the collection $\{J_k \times \{\frac{1}{k}\}\}_{k=1}^\infty$ is a null sequence in Z . For every k let $g_k : J_k \times \{\frac{1}{k}\} \rightarrow I^k$ be defined by $g_k(x, \frac{1}{k}) = f_k(x)$, i.e., g_k is a copy of the map f_k . Now consider the collection

$$\mathcal{H} = \{ \{y : y = x \text{ or } g_k(y) = g_k(x) \text{ for some } k \in \mathbb{N}\} : x \in Z \}.$$

Then \mathcal{H} is an upper semicontinuous decomposition of Z , so we can consider the quotient space $X = Z/\mathcal{H}$ with corresponding quotient map $\rho : Z \rightarrow X$.

Observe that X is a disc with the mapping cylinders of the maps f_k attached to it. Indeed, for every k we have that the set

$$\rho(J_k \times [0, 1/k])$$

is a canonical copy of the mapping cylinder of f_k . Observe that X is an AR for the same reason as Y is. Also, observe that Z is strongly countable dimensional. If we remove $\bigcup_{k=1}^\infty \rho(J_k \times \{\frac{1}{k}\})$ from Z then what remains is (homeomorphic to) a σ -compact subspace of $D \times I$.

Finally, we define the map f . For every k let $c_k: \rho(J_k \times [0, \frac{1}{k}]) \rightarrow I^k$ be the collapse to the base. Then c_k is clearly an AR-map (its point inverses are cones of finite sets). Put

$$\mathcal{F} = \{ \{y : y = x \text{ or } c_k(y) = c_k(x) \text{ for some } k \in \mathbf{N} \} : x \in X \}.$$

Then \mathcal{F} is upper semicontinuous, and the decomposition space X/\mathcal{F} is clearly homeomorphic to Y . Since the quotient map $X \rightarrow X/\mathcal{F}$ is an AR-map, we are done.

For a σ -compact space X , let $\gamma(X)$ be the minimum ordinal α such that X is a countable union of compact subsets with transfinite dimension $\text{ind} < \alpha$ (cf. [2, p. 282]). Let us recall that the transfinite dimension ind is the extension by transfinite induction of the classical Menger-Urysohn inductive dimension ($\text{ind } X = -1$ means $X = \emptyset$, $\text{ind } X \leq \alpha$, where α is a countable ordinal, if and only if X has an open basis consisting of sets B with $\text{ind}(\partial B) < \alpha$). Obviously, we have $\gamma(X) \leq \text{ind } X$ and a compactum X is strongly countable dimensional if and only if $\gamma(X) \leq \omega_0$. In Proposition 2.1 we constructed an AR-map of a compact AR X , with $\gamma(X) = \omega_0$, onto a compact AR Y , with $\gamma(Y) = \omega_0 + 1$. By a slight modification of our construction we obtain:

Proposition 2.2. *For every countable ordinal α , $\alpha \geq \omega_0$, there exists an AR-map f_α from a strongly countable dimensional compact AR X_α onto a compact AR Y_α with $\gamma(Y_\alpha) \geq \alpha$.*

Proof. As in the proof of Proposition 2.1, we pick a null-sequence $\{J_k\}_{k=1}^\infty$ consisting of pairwise disjoint arcs in a disk D . Let H^α , for $\alpha < \omega_1$, stand for a slight modification of Henderson’s “transfinite cubes.” We define H^α together with a point $p^\alpha \in H^\alpha$ as follows. $H^0 = \{0\}$, $H^{\alpha+1} = H^\alpha \times I$, and $p^{\alpha+1} = (p^\alpha, 0)$. If α is a limit ordinal then we write $\{\beta : \beta < \alpha\} = \{\beta_i : i \in \mathbf{N}\}$ and we connect in the topological sum $\bigoplus_{i=1}^\infty H^{\beta_i}$ every p^{β_i} with $p^{\beta_{i+1}}$ by an arc. We then take the one point compactification of the resulting space. This is H^α and p^α is the point at infinity. The space H^α is a strongly countable dimensional compact AR with $\text{ind} = \alpha$ (see [8]).

Now, let α_k be a nondecreasing sequence converging to α . For every k , let $g_k: J_k \rightarrow H^{\alpha_k}$ be a continuous surjection with finite fibres (it is not hard to construct g_k using the map p described in the proof of Proposition 2.1). We construct the spaces X_α , Y_α and the map f_α in the same way as X , Y , and f were constructed. Hence X_α and Y_α are compact ARs and f_α is an AR-map. By strong countable dimensionality of H^{α_k} , for $k = 1, 2, \dots$, X_α is strongly countable dimensional. Since each nonempty subset of Y_α contains infinitely many H^{α_k} ’s we have $\gamma(Y_\alpha) \geq \alpha$.

3. A SHRINKABLE DECOMPOSITION OF l_f^2

The pre-Hilbert space l_f^2 is strongly countable dimensional and universal for all strongly countable dimensional compacta. Henderson and Walsh [9] constructed a cell-like map f from l_f^2 onto an AR M such that M is not homeomorphic to l_f^2 but $M \times \mathbf{R} \cong l_f^2$. Proposition 2.1 allows us to construct an AR-map of l_f^2 onto an AR Z such that not even $Z \times l_f^2$ is homeomorphic to l_f^2 . Our map has another interesting property. Let us recall that a proper map $f : X \rightarrow Y$ is *shrinkable* if for every open cover \mathcal{U} of Y and every open cover \mathcal{V} of X there exists a homeomorphism $h : X \rightarrow X$ such that $f \circ h$ is \mathcal{U} -close to f and $\{h(f^{-1}(y))\}_{y \in Y}$ refines \mathcal{V} . Bing's Shrinking Criterion states that if X is complete and f is shrinkable, then Y is homeomorphic to X and f is a near homeomorphism. The following proposition shows that the completeness assumption is essential and it answers questions NCL 12, 13, and 14 in West [13].

Proposition 3.1. *There exists a shrinkable AR-map from l_f^2 onto an absolute retract Z such that $Z \times l_f^2$ is not homeomorphic to l_f^2 .*

Proof. Let X be the strongly countable dimensional AR from Proposition 2.1. We may assume that X is a subset of l_f^2 . By compactness X is a Z -set in l_f^2 . If $f : X \rightarrow Y$ is the AR-map from Proposition 2.1, then the decomposition

$$\mathcal{F} = \{f^{-1}(y) : y \in Y\} \cup \{\{x\} : x \in l_f^2 \setminus X\}$$

is upper semicontinuous and the quotient map $F : l_f^2 \rightarrow l_f^2/\mathcal{F}$ is an AR-map. By [10] the quotient space $Z = l_f^2/\mathcal{F}$ is an AR. Of course, Z and $Z \times l_f^2$ contain topological copies of the space Y which is not strongly countable dimensional. Hence, $Z \times l_f^2$ is not homeomorphic to the strongly countable dimensional space l_f^2 . We prove that F is shrinkable. Let \mathcal{U} and \mathcal{V} be open covers of Z and l_f^2 , respectively. Since F is a fine homotopy equivalence (see [7] or Theorem 4.2) there exists a map $G : Z \rightarrow l_f^2$ such that $G \circ F$ is $F^{-1}(\mathcal{U})$ -homotopic to the identity. The map $G \circ F|_X$ can be approximated by an embedding $v : X \rightarrow l_f^2$ such that v is $F^{-1}(\mathcal{U})$ -homotopic and \mathcal{V} -close to $G \circ F|_X$. Since X and $v(X)$ are Z -sets there exists a homeomorphism $h : l_f^2 \rightarrow l_f^2$ which is $F^{-1}(\text{st } \mathcal{U})$ -homotopic to the identity and such that $h|_X = v$. Thus $F \circ h$ and F are $\text{st } \mathcal{U}$ -close. Since $h|_X$ is \mathcal{V} -close to $G \circ F$ and $G \circ F(F^{-1}(y))$ is a singleton for all $y \in Y$ the family $\{h(F^{-1}(y)) : y \in Y\}$ refines $\text{st } \mathcal{V}$.

4. TRANSFINITE DIMENSION AND CELL-LIKE MAPS

The spaces in this section are assumed to be compact. The following question is closely related to the example of §2.

Question 4.1. Let $f : X \rightarrow Y$ be a cell-like map of a countable dimensional ANR X onto an ANR Y . Is Y countable dimensional?

Let us recall that a cell-like map $f : X \rightarrow Y$ between absolute neighborhood retracts is a *fine homotopy equivalence* (i.e., for every open cover \mathcal{U} of Y there exists a map $g : Y \rightarrow X$ such that $f \circ g$ is \mathcal{U} -homotopic to id_Y and $g \circ f$

is $f^{-1}(U)$ -homotopic to id_X) and a *hereditary shape equivalence* (i.e., for each ANR Z the map f produces a one-to-one correspondence between the homotopy classes of $C(Y, Z)$ and $C(X, Z)$). Hereditary shape equivalence is the natural extension to arbitrary compacta of fine homotopy equivalence between ANRs. Let us have a closer look at this. Let X be a countable dimensional compactum and let $f: X \rightarrow Y$ be a cell-like map. By the Freudenthal Expansion Theorem (see [3]) X is the inverse limit of finite dimensional ANRs, say $X = \varprojlim \{X_n, f_n\}$, with each X_n ANR. Let \tilde{X} be the infinite mapping cylinder of the sequence $\{X_n, f_n\}_{n=1}^\infty$ with a copy of X attached at its end. Then $\tilde{X} \in \text{ANR}$ and \tilde{X} is countable dimensional (observe that we added a countable dimensional set to X). Let $\mathcal{G}_f = \{f^{-1}(y) : y \in Y\} \cup \{\text{points}\}$, then \mathcal{G}_f is a cell-like decomposition of \tilde{X} and the quotient map $\tilde{f}: \tilde{X} \rightarrow \tilde{X}/\mathcal{G}_f$ is a cell-like map. Of course Y embeds in \tilde{X}/\mathcal{G}_f . Let us recall the following theorem of Kozłowski [10] (see also [5, p. 133]).

Theorem 4.2. *The following statements are equivalent:*

- (1) f is a hereditary shape equivalence;
- (2) $\tilde{X}/\mathcal{G}_f \in \text{ANR}$;
- (3) \tilde{f} is a fine homotopy equivalence;
- (4) \tilde{f} is a hereditary shape equivalence.

By Theorem 4.2, Question 4.1 is equivalent to the following question (posed by Henderson, Kozłowski, and Walsh at the problem session of the AMS meeting in Norman, 1983).

Question 4.3. Do hereditary shape equivalences preserve countable dimensionality?

A complete space X is countable dimensional if and only if $\text{ind } X$ exists. Because hereditary shape equivalences do not raise finite dimension we ask

Question 4.4. Let $f: X \rightarrow Y$ be a hereditary shape equivalence between countable dimensional compacta. Is $\text{ind } Y \leq \text{ind } X$?

Ancel [1] proved that a cell-like map with a countable dimensional range is a hereditary shape equivalence. We define

$$\eta(X) = \sup\{\text{ind } Y : Y \text{ is a countable dimensional cell-like image of } X\}.$$

We present the following partial result concerning Question 4.3.

Proposition 4.5. *If for every countable dimensional compactum X we have $\eta(X) < \omega_1$, then hereditary shape equivalences preserve countable dimensionality.*

Proof. Let $f: X \rightarrow Y$ be a hereditary shape equivalence of a countable dimensional compactum X onto Y . In order to prove that Y is countable dimensional we shall apply Pol's characterization of countable dimensionality (see [12]) but we modify his original construction slightly. Let $Y = \varprojlim \{Y_n, g_n\}$, where each Y_n is a finite dimensional ANR. As in the beginning of this section, \tilde{Y} stands for the infinite mapping cylinder of the sequence $\{Y_i, g_i\}_{i=1}^\infty$ with the copy of Y attached at its end. If \tilde{Y}_n is a finite mapping cylinder of the finite sequence $\{Y_i, g_i\}_{i=1}^n$ and if π_n is the natural collapse of \tilde{Y} onto \tilde{Y}_n , then

\tilde{Y}_n is a finite dimensional ANR, π_n is a cell-like map which is a hereditary shape equivalence and the sequence $\delta_n = \sup_{z \in \tilde{Y}_n} \text{diam}(\pi_n^{-1}(z))$, converges to 0. Let C be the Cantor set and let $C_0 = \{c_1, c_2, \dots\}$ be a dense countable subset of C . We shall denote by \mathcal{P} the upper semicontinuous decomposition of the space $C \times \tilde{Y}$ into singletons $\{(c, y)\}$, where $c \notin C_0$, and the sets $\{c_i\} \times \pi_i^{-1}(z)$, where $z \in \tilde{Y}_i$ and $i = 1, 2, \dots$. The space $C \times \tilde{Y}/\mathcal{P}$ is compact and the quotient map $\pi : C \times \tilde{Y} \rightarrow C \times \tilde{Y}/\mathcal{P}$ is a hereditary shape equivalence. According to Pol's characterization the space \tilde{Y} is countable dimensional if and only if

$$\sup_{D \in \widehat{C}_0} \text{ind}(\pi(D \times \tilde{Y})) < \omega_1,$$

where $\widehat{C}_0 = \{D : D \text{ is a compact subset of } C_0\}$. Now, let \widehat{X} be a compactum which is the infinite mapping cylinder of the sequence $\{Y_n, g_n\}$ with a copy of the space X attached at its end (the compactum \widehat{X} can be constructed as follows: assume that $X \subset I^\omega \times \{0\} \subset I^\omega \times [0, 1]$, then the quotient space $I^\omega \times [0, 1]/\mathcal{F}$ of the decomposition $\mathcal{F} = \{(q, t) : t \in (0, 1]\} \cup \{(q, 0) : (q, 0) \notin X\} \cup \{f^{-1}(y) : y \in Y\}$, is homeomorphic to $I^\omega \times [0, 1]$ (see [11, Lemma 7.5.2]) and $\hat{f}(X)$ is a Z -set, where \hat{f} is the quotient map; we set $\widehat{X} = \hat{f}^{-1}(\tilde{Y})$ assuming that \tilde{Y} is a subset of $I^\omega \times [0, 1]/\mathcal{F}$ whose intersection with $\hat{f}(I^\omega \times \{0\})$ is Y). By the construction \widehat{X} is countable dimensional, $\hat{f}|_X = f$ and \hat{f} maps $\widehat{X} \setminus X$ onto $\tilde{Y} \setminus Y$. Then $\phi = \text{id}_C \times (\hat{f}|_{\widehat{X}})$ is a hereditary shape equivalence of $C \times \widehat{X}$ onto $C \times \tilde{Y}$. Thus $\pi \circ \phi$ is a hereditary shape equivalence. If $D \in \widehat{C}_0$, then

$$\text{ind} \pi(D \times \tilde{Y}) = \text{ind}(\pi \circ \phi(D \times \widehat{X}))$$

and the space $\pi \circ \phi(D \times \widehat{X})$ is a closed subset of a countable dimensional space which is a cell-like image of $C \times \widehat{X}$. Thus

$$\text{ind}(\pi(D \times \tilde{Y})) \leq \eta(C \times \widehat{X}).$$

By our assumption $\eta(C \times \widehat{X}) < \omega_1$.

Added in proof. It can be shown that the converse of Proposition 4.5 is also valid and that the answer to Question 4.4 is no, [14].

REFERENCES

1. F. D. Ancel, *The role of countable dimensionality in the theory of cell-like relations*, Trans. Amer. Math. Soc. **287** (1985), 1–40.
2. C. Bessaga and A. Pełczyński, *Selected topics in infinite-dimensional topology*, PWN, Warsaw, 1975.
3. K. Borsuk, *Shape theory*, PWN, Warsaw, 1975.
4. A. N. Dranišnikov, *On a problem of P. S. Alexandrov*, Mat. Sb. **135** (1988), 551–557.
5. J. Dydak and J. Segal, *Shape theory: An introduction*, Lectures Notes in Math., vol. 688, Springer-Verlag, Berlin, 1978.
6. R. Engelking and E. Pol, *Countable dimensional spaces. A survey*, Dissertationes Math. **216** (1983), 1–45.
7. W. E. Haver, *Mappings between ANR's that are fine homotopy equivalence*, Pacific J. Math. **58** (1975), 457–461.

8. D. W. Henderson, *A lower bound for transfinite dimension*, *Fund. Math.* **64** (1968), 167–173.
9. J. P. Henderson and J. J. Walsh, *Examples of cell-like decompositions of the infinite dimensional manifolds σ and Σ* , *Topology Appl.* **16** (1983), 143–154.
10. G. Kozłowski, *Images of ANR's*, unpublished manuscript.
11. J. van Mill, *Infinite-dimensional topology, prerequisites and introduction*, North-Holland, Amsterdam, 1989.
12. R. Pol, *On classification of weakly infinite dimensional compacta*, *Fund. Math.* **116** (1983), 169–188.
13. J. E. West, *Open problems in infinite-dimensional topology*, *Open Problems in Topology* (J. van Mill and G. M. Reed, eds.), North-Holland, Amsterdam, 1990, pp. 523–597.
14. J. J. Dijkstra and J. Mogilski, in preparation.

(J. J. Dijkstra, J. Mogilski) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALABAMA, TUSCALOOSA, ALABAMA, 35487-0350

Current address: J. Mogilski: Department of Mathematics, Bradley University, Peoria, Illinois 61625

E-mail address J. J. Dijkstra: jdijkstr@ualvm.bitnet

E-mail address: jdijkstr@ualvm.bitnet

E-mail address J. Mogilski: jmogilsk@bradley.edu

(J. van Mill) FACULTEIT WISKUNDE EN INFORMATICA, VRIJE UNIVERSITEIT, POSTBUS 7161, 1007 MC AMSTERDAM, THE NETHERLANDS

E-mail address: vanmill@cs.vu.nl