

# Quantum Information Theory, Spring 2019

## Exercise Set 13

## in-class practice problems

### 1. Symmetric subspace:

- Write out  $\Pi_2$  and  $\Pi_3$ .
- In class we wrote down a basis for  $\text{Sym}^2(\mathbb{C}^2)$ . Write down bases of  $\text{Sym}^2(\mathbb{C}^d)$  and  $\text{Sym}^3(\mathbb{C}^2)$ .
- Verify that  $R_\pi R_\tau = R_{\pi\tau}$  and  $R_\pi^\dagger = R_{\pi^{-1}}$  for all  $\pi, \tau \in S_n$ .
- Verify that  $\Pi_n = \frac{1}{n!} \sum_{\pi \in S_n} R_\pi$  is the orthogonal projection onto the symmetric subspace.

### 2. Integral formula: In this exercise you can prove the integral formula:

$$\Pi_n = \binom{n+d-1}{n} \int |\psi\rangle^{\otimes n} \langle\psi|^{\otimes n} d\psi =: \tilde{\Pi}_n$$

- Show that  $\tilde{\Pi}_n = \Pi_n \tilde{\Pi}_n$ .
  - In class we mentioned the following important fact: *If  $A \in L(\mathcal{X}^{\otimes n})$  is an operator such that  $[A, U^{\otimes n}] = 0$  for all unitaries  $U \in \text{U}(\mathcal{X})$ , then  $A$  is a linear combination of permutation operators  $R_\pi$ ,  $\pi \in S_n$ .* Use this fact to show that  $\tilde{\Pi}_n = \sum_\pi c_\pi R_\pi$  for suitable  $c_\pi \in \mathbb{C}$ .
  - Use parts (a) and (b) to prove the integral formula. That is, show that  $\tilde{\Pi}_n = \Pi_n$ .
3. **Haar measure:** There is a unique probability measure  $dU$  on the unitary operators  $\text{U}(\mathcal{X})$  that is invariant under  $U \mapsto VUW$  for every pair of unitaries  $V, W$ . It is called the *Haar measure*. Its defining property can be stated as follows: For every continuous function  $f$  on  $\text{U}(\mathcal{X})$  and for all unitaries  $V, W \in \text{U}(\mathcal{X})$ , it holds that  $\int f(U) dU = \int f(VUW) dU$ . Now let  $A \in L(\mathcal{X})$ .

- Argue that  $\int UAU^\dagger dU$  commutes with all unitaries.
- Deduce that  $\int UAU^\dagger dU = \frac{\text{Tr}[A]}{d} I$ , where  $d = \dim \mathcal{X}$ .

4. **De Finetti theorem and quantum physics (optional):** Given a Hermitian operator  $h$  on  $\mathbb{C}^d \otimes \mathbb{C}^d$ , consider the operator  $H = \frac{1}{n-1} \sum_{i \neq j} h_{i,j}$  on  $(\mathbb{C}^d)^{\otimes n}$ , where  $h_{i,j}$  acts by  $h$  on subsystems  $i$  and  $j$  and by the identity on the remaining subsystems (e.g.,  $h_{1,2} = h \otimes I^{\otimes(n-2)}$ ).

- Show that  $\frac{E_0}{n} \leq \frac{1}{n} \langle \psi^{\otimes n} | H | \psi^{\otimes n} \rangle = \langle \psi^{\otimes 2} | h | \psi^{\otimes 2} \rangle$  for every pure state  $\psi$  on  $\mathbb{C}^d$ .

Let  $E_0$  denote the smallest eigenvalue of  $H$  and  $|E_0\rangle$  a corresponding eigenvector. If the eigenspace is one-dimensional and  $n > d$  then  $|E_0\rangle \in \text{Sym}^n(\mathbb{C}^d)$  (you do not need to show this).

- Use the de Finetti theorem to show that  $\frac{E_0}{n} \approx \min_{\|\psi\|=1} \langle \psi^{\otimes 2} | h | \psi^{\otimes 2} \rangle$  for large  $n$ .

*Interpretation: The Hamiltonian  $H$  describes a mean-field system. Your result shows that in the thermodynamic limit the ground state energy density can be computed using states of form  $\psi^{\otimes n}$ .*