

Portfolio Theory

P.J.C. Spreij

this version: January 28, 2019

Contents

1	Valuation in a one-period model	1
1.1	Arbitrage	1
1.2	Contingent claims and derivatives	4
1.3	Complete markets	8
1.4	Exercises	9
2	Preferences	12
2.1	Preference relations	12
2.2	Numerical representations	13
2.3	Exercises	17
3	Lotteries	20
3.1	Von Neumann-Morgenstern representations	20
3.2	Exercises	25
4	Utility and expected utility	26
4.1	Risk aversion	26
4.2	Arrow-Pratt coefficient	29
4.3	Exercises	31
5	Stochastic dominance	33
5.1	Uniform order	33
5.2	Monotone order	37
5.3	Exercises	38
6	Portfolio optimization	40
6.1	Optimization and absence of arbitrage	40
6.2	Exponential utility and relative entropy	45
6.3	Exercises	47
7	Optimal contingent claims	50
7.1	An expected utility optimization problem	50
7.2	Optimization under uniform order restrictions	53
7.3	Exercises	57
8	Dynamic arbitrage theory	60
8.1	Self-financing trading strategies	60
8.2	Arbitrage	62
8.3	Proof of Lemma 8.11	67
8.4	European contingent claims	72
8.5	Complete markets	77
8.6	CRR model	79
8.7	Exercises	82

9	Optimization in dynamic models	85
9.1	Dynamic programming	85
9.2	Optimal portfolios via dynamic programming	89
9.3	Consumption-investment and the martingale method	91
9.4	Exercises	98
A	Appendix	100
A.1	Separating hyperplanes	100
A.2	Hahn-Banach theorem and some ramifications	101
A.3	Existence of essential supremum	104
A.4	Results on the weak topology for measures	105
A.5	Proofs of results of Section 9.1	106
A.6	Exercises	108

Preface

These lecture notes have been written for and during the course *Portfolio Theory* at the Universiteit van Amsterdam in fall 2007.

The aim of the course is to introduce the fundamental concepts that underly the problem of portfolio optimization. Needed for this is also an exposition of the fundamental notions of financial markets, such as absence of arbitrage and completeness. Other concepts that will be developed are preference relations and utility. All this will first be done for a one-period market and later on extended to markets with a larger horizon. In the latter case we show how to use Dynamic Programming for optimization problems. We confine ourselves to models in discrete time, although portfolio optimization in continuous time is a topic that equally well deserves a place in this set of lecture notes. This will be a topic of future consideration.

The lecture notes are for a large deal based on the book *Stochastic Finance, An Introduction in Discrete Time* by Alexander Schied and Hans Föllmer. But also other sources like *Introduction to Mathematical Finance* by Stanley R. Pliska have been consulted, as well as *Stochastic systems: estimation, identification and adaptive control* by P.R. Kumar and Pravin Varaiya.

Finally, these lectures notes will be updated and adapted for a next course. Many errors and omissions in earlier versions have been corrected, thanks to careful reading by Attila Herczegh, Demeter Kiss and Kamil Kosiński, who were among the first students that took this course. Later on, Johan du Plessis, Hailong Bao and Nicos Starreveld provided me with more very useful feedback that resulted in eliminating other errors. But it is almost inevitable that some remained. Readers are kindly requested to report any inaccuracies. Suggestions for improvements are equally welcome.

Amsterdam, January 2013

Peter Spreij

1 Valuation in a one-period model

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space. We will assume that all random variables that we encounter below are defined on this space and real valued. First we describe a market consisting of $d + 1$ assets. We assume that there are only two relevant time instants, $t = 0$ and $t = 1$. The assets are numbered from 0 to d . The zero-th asset is a non-risky asset. Given an interest rate $r > -1$ and a price π_0 of the asset at time $t = 0$, its price at $t = 1$ is given by $S_0 = \pi_0(1+r) > 0$. We will make the convention that $\pi_0 = 1$. The other d assets are risky. This means that their prices π_i at $t = 0$ are known deterministic nonnegative numbers, but their prices S_i at $t = 1$ are not exactly known at $t = 0$ and are modeled as nonnegative random variables. Next to prices, we also have quantities, which are for the i -th asset given by real constants ξ_i . We introduce the following vectors.

The price vector at $t = 0$ of these assets will be denoted by $\bar{\pi} = (\pi_0, \pi)$, where π denotes the vector of the d risky assets. At $t = 1$ we have with similar notation $\bar{S} = (S_0, S)$. Likewise we have for the quantities $\bar{\xi} = (\xi_0, \xi)$. The vector $\bar{\xi}$ will often be referred to as a *portfolio*. The *value* of the portfolio at $t = 0$ is then $W_0 = \bar{\pi} \cdot \bar{\xi}$ and at $t = 1$ it is $W_1 = \bar{\xi} \cdot \bar{S}$. The dot here denotes the ordinary inner product.

1.1 Arbitrage

In a realistic market there will not exist arbitrage opportunities, making a sure profit by investing in a portfolio. We give a formal definition of this.

Definition 1.1 A portfolio $\bar{\xi}$ is an arbitrage opportunity if $W_0 \leq 0$, $W_1 \geq 0$ a.s. and $\mathbb{P}(W_1 > 0) > 0$. A market is called arbitrage free, if arbitrage opportunities don't exist.

Remark 1.2 Consider an arbitrage free market, with $\pi_i = 0$ for some asset i . Take the portfolio consisting of 1 unit of asset i only. Then $W_1 = S_i$. Since this portfolio is not an arbitrage opportunity, we must have that $S_i = 0$ a.s. Hence this asset is always worthless, and therefore we exclude zero initial prices. All π_i will be assumed strictly positive.

It will turn out useful to characterize arbitrage opportunities in terms of the risky assets only.

Lemma 1.3 *The existence of an arbitrage opportunity is equivalent to the existence of a vector ξ having the properties $\xi \cdot S \geq (1+r)\xi \cdot \pi$ a.s. and $\mathbb{P}(\xi \cdot S > (1+r)\xi \cdot \pi) > 0$.*

Proof Let $\bar{\xi} = (\xi_0, \xi)$ be an arbitrage opportunity. Then

$$\xi \cdot S - (1+r)\xi \cdot \pi = W_1 - (1+r)W_0 \geq W_1.$$

Since also $W_1 \geq 0$ a.s. and $\mathbb{P}(W_1 > 0) > 0$, the characterization follows. Conversely, assume that the characterization holds true for some vector ξ . Choose $\xi_0 = -\xi \cdot \pi$. Then $W_0 = 0$ and $W_1 = \xi \cdot S - (1+r)\xi \cdot \pi$. It follows that $\bar{\xi}$ is an arbitrage opportunity. \square

We now proceed with another characterization of arbitrage opportunities. We need the vector Y of *discounted net gains*. Its elements Y_i ($i = 1, \dots, d$) are given by

$$Y_i = \frac{S_i}{1+r} - \pi_i.$$

Corollary 1.4 *Existence of an arbitrage opportunity is equivalent to the existence of a vector ξ such that $\xi \cdot Y \geq 0$ a.s and $\mathbb{P}(\xi \cdot Y > 0) > 0$. An arbitrage free market is characterized by the implication $\xi \cdot Y \geq 0$ a.s. $\Rightarrow \xi \cdot Y = 0$ a.s.*

Proof This is an immediate consequence of Lemma 1.3. \square

Definition 1.5 A probability measure \mathbb{P}^* on \mathcal{F} is called a risk-neutral measure, or a martingale measure, if

$$\pi_i = \mathbb{E}^* \frac{S_i}{1+r}, \quad i = 0, \dots, d.$$

It follows that \mathbb{P}^* on \mathcal{F} is a risk-neutral measure, iff $\mathbb{E}^* Y = 0$. Notice that $\mathbb{E}^* Y$ is well defined for any \mathbb{P}^* , since each Y_i is lower bounded by $-\pi_i$. By \mathcal{P} we denote the set of all risk-neutral measures that are equivalent to \mathbb{P} (they define the same null sets). Elements of \mathcal{P} are called *equivalent martingale measures*.

Theorem 1.6 below is a version of the (first) *Fundamental Theorem of Asset Pricing* (FTAP).

Theorem 1.6 *A market is free of arbitrage iff the set \mathcal{P} is nonempty. In this case there exists a $\mathbb{P}^* \in \mathcal{P}$ such that the Radon-Nikodym derivative $\frac{d\mathbb{P}^*}{d\mathbb{P}}$ is bounded.*

Proof Let \mathcal{P} be non-empty and take $\mathbb{P}^* \in \mathcal{P}$. Let ξ be such that $\xi \cdot Y \geq 0$ \mathbb{P} -a.s. Then the same is true under \mathbb{P}^* . Since $\mathbb{P}^* \in \mathcal{P}$, we have $\mathbb{E}^* \xi \cdot Y = 0$ and hence $\xi \cdot Y = 0$ \mathbb{P}^* -a.s., but then also under \mathbb{P} . The result follows from Corollary 1.4.

Conversely, assume that the market is arbitrage free. We have to show the existence of a $\mathbb{P}^* \sim \mathbb{P}$ such that $\mathbb{E}^* Y = 0$. We first assume that $\mathbb{E}|Y| < \infty$. Put

$$\mathcal{Q} = \{\mathbb{Q} : \mathbb{Q} \sim \mathbb{P} \text{ and } \frac{d\mathbb{Q}}{d\mathbb{P}} \text{ bounded}\},$$

and

$$\mathcal{C} = \{\mathbb{E}_{\mathbb{Q}} Y : \mathbb{Q} \in \mathcal{Q}\}.$$

Notice that \mathcal{C} is well defined, since $\mathbb{E}_{\mathbb{Q}} |Y| < \infty$ for all $\mathbb{Q} \in \mathcal{Q}$. One easily shows that \mathcal{Q} is a convex set, and hence \mathcal{C} is a convex subset of \mathbb{R}^d . For any \mathbb{P}^* one has $\mathbb{E}^* Y = 0$, and we therefore show that $0 \in \mathcal{C}$.

Assume the contrary, $0 \notin \mathcal{C}$. By virtue of the *Separating Hyperplane Theorem*, Theorem A.1, there exists a vector $\xi \in \mathbb{R}^d$ such that $\xi \cdot x \geq 0$ for all $x \in \mathcal{C}$ and $\xi \cdot x_0 > 0$ for some $x_0 \in \mathcal{C}$. But elements of \mathcal{C} are expectations, so we have $\mathbb{E}_{\mathbb{Q}} \xi \cdot Y \geq 0$ for all $\mathbb{Q} \in \mathcal{Q}$ and $\mathbb{E}_{\mathbb{Q}_0} \xi \cdot Y > 0$ for some $\mathbb{Q}_0 \in \mathcal{Q}$. Since the latter expectation is strictly positive, we must have $\mathbb{Q}_0(\xi \cdot Y > 0) > 0$, and by equivalence we then also have $\mathbb{P}(\xi \cdot Y > 0) > 0$. If we can show that $\xi \cdot Y \geq 0$ a.s., then we have shown existence of an arbitrage opportunity in view of Lemma 1.3, a contradiction. Consider thereto the set $A = \{\xi \cdot Y < 0\}$. The following arguments are aimed at showing $\mathbb{P}(A) = 0$.

Define for $n \geq 2$ the functions ϕ_n by

$$\phi_n = \left(1 - \frac{1}{n}\right) \mathbf{1}_A + \frac{1}{n} \mathbf{1}_{A^c}.$$

Notice that $\mathbb{E} \phi_n \geq \mathbb{P}(A^c)/n > 0$. We can therefore define the probability measures \mathbb{Q}_n by $\frac{d\mathbb{Q}_n}{d\mathbb{P}} = c_n \phi_n$, where $c_n = 1/\mathbb{E} \phi_n$. Note also that $\frac{1}{n} \leq \phi_n \leq 1 - \frac{1}{n}$ as $n \geq 2$, from which we obtain that the \mathbb{Q}_n are in \mathcal{Q} as now $\frac{d\mathbb{Q}_n}{d\mathbb{P}} \leq n - 1$. Then we have the following string of equalities (we also use the Dominated Convergence Theorem)

$$\begin{aligned} \mathbb{E}(\xi \cdot Y \mathbf{1}_{\{\xi \cdot Y < 0\}}) &= \mathbb{E}(\xi \cdot Y \mathbf{1}_A) \\ &= \mathbb{E}(\xi \cdot Y \lim \phi_n) \\ &= \lim \mathbb{E}(\xi \cdot Y \phi_n) \\ &= \lim \frac{1}{c_n} \mathbb{E}_{\mathbb{Q}_n}(\xi \cdot Y) \geq 0. \end{aligned}$$

So the nonpositive random variable $\xi \cdot Y \mathbf{1}_{\{\xi \cdot Y < 0\}}$ has a nonnegative expectation. This implies $\mathbb{P}(\xi \cdot Y \geq 0) = 1$, which we wanted to prove.

The case $\mathbb{E}|Y| = \infty$ is left as Exercise 1.3. Note that the assertion on the existence of a bounded Radon-Nikodym derivative follows by the definition of the set \mathcal{C} . \square

The following example shows that in an arbitrage free market, risk neutral measures are in general not unique.

Example 1.7 Suppose that $\Omega = \{\omega_1, \dots, \omega_n\}$ with $n \geq 2$. Assume that $p_i = \mathbb{P}(\{\omega_i\}) > 0$ for all i , there is only one risky asset S_1 and $s_i = S_1(\omega_i) > 0$ for all i . Assume $s_1 < \dots < s_n$. Theorem 1.6 says that there is no arbitrage iff $(1+r)\pi_1 \in (s_1, s_n)$ and that for every \mathbb{P}^* , represented by a probability vector (p_1^*, \dots, p_n^*) , the p_i^* solve $\sum_i p_i^* s_i = (1+r)\pi_1$. Moreover, the risk-neutral measure is unique iff $n = 2$.

Definition 1.8 The set of attainable pay-offs is $\mathcal{W} = \{\bar{\xi} \cdot \bar{S} : \bar{\xi} \in \mathbb{R}^{d+1}\}$.

Lemma 1.9 Assume that the market is arbitrage free. Suppose that $W \in \mathcal{W}$ can be represented both as $\bar{\xi} \cdot \bar{S}$ and as $\bar{\zeta} \cdot \bar{S}$. Then the values of the two portfolios at $t = 0$ are equal, $\bar{\xi} \cdot \bar{\pi} = \bar{\zeta} \cdot \bar{\pi}$.

Proof Any $\mathbb{P}^* \in \mathcal{P}$ (which is non-empty) satisfies $\mathbb{E}^* \bar{S} = (1+r)\bar{\pi}$. Hence $\mathbb{E}^* W = \mathbb{E}^*(\xi \cdot \bar{S}) = \mathbb{E}^*(\bar{\zeta} \cdot \bar{S})$. The result follows. \square

The above lemma yields *the principle, or law, of one price*. The price at $t = 0$ of an attainable pay-off W is equal to and defined by $\frac{\mathbb{E}^* W}{1+r}$, for any $\mathbb{P}^* \in \mathcal{P}$. This can be rephrased by saying that two portfolios which generate the same pay-off at $t = 1$ must have the same price at $t = 0$, which is then given by the expectation under any of the risk-neutral measures. Check that if the initial price vector $\bar{\pi}$ is different from $\frac{\mathbb{E}^* \bar{S}}{1+r}$, an arbitrage opportunity can explicitly be constructed.

1.2 Contingent claims and derivatives

There are many financial products other than portfolios (whose pay-off is attainable by definition). Some of these depend on the underlying risky assets, like call options. An example is the European call option whose pay-off is $C := (S_1 - K)^+$ (the constant K is called the strike price). We see that this pay-off is a function of S_1 . We will also consider pay-offs that are (in principle) not functions of \bar{S} .

Definition 1.10 A contingent claim C is by definition a nonnegative random variable (so C is \mathcal{F} -measurable). Such a C is called a derivative if C is $\sigma(\bar{S}) = \sigma(S)$ -measurable.

In this definition we require nonnegativity to ensure existence of an expectation. Alternatives to this are conceivable, like C lower bounded, or $\mathbb{E}|C| < \infty$. Another reason for nonnegativity is that we now treat C similar to the given assets, which are always assumed to have a nonnegative payoff.

Since we have seen how to price portfolios in arbitrage free markets (at $t = 0$), the natural question is how to price contingent claims. By analogy, an obvious candidate is $\frac{\mathbb{E}^* C}{1+r}$. It turns out that this is true, but also that in general this price is not unique and that, unlike for attainable pay-offs, it depends on the specific choice of the risk neutral measure.

Since absence of arbitrage is the key to finding a pricing rule for a contingent claim, we will look at the *extended market*. Next to the assets we already have, we consider the extra security $S_{d+1} = C$ and its price π_{d+1} at $t = 0$. No arbitrage considerations in the extended market will give the possible values of π_{d+1} .

Definition 1.11 A (nonnegative) real number π^C is called an arbitrage free price of the contingent claim C if the extended market with $\pi_{d+1} = \pi^C$ is free of arbitrage. We denote by $\Pi(C)$ the set of all prices π^C .

The next theorem confirms the conjecture made at the beginning of this section and gives a precise formulation.

Theorem 1.12 *Let C be a contingent claim. Suppose that the original market is arbitrage free (so $\mathcal{P} \neq \emptyset$). Then also $\Pi(C)$ is non-empty and in fact, one has*

$$(1.1) \quad \Pi(C) = \left\{ \mathbb{E}^* \frac{C}{1+r} : \mathbb{P}^* \in \mathcal{P} \text{ such that } \mathbb{E}^* C < \infty \right\}.$$

Proof We first show that the set on the right hand side of (1.1) is not empty. Call this set E . Without loss of generality we assume that $C \geq 0$. Introduce a probability measure $\tilde{\mathbb{P}}$ by $d\tilde{\mathbb{P}} = \frac{c}{1+C} d\mathbb{P}$, where c is the normalizing constant. One sees that $\tilde{\mathbb{P}} \sim \mathbb{P}$ and that $\tilde{\mathbb{E}} C < \infty$. From the equivalence it follows that the market is also free of arbitrage under $\tilde{\mathbb{P}}$. Then Theorem 1.6 yields the existence of a risk-neutral measure \mathbb{P}^* equivalent to $\tilde{\mathbb{P}}$ (and then also to \mathbb{P}) with $\frac{d\mathbb{P}^*}{d\tilde{\mathbb{P}}}$ bounded, by B say. Then we have $\mathbb{E}^* C = \tilde{\mathbb{E}} \left(\frac{d\mathbb{P}^*}{d\tilde{\mathbb{P}}} C \right) \leq B \tilde{\mathbb{E}} C < \infty$. Hence the number $\frac{\mathbb{E}^* C}{1+r}$ belongs to the set E , which is thus not empty.

We now show that $\Pi(C) \subset E$. If $\Pi(C) = \emptyset$, there is nothing to prove. Therefore we assume that we can pick $\pi^C \in \Pi(C)$. By definition of the arbitrage price, the extended market is free of arbitrage, so in view of Theorem 1.6, there exists a probability measure $\hat{\mathbb{P}}$ equivalent to \mathbb{P} such that $\hat{\mathbb{E}} \frac{S_i}{1+r} = \pi_i$, for all $i = 0, \dots, d+1$. In particular we have that $\hat{\mathbb{E}} \frac{C}{1+r} = \pi^C < \infty$ (take $i = d+1$). Since $\hat{\mathbb{P}}$ is then also a risk-neutral measure for the original market, we have $\Pi(C) \subset E$.

To show the reversed inclusion, we take $\mathbb{P}^* \in \mathcal{P}$ and define $\pi_{d+1} = \pi^C := \frac{\mathbb{E}^* C}{1+r}$. This definition turns \mathbb{P}^* into a risk-neutral measure for the extended market as well, and so we have $E \subset \Pi(C)$. \square

In the proof of Theorem 1.15 the concept of *non-redundancy* comes in handy.

Definition 1.13 A market is called *non-redundant* if the implication $\bar{\xi} \cdot \bar{S} = 0$ \mathbb{P} -a.s. $\Rightarrow \bar{\xi} = 0$ holds.

If the above implication doesn't hold for some portfolio $\bar{\xi}$, then it has a nonzero element, ξ_i say. But then we also have $S_i = -\frac{1}{\xi_i} \sum_{j \neq i} \xi_j S_j$. So, the i -th asset is a linear combination of the other ones, a form of redundancy. In an arbitrage free market we then also have (take expectations under any risk-neutral measure) $\pi_i = -\frac{1}{\xi_i} \sum_{j \neq i} \xi_j \pi_j$, again a linear combination and with the same coefficients, and likewise Y_i is the same linear combination of the Y_j .

Proposition 1.14 *The following holds.*

- (i) *Any (finite) market can be reduced to a non-redundant market in the sense that there exists a non-redundant market such that for any portfolio of assets in the original market one can find a portfolio in the non-redundant market with exactly the same payoff.*
- (ii) *In a non-redundant market the implication $\xi \cdot Y = 0$ \mathbb{P} -a.s. $\implies \xi = 0$ holds. Conversely, if this implication holds and the market is arbitrage free, then the market is also non-redundant.*

Proof Exercise 1.5. \square

We will see later that $\Pi(C)$ is an interval. We then need the lower and upper limits. This motivates to study $\inf \Pi(C)$ and $\sup \Pi(C)$. Of course these quantities are of independent interest. It gives upper and lower bounds for the possible arbitrage free prices of a contingent claim C .

Theorem 1.15 *Assume that the market is arbitrage free. Let*

$$M_0 = \{m \in [0, \infty] : \exists \xi \in \mathbb{R}^d \text{ with } m + \xi \cdot Y \leq \frac{C}{1+r} \mathbb{P}\text{-a.s.}\}$$

and

$$M_1 = \{m \in [0, \infty] : \exists \xi \in \mathbb{R}^d \text{ with } m + \xi \cdot Y \geq \frac{C}{1+r} \mathbb{P}\text{-a.s.}\}.$$

Then $\inf \Pi(C) = \max M_0$ and $\sup \Pi(C) = \min M_1$.

Proof We only give the proof of the characterization of the supremum. The other assertion follows by similar arguments. We first notice that $M_1 \neq \emptyset$, since $\infty \in M_1$ (take $\xi = 0$). Take $m \in M_1$ and $\xi \in \mathbb{R}^d$ such that $m + \xi \cdot Y \geq \frac{C}{1+r}$ \mathbb{P} -a.s. For any $\mathbb{P}^* \in \mathcal{P}$ we then have $m \geq \frac{\mathbb{E}^* C}{1+r}$. Taking the supremum on the right hand side over all \mathbb{P}^* and then the infimum on the left hand side over all $m \in M_1$ yields $\inf M_1 \geq \sup \Pi(C)$ in view of Theorem 1.12. We proceed by showing the reversed inequality. Since this is trivial if $\sup \Pi(C) = \infty$, we assume that $\sup \Pi(C) < \infty$. Pick $m > \sup \Pi(C)$. If we can show that $m \geq \inf M_1$, we are done by taking the limit $m \downarrow \sup \Pi(C)$. Since $m \notin \Pi(C)$, there exists an arbitrage opportunity in the extended market with $\pi_{d+1} = m$ and $S_{d+1} = C$. From Corollary 1.4, applied to the extended market, we obtain the existence of a vector $\xi \in \mathbb{R}^d$ and a real number ξ_{d+1} with the properties that

$$(1.2) \quad \xi \cdot Y + \xi_{d+1} \left(\frac{C}{1+r} - m \right) \geq 0 \quad \mathbb{P}\text{-a.s.}$$

and strictly positive with positive \mathbb{P} -probability. Since the original market is arbitrage free, we can take $\mathbb{P}^* \in \mathcal{P}$ under which $\xi \cdot Y + \xi_{d+1} \left(\frac{C}{1+r} - m \right)$ has strictly positive expectation $\mathbb{E}^* \left(\xi_{d+1} \left(\frac{C}{1+r} - m \right) \right)$. Because $m > \sup \Pi(C) \geq \mathbb{E}^* \frac{C}{1+r}$, we find that $\xi_{d+1} < 0$. Consider the portfolio $\zeta = -\frac{\xi}{\xi_{d+1}}$. From (1.2) we obtain that $m + \zeta \cdot Y \geq \frac{C}{1+r}$. So $m \in M_1$ and thus $m \geq \inf M_1$, which we wanted to show.

The last thing to do is to show that the infimum is attained. If $\inf M_1 = \infty$, this is trivial. So, we assume that $\inf M_1 < \infty$. Choose a sequence of $m^n \in M_1$ that decreases to $\inf M_1$ and pick the corresponding ξ^n . We then have

$$(1.3) \quad m^n + \xi^n \cdot Y \geq \frac{C}{1+r} \mathbb{P}\text{-a.s.}$$

If we would know that the sequence (ξ^n) had a finite limit ξ , we could take limits in (1.3) to get $m + \xi \cdot Y \geq \frac{C}{1+r}$, which case it follows that $\inf M_1 \in M_1$ and is thus attained. In general, the existence of a limit ξ cannot be guaranteed,

but in the sequel we show that we can apply the above arguments along some subsequence.

Without loss of generality, we may assume that the market is non-redundant. Otherwise, we could replace the $\xi^n \cdot Y$ by a linear combination of non-redundant discounted net gains. Suppose that $\liminf \|\xi^n\| = \infty$. Then the vectors $\eta^n := \frac{\xi^n}{\|\xi^n\|}$ all lie on the (compact) unit circle and therefore converge along a subsequence (η^{n_k}) to some vector η . Divide the inequality (1.3) by $\|\xi^{n_k}\|$ and take the limit for $k \rightarrow \infty$ to obtain $\eta \cdot Y \geq 0$ \mathbb{P} -a.s. Absence of arbitrage entails $\eta \cdot Y = 0$ \mathbb{P} -a.s. and non-redundancy then yields $\eta = 0$ (see Proposition 1.14). This contradicts $\|\eta\| = 1$. Hence $\liminf \|\xi^n\| < \infty$. Choose then a subsequence (again denoted by) (ξ^{n_k}) converging to a finite limit ξ . Take along the same subsequence limits in (1.3) to arrive at $\inf M_1 + \xi \cdot Y \geq \frac{C}{1+r}$ \mathbb{P} -a.s. This shows that $\inf M_1 \in M_1$. \square

Remark 1.16 The sets M_0 and M_1 in Theorem 1.15 are of interest in their own rights. One can show that the set M_1 coincides with the set of prices (at $t = 0$) of portfolios that *superhedge* the claim C , where a portfolio $\bar{\xi}$ superhedges C if $\bar{\xi} \cdot \bar{S} \geq C$ a.s. A similar characterization can be given for M_0 . See also Exercise 1.2.

Definition 1.17 A contingent claim C is called *attainable* if C a.s. belongs to the space of attainable pay-offs \mathcal{W} , so $C = \bar{\xi} \cdot \bar{S}$ for some portfolio $\bar{\xi}$. Such a portfolio is called *replicating* portfolio, or *hedge*.

It is obvious that in an arbitrage free market, the arbitrage-free price of an attainable claim equals that of the replicating portfolio and is thus unique, due to the law of one price.

Proposition 1.18 *Let C be a contingent claim in an arbitrage free market. Then*

- (i) C is attainable iff it admits a unique arbitrage-free price.
- (ii) If C is not attainable, $\Pi(C)$ is the open interval $(\inf \Pi(C), \sup \Pi(C))$.

Proof (i) If C admits a unique price, the set $\Pi(C)$ is not an open interval, so (i) follows from (ii). We now prove the latter assertion. Let C be not attainable. From Theorem 1.12 it follows that $\Pi(C)$ is a non-empty convex subset of \mathbb{R} , and so it is an interval. We only have to show that it is open, which happens if both $\inf \Pi(C)$ and $\sup \Pi(C)$ are not contained in it. We only consider $\inf \Pi(C)$ and suppose that $\inf \Pi(C) \in \Pi(C)$. Theorem 1.15 then implies that there exists $\xi \in \mathbb{R}^d$ such that $\inf \Pi(C) + \xi \cdot Y \leq \frac{C}{1+r}$ \mathbb{P} -a.s. Since C is not attainable we have $\mathbb{P}(\inf \Pi(C) + \xi \cdot Y < \frac{C}{1+r}) > 0$. Extend the original market with the risky asset having pay-off C endowed with the price $\inf \Pi(C)$ and consider the risky portfolio $-\xi$ augmented with one unit of the extra risky asset. The total net discounted gain of this extended portfolio is equal to $-\xi \cdot Y + \frac{C}{1+r} - \inf \Pi(C)$. The above inequalities show that we have constructed an arbitrage opportunity in the extended market. Hence $\inf \Pi(C)$ is not an arbitrage free price for C and thus not an element of $\Pi(C)$. \square

1.3 Complete markets

In arbitrage free *complete* markets, as we shall see, every contingent claim has a unique price. We start with a definition.

Definition 1.19 A market is called *complete* if every contingent claim is attainable.

Given a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, we denote for $p \geq 1$ by $L^p(\Omega, \mathcal{F}, \mathbb{P})$ the usual Banach space with the L^p -norm $\|\cdot\|_p$ defined by $\|X\|_p = (\mathbb{E}|X|^p)^{1/p}$. For $p = \infty$ we have $\|X\|_\infty = \inf\{c \geq 0 : \mathbb{P}(|X| > c) = 0\}$. By $L^0(\Omega, \mathcal{F}, \mathbb{P})$ we simply denote the linear space of a.s. finitely valued random variables. It can be considered as a (complete) *metric* space by defining the metric ρ by $\rho(X, Y) = \mathbb{E} \frac{|X-Y|}{1+|X-Y|}$. It then holds that $X_n \xrightarrow{\rho} X$ iff $X_n \xrightarrow{\mathbb{P}} X$. We will often use the same notation $L^p(\Omega, \mathcal{F}, \mathbb{P})$ when its elements are finite-dimensional random vectors $X = (X_1, \dots, X_k)$ with $|X(\omega)|$ denoting any norm of the vector $X(\omega)$, usually $|X(\omega)| = (\sum_{j=1}^k X_j(\omega)^2)^{1/2}$.

In every market \mathcal{W} is a subset of the set of \mathcal{F} -measurable random variables. If \mathcal{C} is the set of attainable claims, we have the inclusions $\mathcal{C} \subset \mathcal{W} \subset L^0(\Omega, \mathcal{F}, \mathbb{P})$. In a complete market we have $L^0(\Omega, \mathcal{F}, \mathbb{P}) = \mathcal{C} + (-\mathcal{C}) \subset \mathcal{W}$ and so $L^0(\Omega, \mathcal{F}, \mathbb{P}) = \mathcal{W}$. It follows that in this case $L^0(\Omega, \mathcal{F}, \mathbb{P})$ is finite dimensional, with dimension less than or equal to $d + 1$ and also that \mathcal{F} and $\sigma(S)$ are identical modulo null sets.

Recall the definition of an atom. A set $A \in \mathcal{F}$ is called an atom of $(\Omega, \mathcal{F}, \mathbb{P})$ if $\mathbb{P}(A) > 0$ and every measurable subset B of A either has probability zero or $\mathbb{P}(A)$.

Proposition 1.20 Let \mathcal{N} be the set of integers m for which there exists a measurable partition of Ω into m sets with positive probability. For any $p \in [0, \infty]$ one has $\dim L^p(\Omega, \mathcal{F}, \mathbb{P}) = \sup \mathcal{N}$. Moreover, $n := \dim L^p(\Omega, \mathcal{F}, \mathbb{P}) < \infty$ iff there exists a partition of Ω into n atoms.

Proof Suppose that there exists a measurable partition of m sets with positive probability, A_1, \dots, A_m say. Then the set of corresponding indicators $\mathbf{1}_{A_1}, \dots, \mathbf{1}_{A_m}$ is linearly independent in any $L^p(\Omega, \mathcal{F}, \mathbb{P})$. Therefore $n \geq m$. If \mathcal{N} is unbounded, we trivially have $n = \sup \mathcal{N}$. Assume therefore that $n_0 := \sup \mathcal{N} < \infty$. Since $n_0 \in \mathcal{N}$ there exists a partition A_1, \dots, A_{n_0} of *atoms*. Indeed, if one of the A_i were not an atom, we split it into more sets of positive probability, but then n_0 would not be maximal. Hence we have that $\mathcal{F} = \sigma(A_1, \dots, A_{n_0})$ modulo null sets and every \mathcal{F} -measurable function is almost surely constant on the A_i and hence a linear combination of the linearly independent $\mathbf{1}_{A_i}$. It follows that $n = n_0$. \square

The following theorem is known as the second Fundamental Theorem of Asset Pricing.

Theorem 1.21 An arbitrage free market is complete iff there exists exactly one risk-neutral measure. In this case $\dim L^0(\Omega, \mathcal{F}, \mathbb{P}) \leq d + 1$.

Proof Let the market be complete. Consider the claim $C = \mathbf{1}_A(1+r)$ for some $A \in \mathcal{F}$. Then C is attainable and thus admits the unique fair price $\mathbb{P}^*(A)$, by virtue of Proposition 1.18, valid for any $\mathbb{P}^* \in \mathcal{P}$. But since A is arbitrary, this shows that \mathbb{P}^* is unique.

Conversely, assume that $\mathcal{P} = \{\mathbb{P}^*\}$. Let C be a *bounded* contingent claim. Then its arbitrage free price is $\frac{\mathbb{E}^*C}{1+r} < \infty$. Proposition 1.18 says that C is attainable. But then we have that $L^\infty(\Omega, \mathcal{F}, \mathbb{P}) \subset \mathcal{W}$, and so $\dim L^\infty(\Omega, \mathcal{F}, \mathbb{P}) \leq d+1$, since $\dim \mathcal{W} \leq d+1$. Proposition 1.20 then implies that $(\Omega, \mathcal{F}, \mathbb{P})$ has at most $d+1$ atoms. It follows that every contingent claim must be bounded, and by the above reasoning, it is attainable as well. \square

It follows from Theorem 1.21 that for a complete market with finitely many securities we can as well set up a model with a finite space Ω . Let us take an example with n atoms, $\Omega = \{\omega_1, \dots, \omega_n\}$. Let there be one risky asset with initial price π and S as its price at $t = 1$. Let $s_1 = S(\omega_1) < \dots < s_n = S(\omega_n)$. We have seen before (Example 1.7) that any risk-neutral measure \mathbb{P}^* , represented by a vector (p_1, \dots, p_n) on the simplex, must satisfy $\sum_i p_i s_i = (1+r)\pi \in (s_1, s_n)$. It is obvious that \mathbb{P}^* is unique iff $n = 2$, which is then the only case where the market is complete, in view of Theorem 1.21.

Moreover, in this case one easily computes $p_1 = \frac{s_2 - (1+r)\pi}{s_2 - s_1}$ and $p_2 = \frac{(1+r)\pi - s_1}{s_2 - s_1}$. It is also straightforward to compute the replicating portfolio for a given contingent claim C . Let $c_i = C(\omega_i), i = 1, 2$. If the claim is attainable one should find ξ_0 and ξ_1 such that $c_i = \xi_0(1+r) + \xi_1 s_i, i = 1, 2$. This linear system of equations is solved by

$$\begin{aligned}\xi_0 &= \frac{c_1 s_2 - c_2 s_1}{(s_2 - s_1)(1+r)} \\ \xi_1 &= \frac{c_2 - c_1}{s_2 - s_1}.\end{aligned}$$

One also easily computes an explicit expression for $\pi^C = \xi_0 + \frac{\xi_1 \mathbb{E}^* S}{1+r}$. Furthermore, here we have $\dim L^p(\Omega, \mathcal{F}, \mathbb{P}) = d+1 = 2$.

1.4 Exercises

1.1 Consider a financial market as in Example 1.7 and assume $N = 3$.

- (a) Characterize explicitly the set \mathcal{P} as a set of vectors (p_1^*, p_2^*, p_3^*) .
- (b) Consider a call option $C = (S_1 - K)^+$ for some $K > 0$. The set $\Pi(C)$ will turn out to be an interval. What are the upper and lower limits?

1.2 Let C be a contingent claim in an arbitrage free market with discounted net gains vector Y . Consider the set M_1 of Theorem 1.15. Put $\pi^* := \inf M_1$. Show that π^* is the lowest price of all portfolios ξ that are such that $\xi \cdot \bar{S} \geq C$ a.s.

1.3 Finish the proof of Theorem 1.6, by considering the case $\mathbb{E}|Y| = \infty$. *Hint:* Consider the auxiliary probability measure $\tilde{\mathbb{P}}$ defined by $\frac{d\tilde{\mathbb{P}}}{d\mathbb{P}} = \frac{c}{1+|Y|}$, with c the normalizing constant, and reason as in the proof of Theorem 1.12

1.4 Write $\pi(W) = \bar{\pi} \cdot \bar{\xi}$ for $W = \bar{\xi} \cdot \bar{S}$. If $\pi(W) \neq 0$ we define the *return* of W by $R(W) := \frac{W - \pi(W)}{\pi(W)}$.

- (a) Let $W = S_0$, so $\xi_0 = 1$ and the other ξ_i are zero. Compute $R(W)$.
- (b) Assume an arbitrage free market. Show that $\mathbb{E}^*(W) = (1+r)\pi(W)$, for any $W \in \mathcal{W}$. Compute, for every $W \in \mathcal{W}$ with $\pi(W) \neq 0$, the expected return $\mathbb{E}^*R(W)$.

1.5 Here you prove Proposition 1.14.

- (a) To prove the first statement you may proceed along the following pattern. Let $N = \{\bar{\xi} \in \mathbb{R}^{d+1} : \bar{\xi} \cdot \bar{S} = 0\}$ and assume that $\dim N > 0$. Let N^\perp be the orthogonal complement of N in \mathbb{R}^{d+1} with $\dim N^\perp = k+1 < d+1$. Show that there exists a matrix $B \in \mathbb{R}^{(d+1) \times (k+1)}$ of full column rank and a $(k+1)$ -dimensional random vector \bar{S}' such that $\bar{S} = B\bar{S}'$.
- (b) Show that any portfolio of assets in the original market has a counterpart, a portfolio of assets in \bar{S}' with the same payoff.
- (c) Show that the $(k+1)$ -dimensional market described by \bar{S}' is non-redundant. Does it have a riskless asset?

1.6 Consider a market defined on a finite probability space (as in Example 1.7) with 1 riskless and d (instead of one) risky assets. Find a relation between n and d if the market is non-redundant and complete. Show that this condition is not sufficient and provide a sufficient condition for non-redundancy and completeness.

1.7 Consider next to the given market the alternative market with the original non-risky asset and d risky assets whose values at $t = 1$ are represented by the vector of discounted net gains Y . Assume that at $t = 0$ the non-risky asset has price $\pi'_0 = \pi_0 = 1$ and the alternative risky assets have price vector π' equal to zero.

- (a) Show that any portfolio in the original market has a counterpart in the alternative market with same pay-off at $t = 1$.
- (b) Show that the alternative market is arbitrage free iff the original market is arbitrage free.

1.8 Assume that a market admits an arbitrage opportunity. Show there exists a $\bar{\xi} \in \mathbb{R}^{d+1}$ such that $W_0 = 0$ a.s., $W_1 \geq 0$ a.s., and $\mathbb{P}(W_1 > 0) > 0$. [Conversely, such a $\bar{\xi}$ is an arbitrage opportunity in the sense of Definition 1.1.]

1.9 Suppose that the initial price vector $\bar{\pi}$ is different from $\mathbb{E}^* \frac{\bar{S}}{1+r}$. Construct an arbitrage opportunity.

1.10 Consider the sets M_0 and M_1 of Theorem 1.15. Show by a direct argument that $\inf M_1 \geq \sup M_0$.

1.11 Complete the proof of Theorem 1.15, i.e. show that $\inf \Pi(C) = \max M_0$.

1.12 Complete the proof of Proposition 1.18, i.e. show that $\sup \Pi(C) \notin \Pi(C)$.

1.13 Consider a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and assume Ω has three elements, $\Omega = \{\omega_1, \omega_2, \omega_3\}$. Suppose there is, next to the riskless S_0 , one risky asset S_1 that is not constant. Construct a non-attainable claim C and show that the market extended with C is complete.

1.14 Consider a market, defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, consisting of d risky assets and one riskless asset. Assume that it is arbitrage free, complete and non-redundant. Show that $\dim L^p(\Omega, \mathcal{F}, \mathbb{P}) = d + 1$ (for any $p \in [0, \infty]$). Conversely, if $\dim L^p(\Omega, \mathcal{F}, \mathbb{P}) = d + 1$, show that the market is arbitrage free, complete and non-redundant. [This is a new exercise, not sure it is correct.]

1.15 The proof of the first part of Theorem A.1 says "one easily shows that $\lambda \geq 1$ ". You may have noticed that in the text before that, convexity of C is not used, and this is what is needed to establish $\lambda \geq 1$. Define $x_t = tx + (1-t)x_0$, $t \in [0, 1]$. For which t is $\|x_t\|$ minimal? Exploit this to prove the assertion.

2 Preferences

In a market commodities are traded and an agent acting in this market will have certain preferences of some commodities over others. Think that he likes apples more than pears. Preferences will be made explicit by introducing *preference relations*. Traded commodities include risky assets as well, or contingent claims. Future pay-offs of these products are uncertain, random. We have seen in the previous section, that in a complete market such claims have a unique (arbitrage free) price, computed as an expectation under the unique risk-neutral measure. In incomplete markets, there is usually an interval of possible arbitrage free prices. To select one of these, preference relations or their numerical counterparts, *utility functions*, are instrumental. Utility functions also describe the attitude of an economic agent towards *risk* (incurred by the uncertain pay-offs). In this case, the resulting price of a contingent claim depends on how risk is quantified and for different utility functions, usually, different prices will be the result. Utility functions can also be used in a complete market to choose between two portfolios that have the same price.

2.1 Preference relations

Let \mathcal{X} be a non-empty set representing commodities or securities, or in general possible choices an economic agent can make. Recall that a binary relation R on \mathcal{X} can be represented as a subset of $\mathcal{X} \times \mathcal{X}$ and that xRy means $(x, y) \in R$.

Definition 2.1 A (*strict*) *preference relation* or *preference order* on \mathcal{X} is a binary relation \succ with the properties

- (i) Asymmetry: If $x \succ y$, then $y \not\succeq x$.
- (ii) Negative transitivity: If $x \succ y$ and $z \in \mathcal{X}$, then $x \succ z$ or $z \succ y$.

Definition 2.2 A *weak preference relation* on \mathcal{X} is a binary relation \succeq with the properties

- (i) Completeness: For all $x, y \in \mathcal{X}$ one has $x \succeq y$ or $y \succeq x$.
- (ii) Transitivity: If $x \succeq y$ and $y \succeq z$, then $x \succeq z$.

Proposition 2.3 *Strict and weak preference relations are connected in the following way. If \succ is a strict preference relation on \mathcal{X} , then $x \succeq y$ defined by $y \not\succeq x$ yields a weak preference relation on \mathcal{X} . Conversely, if \succeq is a weak preference relation, then $x \succ y$ defined by $y \not\succeq x$ yields a strict preference relation.*

Proof Exercise 2.1. □

When dealing with strict and weak preference relations \succ and \succeq , we will always assume that they are related as in Proposition 2.3. Given a weak preference relation \succeq , an *indifference relation* \sim is defined by $x \sim y$ iff $x \succeq y$ and $y \succeq x$. One easily verifies that an indifference relation is an equivalence relation. Note that we also have $x \succ y$ iff $x \succeq y$ and $x \not\sim y$ (see also Exercise 2.6).

Sometimes it is notationally convenient to use reverse preference relations. So, instead of $x \succ y$ we also write $y \prec x$ and likewise we use $y \preceq x$ for $x \succeq y$.

2.2 Numerical representations

In a rather general setting abstract preference orders can be replaced with equivalent *numerical representations* for which the usual order on \mathbb{R} can be used.

Definition 2.4 A function $U : \mathcal{X} \rightarrow \mathbb{R}$ is called a *numerical representation* of a preference relation \succ , if $x \succ y$ is equivalent to $U(x) > U(y)$.

An alternative definition of a numerical representation is obtained by putting

$$(2.1) \quad x \succeq y \text{ iff } U(x) \geq U(y)$$

instead of the equivalence in Definition 2.4. Of course, any strictly increasing transformation of a numerical representation U yields another numerical representation. Numerical representations having additional properties are often called *utility functions*. We'll come back to this in Section 4.

Definition 2.5 Given a preference relation \succ on \mathcal{X} , a subset \mathcal{Z} of \mathcal{X} is called *order dense* (in \mathcal{X}) if for all $x, y \in \mathcal{X}$ with $x \succ y$, there exists a $z \in \mathcal{Z}$ such that $x \succeq z \succeq y$.

Theorem 2.6 A given preference relation \succ on \mathcal{X} admits a numerical representation iff \mathcal{X} contains a countable order dense subset.

Proof Let \mathcal{Z} be a countable order dense subset of \mathcal{X} . Choose a probability measure μ on \mathcal{Z} with $\mu(z) = \mu(\{z\}) > 0$ for all $z \in \mathcal{Z}$. Then we put

$$(2.2) \quad U(x) := \sum_{z: x \succ z} \mu(z) - \sum_{z: z \succ x} \mu(z).$$

Notice that existence of such a probability distribution is guaranteed by countability of \mathcal{Z} , that also makes $U(x)$ well defined in terms of the given summations. By construction we have $x \succ y$ iff $U(x) > U(y)$. First we compute

$$U(x) - U(y) = \sum_{z: x \succ z \succeq y} \mu(z) + \sum_{z: x \succeq z \succ y} \mu(z).$$

If $x \succ y$, then there is $z_0 \in \mathcal{Z}$ such that $x \succeq z_0 \succeq y$. By negative transitivity we also have $z_0 \succ y$ or $x \succ z_0$. Hence we have $x \succ z_0 \succeq y$ or $x \succeq z_0 \succ y$. and we see that at least one of the two sums in the display is strictly positive, which yields $U(x) > U(y)$. If $U(x) > U(y)$, then we can use the displayed formula to conclude that $x \succ y$.

Conversely, we assume that a numerical representation is given. We also assume that \mathcal{X} is uncountable, otherwise there is nothing to prove. Let $\mathcal{J} := \{[a, b] : a, b \in \mathbb{Q}, a < b, U^{-1}([a, b]) \neq \emptyset\}$. Then, for every $I \in \mathcal{J}$, there exists $z_I \in \mathcal{X}$ with $U(z_I) \in I$. Put $A := \{z_I : I \in \mathcal{J}\}$ and observe that A is countable. The set A is almost the set \mathcal{Z} we are after. A naive approach could be as follows. Suppose $y \succ x$, then $U(y) > U(x)$ and there are rational a and b such that $U(x) < a < b < U(y)$. The problem arises that it is not

guaranteed that $U^{-1}([a, b])$ is non-void. To remedy this, we consider the set $C := \{(x, y) \in A^c \times A^c : y \succ x \text{ and } \forall z \in A : x \succeq z \text{ or } z \succeq y\}$. Let $(x, y) \in C$, but suppose that there exists $z \in \mathcal{X} \setminus A$ such that $y \succ z \succ x$. Then we can also find rational a and b such that $U(x) < a < U(z) < b < U(y)$ and therefore $I := [a, b] \in \mathcal{J}$. By definition of A , we can then find $z_I \in A$ that then also has the property $U(x) < a \leq U(z_I) \leq b < U(y)$ and hence $y \succ z_I \succ x$. This contradicts $(x, y) \in C$. We conclude that if $(x, y) \in C$, then for all $z \in \mathcal{X}$ it holds that $x \succeq z$ or $z \succeq y$. This implies the following observation. If $(x, y) \in C$ and $(x', y') \in C$, such that $U(x) \neq U(x')$ or $U(y) \neq U(y')$, then $(U(x), U(y)) \cap (U(x'), U(y')) = \emptyset$. We argue as follows. The situation $x \sim x'$ and $y \sim y'$ is ruled out by assumption. Therefore assume w.l.o.g. that $x \approx x'$. Since $(x, y) \in C$, we must have $x \succeq x'$ or $x' \succeq y$, which implies that either $U(x) \geq U(x')$ or $U(x') \geq U(y)$. In the latter case we are done. Let then the first inequality hold. Since also $(x', y') \in C$, we have $x' \succeq x$ or $x \succeq y'$. The first of these possibilities can not happen, since we ruled out $x \sim x'$, and therefore the second one holds, and we obtain $U(x) \geq U(y')$, from which the conclusion follows as well. Knowing that the intervals of the type $(U(x), U(y))$ with $(x, y) \in C$ are disjoint, we conclude that there are only countably many of them and it follows that the collection of these intervals can be written as a collection of intervals $(U(x), U(y))$, where x and y run through a countable subset of \mathcal{X} , B say. We put $\mathcal{Z} = A \cup B$, a countable set as well, and we will see that it is order dense. Take $x, y \in \mathcal{X} \setminus \mathcal{Z}$ with $y \succ x$. If there is $z \in A$ such that $y \succ z \succ x$, we are done. If such a z doesn't exist, then $(x, y) \in C$, in which case we have for instance $U(x) = U(z)$ for some $z \in B$. But then $y \succ z \succeq x$. \square

Not every preference order admits a numerical representation. It turns out that the *lexicographical order* on $[0, 1] \times [0, 1]$ provides us with a counterexample. This order is defined by $x \succ y$ iff $x_1 > y_1$ or $x_1 = y_1$ and $x_2 > y_2$.

Example 2.7 Let $\mathcal{X} = [0, 1] \times [0, 1]$ endowed with the lexicographical order \succ . Suppose that \succ admits a numerical representation U . Since $(\alpha, 1) \succ (\alpha, 0)$, we have $d(\alpha) := U(\alpha, 1) - U(\alpha, 0) > 0$ for all $\alpha \in [0, 1]$, and hence $[0, 1] = \cup_n A_n$, where $A_n = \{\alpha \in [0, 1] : d(\alpha) > \frac{1}{n}\}$. Since $[0, 1]$ is uncountable, there must be a set A_m with infinitely many elements. In this set we can choose $\alpha_0 < \dots < \alpha_N$. Note that $U(\alpha_{i+1}, 0) > U(\alpha_i, 1)$, and so we get $U(\alpha_{i+1}, 0) - U(\alpha_i, 0) > d(\alpha_i) > \frac{1}{m}$. Hence we get

$$\begin{aligned} U(1, 1) - U(0, 0) &= U(1, 1) - U(\alpha_N, 0) \\ &\quad + \sum_{i=0}^{N-1} (U(\alpha_{i+1}, 0) - U(\alpha_i, 0)) \\ &\quad + U(\alpha_0, 0) - U(0, 0) \\ &> \frac{N}{m}. \end{aligned}$$

Letting $N \rightarrow \infty$ yields $U(1, 1) - U(0, 0) = \infty$, which is excluded.

Definition 2.8 Let \mathcal{X} be a topological space. A preference relation \succ is called *continuous* if for every $x \in \mathcal{X}$ the sets $((x, \rightarrow)) := \{y \in \mathcal{X} : y \succ x\}$ and $((\leftarrow, x)) := \{y \in \mathcal{X} : x \succ y\}$ are open.

We will also use the notation $((x, y))$ for $((\leftarrow, y)) \cap ((x, \rightarrow))$. Likewise we define $[[x, \rightarrow)) = \{y \in \mathcal{X} : y \succeq x\}$, $((\leftarrow, x]] = \{y \in \mathcal{X} : x \succeq y\}$ etc. These sets will be called *preference intervals*.

Remark 2.9 Suppose that \succ admits a numerical representation U . Because of the identity $((x, \rightarrow)) = U^{-1}(U(x), \infty)$, it follows that \succ is continuous, if U is a continuous function. But there are also examples of preference orders that are not continuous. Consider again the lexicographical order on $\mathcal{X} = [0, 1] \times [0, 1]$. Then $\{(y_1, y_2) \in \mathcal{X} : (y_1, y_2) \succ (\frac{1}{2}, \frac{1}{2})\}$ is not open in the ordinary topology (draw a picture).

Proposition 2.10 Let \mathcal{X} be a Hausdorff space. On $\mathcal{X} \times \mathcal{X}$ we use the product topology. Then the following are equivalent.

- (i) \succ is continuous.
- (ii) The set $\{(x, y) : y \succ x\}$ is open.
- (iii) The set $\{(x, y) : y \succeq x\}$ is closed.

Proof First we show (i) \Rightarrow (ii): Let $(x_0, y_0) \in M := \{(x, y) : y \succ x\}$. We show that there are open subsets U and V of \mathcal{X} such that $(x_0, y_0) \in U \times V \subset M$. Suppose first that $((x_0, y_0)) \neq \emptyset$. Pick a z from this preference interval, then $y_0 \succ z \succ x_0$. The sets $U := ((\leftarrow, z))$ and $V := ((z, \rightarrow))$ are open and contain x_0 and y_0 respectively. Moreover, one quickly sees that $U \times V \subset M$. If the intersection $((x_0, y_0))$ is empty, we choose $U = ((\leftarrow, y_0))$ and $V = ((x_0, \rightarrow))$. Take $(x, y) \in U \times V$. Then $y_0 \succ x$ and $y \succ x_0$. To show that $y \succ x$, we assume the contrary. By negative transitivity we must have $y_0 \succ y$. But then $y \in ((x_0, y_0))$, which was empty. Contradiction.

(ii) \Rightarrow (iii): It follows from (ii) that also $\{(x, y) : x \succ y\}$ is open. But its complement is just $\{(x, y) : y \succeq x\}$.

(iii) \Rightarrow (i): Since \mathcal{X} is Hausdorff, every singleton $\{x\}$ is closed and so $\{x\} \times \mathcal{X}$ is closed in the product topology. By assumption, then also $\{x\} \times \{y : y \succeq x\} = \{x\} \times \mathcal{X} \cap \{(u, y) : y \succeq u\}$ is closed. But then, the set $\{y : y \succeq x\}$ is closed in \mathcal{X} since a product set is closed iff all factors are closed, and so $\{y : x \succ y\}$ is open. In a similar way one proves that $\{y : y \succ x\}$ is open. \square

Proposition 2.11 Let \mathcal{X} be a connected topological space endowed with a continuous preference order \succ . Then every dense set \mathcal{Z} of \mathcal{X} is also order dense. If \mathcal{X} is separable, then there exists a numerical representation of \succ .

Proof First we rule out the trivial situation in which all elements of \mathcal{X} are indifferent. So, we can take $x, y \in \mathcal{X}$ with $y \succ x$. Observe that $y \in ((x, \rightarrow))$ and $x \in ((\leftarrow, y))$, so both open preference intervals are non-empty. Moreover, their union is \mathcal{X} , because of negative transitivity. Then we must have that

$((x, \rightarrow)) \cap ((\leftarrow, y)) \neq \emptyset$, because \mathcal{X} is connected. The intersection is open as well, so it must contain a z from \mathcal{Z} , since \mathcal{Z} is dense. Then $y \succ z \succ x$, and so \mathcal{Z} is order dense. If \mathcal{X} is separable, there exists a countable dense and thus order dense subset. Apply Theorem 2.6. \square

Connectedness is essential in Proposition 2.11. Here is a counterexample. Let x_0, y_0 be irrational numbers in \mathbb{R} with $x_0 < y_0$. Let $\mathcal{X} = (-\infty, x_0] \cup [y_0, \infty)$ and let \succ be the usual order on \mathbb{R} . Obviously $\mathbb{Q} \cap \mathcal{X}$ is dense in \mathcal{X} , but not order dense, since $[x_0, y_0]$ (by definition a subset of \mathcal{X}) contains no rational numbers.

The assertion of Proposition 2.11, that every continuous preference relation on a connected topological space has a numerical representation, can be sharpened.

Theorem 2.12 *Let \mathcal{X} be a connected and separable topological space, endowed with a continuous preference order. Then this preference order admits a continuous numerical representation.*

Proof We rule out the trivial case that $x \sim y$ for all $x, y \in \mathcal{X}$. Let \mathcal{Z} be a countable dense subset in \mathcal{X} , write $\mathcal{Z} = \{z_1, z_2, \dots\}$. We will first construct a representation U_0 of \succ restricted to \mathcal{Z} , and then give it a continuous extension U on \mathcal{X} . The construction will be recursive by ‘filling the holes’ and bears some similarity with the construction of the Cantor function.

We define $U_0(z_1) := \frac{1}{2}$. Consider z_2 . Three possibilities arise. If $z_2 \sim z_1$, then $U_0(z_2) = U_0(z_1)$. If $z_1 \succ z_2$, then $U_0(z_2) := \frac{1}{4}$ and if $z_2 \succ z_1$ we put $U_0(z_2) = \frac{3}{4}$. For later use we define $V_n = \{k2^{-n} : k = 1, \dots, 2^n - 1\}$. Notice that $U_0(z_1) \in V_1$ and $U_0(z_2) \in V_2$, whatever z_2 , and that $V_n \subset V_{n+1}$ for all n .

We now give the general pattern. Supposing that $U_0(z_1), \dots, U_0(z_n)$ ($n \geq 2$) have been defined and that $U_0(z_k) \in V_k \subset V_n$, for $k \leq n$. We now define $U_0(z_{n+1})$ after realizing that only four different situations can arise. Case 1: z_{n+1} is indifferent to some z_i with $i \leq n$. Then $U_0(z_{n+1}) := U_0(z_i) \in V_n \subset V_{n+1}$. Case 2: $z_i \succ z_{n+1}$ for all $i \leq n$. Then $U_0(z_{n+1}) = \frac{1}{2} \min\{U_0(z_i) : i = 1, \dots, n\} \in V_{n+1}$. Case 3: $z_{n+1} \succ z_i$ for all $i \leq n$. Then $U_0(z_{n+1}) = \frac{1}{2}(\max\{U_0(z_i) : i = 1, \dots, n\} + 1) \in V_{n+1}$. Case 4: There exist $i_n, j_n \in \{1, \dots, n\}$ such that for all other $i \in \{1, \dots, n\}$ it holds that either $z_{i_n} \succeq z_i$ or $z_i \succeq z_{j_n}$ and $z_{j_n} \succ z_{n+1} \succ z_{i_n}$. In this case we put $U_0(z_{n+1}) = \frac{1}{2}(U_0(z_{i_n}) + U_0(z_{j_n})) \in V_{n+1}$.

Let $V = \cup_n V_n$, $u_0 = \inf U_0(\mathcal{Z})$ and $u_1 = \sup U_0(\mathcal{Z})$. Notice that $u_0 < u_1$. Obviously, $U_0(\mathcal{Z}) \subset V$, but we claim that also $V \cap (u_0, u_1) \subset U_0(\mathcal{Z})$. To see that this holds true, we argue as follows.

We may assume without loss of generality that u_0 and u_1 are not attained. If this were not true, we replace below \mathcal{Z} with the non-empty set $\mathcal{Z} \setminus U_0^{-1}(\{u_0, u_1\})$. If both u_0, u_1 are attained, $u_0 = U_0(z^0)$ and $u_1 = U_0(z^1)$ say, this set is dense in $((z^0, z^1))$ (Exercise 2.4). If only one of the two is attained, a similar argument applies. We continue under the assumption made, u_0 and u_1 are not attained.

We have $U_0(z_1) = \frac{1}{2}$. The sets $((\leftarrow, z_1))$ and $((z_1, \rightarrow))$ are both open and non-empty, otherwise we would have $u_0 = U_0(z_1)$ or $u_1 = U_0(z_1)$, contradicting our assumption that u_0 and u_1 are not attained. Hence there must be z and z'

in \mathcal{Z} such that $U_0(z) = \frac{1}{4}$ and $U_0(z') = \frac{3}{4}$. Hence all values in $V_2 \cap (u_0, u_1)$ are attained. We continue by induction. Suppose that all values in $V_n \cap (u_0, u_1)$ are attained. Then for all $1 \leq k < 2^n - 1$, there are $z, z' \in \mathcal{Z}$ such that $U_0(z) = k2^{-n}$ and $U_0(z') = (k+1)2^{-n}$. The set $((z, z'))$ is open and, by the fact that \mathcal{X} is connected, not empty. Since \mathcal{Z} is dense, there is $z'' \in \mathcal{Z} \cap ((z, z'))$. We can even choose z'' such that $U_0(z'') = (2k+1)2^{-n-1} \in V_{n+1} \setminus V_n$, by the construction of U_0 . If $2^{-n} \in V_n \cap (u_0, u_1)$ and z is such that $U_0(z) = 2^{-n}$, then $2^{-n} > u_0$ (the infimal value) and there must be a z' such that $U_0(z') < 2^{-n}$. By the construction of U_0 , we can choose z' even such that $U_0(z') = 2^{-n-1}$. For z such that $U_0(z) = 1 - 2^{-n}$ there is a similar reasoning. This shows that all values in V_{n+1} are attained as well.

Having thus proved the claim, we take closures and it follows that $[u_0, u_1] = \text{Cl}U_0(\mathcal{Z})$, so $U_0(\mathcal{Z})$ is dense in $[u_0, u_1]$. It is obvious that the equivalence (2.1) holds for all $x \in \mathcal{Z}$, by construction of U_0 on \mathcal{Z} . We extend U_0 to have domain \mathcal{X} by setting

$$U(x) = \sup\{U_0(z) : z \in \mathcal{Z}, z \preceq x\}.$$

First we check that $U(z) = U_0(z)$, for $z \in \mathcal{Z}$. It is obvious that $U(z) \geq U_0(z)$. Suppose that $U(z) > U_0(z)$. Then there would be $z' \prec z$ such that $U_0(z') > U_0(z)$ and then also $z' \succ z$, a contradiction. Now we show that U is a numerical representation of \succ . Suppose that $x \succeq y$. Then $\{U_0(z) : z \in \mathcal{Z}, z \preceq y\} \subset \{U_0(z) : z \in \mathcal{Z}, z \preceq x\}$ and $U(x) \geq U(y)$ obviously holds. Let now $x \succ y$. Then, by \mathcal{Z} dense and \mathcal{X} connected, there are $z, z' \in \mathcal{Z}$ such that $x \succeq z \succ z' \succeq y$ (Exercise 2.3), and hence $U(x) \geq U(z) > U(z') \geq U(y)$.

We finally show that U is continuous, for which it is sufficient to show that the sets of the form $U^{-1}((-\infty, u))$ and $U^{-1}((u, \infty))$ are open for $u \in [u_0, u_1]$. We focus on $U^{-1}((-\infty, u))$, and we only have to consider $u > u_0$. Take x from this set. By V dense, there is $v \in V$ such that $U(x) < v$ and $v < u$. But $v = U(z)$ for some $z \in \mathcal{Z}$. Hence, we have $U^{-1}((u, \infty))$ is a union of sets $((\leftarrow, z))$, which are open by assumption. \square

2.3 Exercises

2.1 Prove Proposition 2.3.

2.2 Assume that \succ is a continuous preference relation on a connected set \mathcal{X} , which is endowed with a topology that is first-countable (this allows you to work below with sequences). Let \mathcal{Z} be a dense subset of \mathcal{X} . If $U : \mathcal{X} \rightarrow \mathbb{R}$ is continuous and its restriction to \mathcal{Z} is a numerical representation of \succ , then U is also a numerical representation of \succ on all of \mathcal{X} . To show this, you verify the following implications.

- (a) $x \succ y \Rightarrow U(x) > U(y)$
- (b) $U(x) > U(y) \Rightarrow x \succ y$.

Hints: To show (a) you complete the following steps. Show that there are $z, w \in \mathcal{Z}$ such that $x \succ z \succ w \succ y$. Choose then $z_n, w_n \in \mathcal{Z}$ such that $z_n \rightarrow x$ and $w_n \rightarrow y$ and finish the proof.

For (b) you show first that $U^{-1}(U(y), \infty) \cap U^{-1}(-\infty, U(x))$ is non-void and select $z, w \in \mathcal{Z}$ such that $U(x) > U(z) > U(w) > U(y)$. Use again convergent sequences.

2.3 Let \mathcal{X} be a connected topological space and \succ a continuous strict preference relation on it.

- (a) Let $x \succ y$. Show that $\mathcal{X} = ((y, \rightarrow))^c \cup ((y, x)) \cup ((\leftarrow, x))^c$.
- (b) Show that $((y, x))$ is not empty.
- (c) Let \mathcal{Z} be dense in \mathcal{X} . Show that there are $z, z' \in \mathcal{Z}$ such that $x \succ z \succ z' \succ y$.

2.4 Show that the set $\{z \in \mathcal{Z} : U_0(z) \in (u_0, u_1)\}$ in the proof of Theorem 2.12 is dense in $((z^0, z^1))$.

2.5 Let \mathcal{X} be a topological space with a continuous preference order \succ and topology \mathcal{T} . Let \mathcal{S} be the set of all preference intervals $((\leftarrow, x))$ or $((x, \rightarrow))$. Let \mathcal{T}_0 be the smallest topology that contains \mathcal{S} . Show that \mathcal{T}_0 is the coarsest topology for which Proposition 2.11 and Theorem 2.12 are valid.

2.6 Let \succ be a strict preference order on a set \mathcal{X} and \succeq its associated weak preference order. Prove, using the definitions of these preference orders, the following statements.

- (a) \sim is an equivalence relation on \mathcal{X} .
- (b) $x \succ y$ iff $x \succeq y$ and $x \not\sim y$.
- (c) $x \succeq y$ iff $x \succ y$ or $x \sim y$.

2.7 A subset K of \mathbb{R} is compact in the ordinary topology iff it is closed and bounded, in which case there is a maximal number in K . Here is the counterpart for preference relations. Let \succeq be a continuous weak preference relation on a set \mathcal{X} , which is endowed with a topology under which it becomes a Hausdorff space (a compact subset of \mathcal{X} is then closed). Let K be a non-empty compact subset of \mathcal{X} . Show that there is a most preferred element \bar{x} in K (i.e. $\bar{x} \succeq x$ for all $x \in K$) and that the set of all such \bar{x} is compact. *Hint:* Consider for each $x \in K$ the preference intervals in K , $[[x, \rightarrow)) \cap K$, and show that any finite intersection of them is non-empty.

2.8 Let \succeq be a continuous weak preference relation on $\mathcal{X} = (0, \infty)^m$ that has the additional property that it is monotonic: if $x = (x_1, \dots, x_m)$ and $y = (y_1, \dots, y_m)$ are such that $x_i \geq y_i$ for all i and $x_i \neq y_i$ for some i , then x is strictly preferred over y , $x \succ y$. Let $\mathbf{1}$ be the vector $(1, \dots, 1)$. Define $U(x) = \sup\{\alpha > 0 : x \succeq \alpha \mathbf{1}\}$, $x \in \mathcal{X}$.

- (a) Show that U is a numerical representation of \succeq .
- (b) Consider for given $x \in \mathcal{X}$ the sets $U = \{\alpha \geq 0 : \alpha \mathbf{1} \preceq x\}$ and $L = \{\alpha \geq 0 : \alpha \mathbf{1} \succeq x\}$. Show that (i) $L \cap U \neq \emptyset$ and that (ii) actually this intersection consists of one point only.
- (c) Define $U'(x) = \inf\{\alpha > 0 : x \preceq \alpha \mathbf{1}\}$, $x \in \mathcal{X}$. Show that $U = U'$.

2.9 Let $\mathcal{X} = [0, 1] \times [0, 1]$ endowed with the lexicographical order \succ . Show by a direct argument, not referring to Theorem 2.6, that \mathcal{X} has no countable order dense subset.

2.10 Consider the function U as in the proof of Theorem 2.12, whose assumptions are in force. Define $V(x) = \inf\{U_0(z) : z \in \mathcal{Z}, z \succeq x\}$. Show that $U(x) \leq V(x)$, for all $x \in \mathcal{X}$. Is $U = V$?

3 Lotteries

In this section we take the situation of the previous section as a starting point. We assume that the set \mathcal{X} is a convex subset of the set of all probability measures on some measurable space (S, \mathcal{S}) . We write \mathcal{M} instead of \mathcal{X} . Every probability measure can be considered as a *lottery*, in the ‘argot du métier’. Our aim is to consider preference relations on the space of lotteries that admit a numerical representation of a special kind.

3.1 Von Neumann-Morgenstern representations

Definition 3.1 Let \succ be a preference relation on \mathcal{M} . A numerical representation U is called a *Von Neumann-Morgenstern* representation if there is a measurable function $u : S \rightarrow \mathbb{R}$ such that

$$(3.1) \quad U(\mu) = \int u \, d\mu, \forall \mu \in \mathcal{M}.$$

It is easy to check that a Von Neumann-Morgenstern representation U is an affine function. But, if a numerical representation U of \succ is affine, then it implies two additional properties of \succ (see Proposition 3.3), that we define now.

Definition 3.2 Let \succ be a preference relation on \mathcal{M} . It satisfies the *independence axiom* if for all $\mu \succ \nu$ it holds that

$$(3.2) \quad t\mu + (1-t)\lambda \succ t\nu + (1-t)\lambda,$$

for all $\lambda \in \mathcal{M}$ and $t \in (0, 1]$. The preference relation satisfies the *Archimedean axiom*, if for all $\mu \succ \lambda \succ \nu$, there are $t, s \in (0, 1)$ such that

$$(3.3) \quad t\mu + (1-t)\nu \succ \lambda \succ s\mu + (1-s)\nu.$$

Proposition 3.3 Assume that \succ admits an affine numerical representation. Then \succ satisfies the axioms of Definition 3.2.

Proof Exercise 3.1. □

The nice thing is that Proposition 3.3 has a converse as well. This is the content of the next theorem.

Theorem 3.4 Suppose that a preference relation \succ on \mathcal{M} satisfies both the independence and Archimedean axioms. Then it has an affine numerical representation, U say. Moreover, for any other affine numerical representation \tilde{U} , there exist $a > 0$ and $b \in \mathbb{R}$ such that $\tilde{U} = aU + b$.

To prepare for the proof of this theorem, we present a lemma. In the proof we use a couple of times a consequence of the independence axiom. If $\mu \succ \nu$, then $\mu \succ t\mu + (1-t)\nu$ for all $t \in [0, 1)$ and $t\mu + (1-t)\nu \succ \nu$ for all $t \in (0, 1]$.

Lemma 3.5 Suppose that a preference relation \succ on \mathcal{M} satisfies both the independence and Archimedean axioms. Then the following hold true.

- (i) If $\mu \succ \nu$, then for all $0 \leq t < s \leq 1$ it holds that $s\mu + (1-s)\nu \succ t\mu + (1-t)\nu$.
- (ii) If $\mu \succ \nu$ and $\lambda \in [[\nu, \mu]]$, then there exists a unique $t \in [0, 1]$ such that $\lambda \sim t\mu + (1-t)\nu$.
- (iii) If $\mu \sim \nu$, then $t\mu + (1-t)\lambda \sim t\nu + (1-t)\lambda$, for all $\lambda \in \mathcal{M}$ and $t \in [0, 1]$.
- (iv) For any $\rho \succeq \lambda$, the preference interval $[[\lambda, \rho]]$ is convex.

Proof (i) We use (3.2), with s instead of t and $\lambda = \nu$, to get $\rho := s\mu + (1-s)\nu \succ \nu$. Using (3.2) again, with $\rho = \mu = \lambda$ and $1-u$ instead of t , we get $(1-u)\rho + u\rho \succ (1-u)\nu + u\rho$. Here the left hand side is ρ , whereas the right hand side equals $su\mu + (1-su)\nu$. Take $u = t/s \in (0, 1)$.

(ii) We only have to show existence, uniqueness follows from (i). Existence for the cases $\lambda \sim \mu$ and $\lambda \sim \nu$ is trivial, so we assume $\mu \succ \lambda \succ \nu$. In view of (i), it should hold that $t = \sup A := \sup\{u : \lambda \succ u\mu + (1-u)\nu\}$. Suppose that this t is not the right one, then either $\lambda \succ t\mu + (1-t)\nu$ or $\lambda \prec t\mu + (1-t)\nu$. Consider the first case, which rules out $t = 1$, so $t < 1$. We use the right hand side of (3.3) applied to the triple $\mu \succ \lambda \succ t\mu + (1-t)\nu$ to get existence of $s \in (0, 1)$ such that $\lambda \succ s(t\mu + (1-t)\nu) + (1-s)\mu = (1-s+ts)\mu + (1-t)s\nu$. The definition of t implies $t \geq 1-s+ts$, which is, since $s < 1$, equivalent to $t \geq 1$, a contradiction. In the second of the two above cases, we apply (3.3) to the triple $t\mu + (1-t)\nu \succ \lambda \succ \nu$, to get $s \in (0, 1)$ such that $st\mu + (1-st)\nu = s(t\mu + (1-t)\nu) + (1-s)\nu \succ \lambda$. This means that $st \notin A$, and hence $st \geq \sup A = t$, so $s \geq 1$, another contradiction.

(iii) We rule out the case that all $\rho \in \mathcal{M}$ are equivalent to μ , because then we immediately have $t\mu + (1-t)\lambda \sim \mu$, $t\nu + (1-t)\lambda \sim \mu$ and we are done. So, take $\rho \approx \mu$ and suppose that $\rho \succ \mu$ (the other case can be treated similarly). Then also $\rho \succ \nu$ and we apply (3.2) to obtain $s\rho + (1-s)\nu \succ s\nu + (1-s)\nu = \nu$, for all $s \in (0, 1)$. Then we apply it again to get

$$(3.4) \quad t(s\rho + (1-s)\nu) + (1-t)\lambda \succ t\mu + (1-t)\lambda.$$

If the assertion were not true, then we have for instance $t\mu + (1-t)\lambda \succ t\nu + (1-t)\lambda$ (the other possibility can be treated similarly). Assume that $t\mu + (1-t)\lambda \succ t\nu + (1-t)\lambda$, contrary to what we have to show. Now we can apply (ii), which yields a unique $u \in (0, 1)$ such that $t\mu + (1-t)\lambda \sim u(t(s\rho + (1-s)\nu) + (1-t)\lambda) + (1-u)(t\nu + (1-t)\lambda) = tsu\rho + t(1-su)\nu + (1-t)\lambda$. Equation (3.4) is true for all s , and so we can replace it with su . It now follows that $t\mu + (1-t)\lambda \succ t\mu + (1-t)\lambda$, which is a contradiction, caused by the assumption $t\mu + (1-t)\lambda \succ t\nu + (1-t)\lambda$. Likewise, one can eliminate $t\mu + (1-t)\lambda \prec t\nu + (1-t)\lambda$ to complete the proof.

(iv) Next we show that $[[\lambda, \rho]]$ is convex. We assume $\rho \succ \lambda$, since the case $\rho \sim \lambda$ immediately follows from part (ii). Consider first the interior case, we take $\mu, \nu \in ((\lambda, \rho))$ and $t \in (0, 1)$. Since $\rho \succ \nu$ we use the independence axiom to get $\rho = t\rho + (1-t)\rho \succ t\rho + (1-t)\nu$. And since $\rho \succ \mu$, we use the same axiom to get $t\rho + (1-t)\nu \succ t\mu + (1-t)\nu$. Combining these relations, we obtain $\rho \succ t\mu + (1-t)\nu$. One similarly proves $t\mu + (1-t)\nu \succ \lambda$.

Next we consider a boundary case $\mu \sim \rho$, $\nu \in ((\lambda, \rho))$. Inspection of the proof of the previous case shows that this case is partly handled, and one can use part (iii) of this lemma to complete the proof (you check!). Finally we have the extreme case $\nu \sim \lambda$ and $\mu \sim \rho$. Here one can use part (iii) again (Exercise 3.6). We conclude that $[[\lambda, \rho]]$ is convex. \square

Proof of Theorem 3.4 We exclude the trivial case in which all elements of \mathcal{M} are indifferent. Choose $\rho, \lambda \in \mathcal{M}$ with $\rho \succ \lambda$. Let $\mu \in [[\lambda, \rho]]$. Lemma 3.5(ii) yields a unique $t = t(\mu)$ such that $\mu \sim t\rho + (1-t)\lambda$. We use this to define U on $[[\lambda, \rho]]$ by $U(\mu) := t$. Notice that $U(\lambda) = 0$ and $U(\rho) = 1$. The first thing to show is that we have defined a numerical representation of \succ . Let $U(\mu) > U(\nu)$. In view of Lemma 3.5(i) we have $U(\mu)\rho + (1-U(\mu))\lambda \succ U(\nu)\rho + (1-U(\nu))\lambda$. But the probability measures on both sides are indifferent to μ and ν respectively. Hence $\mu \succ \nu$. To prove the converse implication it is sufficient to show that $U(\mu) = U(\nu)$ implies $\mu \sim \nu$. But this is obvious from the definition of U .

We now show that U is affine. Since $[[\lambda, \rho]]$ is convex (Lemma 3.5(iv)), $U(t\mu + (1-t)\nu)$ is well defined for $\mu, \nu \in [[\lambda, \rho]]$. Recall that $U(\mu)\rho + (1-U(\mu))\lambda \sim \mu$ and $U(\nu)\rho + (1-U(\nu))\lambda \sim \nu$. A double application of Lemma 3.5(iii) gives

$$t\mu + (1-t)\nu \sim t(U(\mu)\rho + (1-U(\mu))\lambda) + (1-t)(U(\nu)\rho + (1-U(\nu))\lambda).$$

Rearranging terms on the right hand side gives $(tU(\mu) + (1-t)U(\nu))\rho + (1-tU(\mu) - (1-t)U(\nu))\lambda$, a convex combination of ρ and λ . But then, by definition of U , we have $U(t\mu + (1-t)\nu) = tU(\mu) + (1-t)U(\nu)$, as desired.

The next step is to show that U is unique up to an affine transformation. Let \tilde{U} be another affine representation of \succ . Let $\mu \in [[\lambda, \rho]]$ and define

$$\hat{U}(\mu) = \frac{\tilde{U}(\mu) - \tilde{U}(\lambda)}{\tilde{U}(\rho) - \tilde{U}(\lambda)},$$

an affine transformation of \tilde{U} , having, like U , the properties $\hat{U}(\lambda) = 0$ and $\hat{U}(\rho) = 1$. Combine this with affinity of \tilde{U} to obtain $\hat{U}(\mu) = \hat{U}(U(\mu)\rho + (1-U(\mu))\lambda) = U(\mu)\hat{U}(\rho) + (1-U(\mu))\hat{U}(\lambda) = U(\mu)$. Therefore U is an affine transformation of \tilde{U} .

The last step is to show that U can be extended to all of \mathcal{M} . Consider a preference interval $[[\lambda_1, \rho_1]] \supset [[\lambda, \rho]]$. We know that \succ has an affine representation U_1 on $[[\lambda_1, \rho_1]]$, which can be taken such that $U_1(\rho) = 1$, $U_1(\lambda) = 0$ (apply an affine transformation to accomplish this). So U_1 must coincide with U on $[[\lambda, \rho]]$. Hence U can be extended to all of \mathcal{M} , since every element of it belongs to some preference interval by transitivity of \succeq . Indeed, if $\mu \notin [[\lambda, \rho]]$, then e.g. $\mu \prec \lambda$ and we can take $[[\lambda_1, \rho_1]] = [[\mu, \rho]] \supset [[\lambda, \rho]]$. \square

Remark 3.6 We note that up to here, we didn't use that \mathcal{M} is a set of probability measure. The above results are valid, under the stated assumptions, for any convex set \mathcal{M} . On the other hand, for \mathcal{M} a set of probability measures, convex combinations of the type $t\mu + (1-t)\nu$ have a nice interpretation as a compound lottery. The results that follow use essential properties of probability measures.

We return to the Von Neumann-Morgenstern representation of a preference order \succ on \mathcal{M} . We first treat a simple case.

Example 3.7 Suppose that \mathcal{M} is the set of all finite mixtures of Dirac measures δ_x , and that an affine representation exists. Define $u(x) = U(\delta_x)$. Let $\mu = \sum t_i \delta_{x_i}$, where the $t_i \geq 0$ and $\sum t_i = 1$. Affinity of U yields $U(\mu) = \sum t_i u(x_i) = \int u d\mu$, which is the desired representation. So, in this case, if there exists an affine representation, it is automatically of Von Neumann-Morgenstern type.

In the remainder of this section we assume that the set S is a separable metric space and that \mathcal{S} is its Borel σ -algebra. As a preparation for the final theorem, we have the following lemma.

Lemma 3.8 Consider the space \mathcal{M} of all probability measures on (S, \mathcal{S}) endowed with the weak topology. Fix $\mu, \nu \in \mathcal{M}$ and consider $A : t \rightarrow t\mu + (1-t)\nu$. Then $A : [0, 1] \rightarrow \mathcal{M}$ is continuous. If \succ is a continuous preference ordering on \mathcal{M} , then it satisfies the Archimedean axiom.

Proof The first assertion follows from the evident identity $\int f d(t\mu + (1-t)\nu) = t \int f d\mu + (1-t) \int f d\nu$, valid for any bounded and continuous function f on S . To prove the second assertion, let $\mu \succ \nu$ and choose $\lambda \in ((\nu, \mu))$. Observe that $t = 1$ is an element of $A^{-1}((\lambda, \rightarrow))$ and that this set is open in $[0, 1]$ by the just shown continuity of A . Hence there is also some $t \in (0, 1)$ belonging to it, and for this t one has $A(t) = t\mu + (1-t)\nu \succ \lambda$, as required in Definition 3.2. The existence of s in that definition follows similarly. \square

Theorem 3.9 Consider the space \mathcal{M} of all probability measures on (S, \mathcal{S}) endowed with the weak topology, where S is assumed to be separable. Let \succ be a continuous preference ordering on \mathcal{M} , satisfying the independence axiom. Then \succ admits a Von Neumann-Morgenstern representation

$$(3.5) \quad U(\mu) = \int u d\mu,$$

where the function $u : S \rightarrow \mathbb{R}$ is bounded, continuous and unique up to affine transformations.

Proof Consider first the subspace \mathcal{M}_S of simple distributions on S , these are the distributions as in Example 3.7. We conclude from Lemma 3.8 and Theorem 3.4 that \succ restricted to \mathcal{M}_S admits an affine representation, which is, by Example 3.7, automatically of Von Neumann-Morgenstern type.

The function u involved will turn out to be bounded. Suppose that this is not the case, then there is a sequence $(x_n) \subset S$ such that $(u(x_n))$ is increasing and $u(x_n) > n$ (the other possibility $u(x_n) < -n$ can be treated similarly). Put $\mu_n = (1 - \frac{1}{\sqrt{n}})\delta_{x_1} + \frac{1}{\sqrt{n}}\delta_{x_n}$. Since $u(x_2) > u(x_1)$, we have $\delta_{x_2} \succ \delta_{x_1}$, so $\delta_{x_1} \in ((\leftarrow, \delta_{x_2}))$. One easily checks that $\mu_n \rightarrow \delta_{x_1}$ weakly. Hence, for n big enough, μ_n belongs to any (nonempty) open neighborhood of δ_{x_1} , so eventually we have $\mu_n \in ((\leftarrow, \delta_{x_2}))$. But then $U(\mu_n) \leq u(x_2)$. However, by

direct computation, we have $U(\mu_n) > (1 - \frac{1}{\sqrt{n}})u(x_1) + \sqrt{n}$, which yields a contradiction.

We now show that u is continuous. Suppose the contrary, then there is a sequence (x_n) converging to some $x \in S$, whereas $u(x_n)$ doesn't converge to $u(x)$. Assume e.g. that one has $\limsup u(x_n) < u(x)$. Then along a subsequence, again denoted by (x_n) , one has $\lim u(x_n) =: a < u(x)$. In particular, there is $m \in \mathbb{N}$ such that $|u(x_n) - a| < \frac{1}{3}(u(x) - a)$, for $n \geq m$; equivalently $\frac{4}{3}a - \frac{1}{3}u(x) < u(x_n) < \frac{2}{3}a + \frac{1}{3}u(x)$, for $n \geq m$. Put $\mu = \frac{1}{2}(\delta_x + \delta_{x_m})$. Then also $U(\delta_x) = u(x) > \frac{2}{3}u(x) + \frac{1}{3}a > \frac{1}{2}(u(x) + u(x_m)) = U(\mu) > \frac{1}{3}u(x) + \frac{2}{3}a > U(\delta_{x_n})$, for $n \geq m$. So, $\delta_x \succ \mu \succ \delta_{x_n}$. This means that δ_{x_n} doesn't belong to the open neighborhood $((\mu, \rightarrow))$ of δ_x , contradicting the fact that $\delta_{x_n} \rightarrow \delta_x$ weakly.

We now show that, knowing the function u , Equation (3.5) defines a numerical representation U of \succ . Since u is bounded and continuous, U , as defined in (3.5), is continuous w.r.t. the weak topology. It is a fact that the set of simple distributions is weak-dense in the set of all probability measures on (S, \mathcal{S}) , see Proposition A.12. Since we know that U is a numerical representation of \succ on the set of simple distributions, we can argue as in the proof of Theorem 2.12, that U is also a numerical representation on the collection of all probability measures on (S, \mathcal{S}) (see also Exercise 2.2).

Finally, u is unique up to affine transformations. This follows from the fact that affine numerical representations are unique up to affine transformations (Theorem 3.4). \square

Later on we need representations of preference orders, where u is unbounded. This cannot happen under the conditions of Theorem 3.9. A way out is obtained, by replacing the weak topology with a stronger one. Let ψ be a continuous function on S with $\psi \geq 1$. Let \mathcal{M}^ψ be the set of probability measures μ such that $\int \psi d\mu < \infty$ and let C^ψ be the space of continuous functions f such that $|f|/\psi$ is bounded. If μ is a probability measure in \mathcal{M}^ψ , then

$$\mu^\psi(B) := \frac{\int \mathbf{1}_B \psi d\mu}{\int \psi d\mu}$$

defines another probability measure on (S, \mathcal{S}) . We say that a sequence $(\mu_n) \subset \mathcal{M}^\psi$ converges in the ψ -weak topology if the corresponding sequence (μ_n^ψ) converges in the weak topology. If the ψ -weak limit is μ , then this means nothing else than $\int f d\mu_n \rightarrow \int f d\mu$, for all $f \in C^\psi$. This immediately yields the following corollary to Theorem 3.9.

Corollary 3.10 *Let \succ be a preference order on \mathcal{M}^ψ that is continuous w.r.t. the ψ -weak topology and that satisfies the independence axiom. Then there exists a Von Neumann-Morgenstern representation of the form (3.5), with $u \in C^\psi$. Also in this case, U and u are unique up to affine transformations.*

Proof The proof is based on the fact that the transformation $\mu \rightarrow \mu^\psi$, can be used to apply the results of Theorem 3.9. Details (Exercise 3.2) are left to the reader. \square

3.2 Exercises

3.1 Prove Proposition 3.3.

3.2 Prove Corollary 3.10.

3.3 Let S be the set of positive integers, $S = \mathbb{N}$. Let \mathcal{M} be the collection of probability measures μ on \mathbb{N} with the property that $U(\mu) := \lim n^2 \mu(n) < \infty$. Then U is affine and induces a preference order \succ on \mathcal{M} .

- (a) Show that \succ satisfies both the independence and Archimedean axioms.
- (b) Show that \succ does not admit a Von-Neumann-Morgenstern representation.
- (c) Why can't we apply Theorem 3.9?

3.4 Finish the proof of Theorem 3.9. [If you want, you may use Exercise 2.2, and that \mathcal{M} is metrizable. See also Corollary A.13.]

3.5 Suppose you are a plumber and you have a client that wants to pay you 1000 euro for installing a drainage system. If you do nothing and stay home, you don't get paid. Let μ be the sure 'lottery' that pays out 1000 euro with certainty, λ the sure 'lottery' that pays zero. Then for you $\mu \succ \lambda$. Let ν be the 'lottery' in which you will be shot. Then, most likely, $\mu \succ \lambda \succ \nu$. Is there for you a $t \in (0, 1)$ such that $t\mu + (1 - t)\nu \succ \lambda$? Same question if ν is the sure 'lottery' to get killed in a car crash on your way to the client.

3.6 Prove convexity for the remaining cases in the proof of Lemma 3.5(iv).

4 Utility and expected utility

In this section we consider a set \mathcal{M} of probability measures on an interval S of \mathbb{R} , and \mathcal{S} will be the Borel σ -algebra on S . We assume that \mathcal{M} is convex and contains all Dirac measures on points in S , and consequently all simple measures.

4.1 Risk aversion

We depart from the common assumption that the *fair price* of a lottery $\mu \in \mathcal{M}$ equals its *expectation* $m(\mu)$. We assume, unless the contrary is explicitly stated, that these expectations exist for all $\mu \in \mathcal{M}$. It frequently happens (but it depends on the circumstances) that somebody who has the choice between a lottery with an average pay-off of let's say 1000 euro and getting this amount of money straight away, prefers the latter option. He then exhibits *risk averse* behavior, as a result of his personal preferences. This notion will be made precise in the next definition.

Definition 4.1 A preference order \succ on \mathcal{M} is called *monotone* if the implication $x > y \Rightarrow \delta_x \succ \delta_y$ holds ($x, y \in S$). It is called *risk averse* if $\delta_{m(\mu)} \succ \mu$, unless μ is degenerate, $\mu = \delta_{m(\mu)}$.

Proposition 4.2 Suppose that a preference order \succ on \mathcal{M} has Von Neumann-Morgenstern representation

$$U(\mu) = \int u \, d\mu,$$

for some Borel measurable function u (the integrals are assumed to be well defined for all $\mu \in \mathcal{M}$). Then

- (i) the preference order is monotone iff u is strictly increasing and
- (ii) the preference order is risk averse iff u is strictly concave.

Proof (i) Notice that $U(\delta_x) = u(x)$. Then $u(x) > u(y)$ iff $U(\delta_x) > U(\delta_y)$ iff $\delta_x \succ \delta_y$.

(ii) Suppose that \succ is risk averse. Take $x, y \in S$ and consider $\mu = t\delta_x + (1-t)\delta_y$ for $t \in (0, 1)$. Then $m(\mu) = tx + (1-t)y$. Then the risk averse \succ yields $U(\delta_{m(\mu)}) > U(\mu)$, or $u(tx + (1-t)y) > tu(x) + (1-t)u(y)$. Hence u is strictly concave. Conversely, for strict concave u Jensen's inequality gives for any nondegenerate $\mu \in \mathcal{M}$ that $U(\delta_{m(\mu)}) = u(m(\mu)) > \int u \, d\mu = U(\mu)$. \square

The function u in the Von Neumann-Morgenstern representation of a monotone risk averse preference relation deserves a name of its own.

Definition 4.3 A function $u : S \rightarrow \mathbb{R}$ is called a *utility function* if u is strictly increasing, strictly concave and continuous on S .

Since any concave function is continuous on the interior of the set on which it is defined, Exercise 4.9, the continuity requirement above only concerns boundary points of S . And since u is increasing, in fact it is only a condition of continuity of u in $\inf S$, if this is an element of S .

Definition 4.4 A preference order \succ on \mathcal{M} admits an *expected utility representation* U if there exists a utility function u such that $U(\mu) = \int u d\mu$, for all $\mu \in \mathcal{M}$.

In the remainder of this section, we assume that preference orders admit expected utility representations.

Continuity of u implies that $u(S)$ is connected and hence for all $\mu \in \mathcal{M}$ there exists a number $c(\mu) \in S$ such that $u(c(\mu)) = U(\mu) \in u(S)$. Moreover, $c(\mu)$ is unique, because u is strictly increasing. Whence the indifferent relation $\delta_{c(\mu)} \sim \mu$. In words, playing a lottery μ is indifferent to obtaining the sure amount $c(\mu)$ under a given preference ordering.

Definition 4.5 The number $c(\mu)$ is called the *certainty equivalent* of the lottery μ and the difference $\rho(\mu) := m(\mu) - c(\mu)$ is called the *risk premium* of μ .

Notice that always $c(\mu) \leq m(\mu)$ for risk averse \succ and that strict inequality holds for nondegenerate μ . Hence, a risk averse person with utility function u will not pay more than $c(\mu)$ to play a lottery μ . Conversely, the risk premium is the amount of money a seller of the lottery μ has to pay to a risk averse agent to convince him to exchange the sure amount $m(\mu)$ for the random pay-off of the lottery μ .

In the present context, we consider the following optimization problem. Find, if it exists, a lottery μ^* that is most preferred among all lotteries in a subset of \mathcal{M} , equivalently, the one with the highest value of U , where U is of expected utility type.

We specialize to a specific case. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be given and a random variable X defined on it, with values in S , that has a nondegenerate distribution μ . Let $c \in \mathbb{R}$ and consider the convex combination $X_\lambda = \lambda c + (1 - \lambda)X$. Note that the distribution function of X_λ is obtained by a location-scale transformation of that of X . Write μ_λ for the distribution of X_λ ($\mu_0 = \mu$). Put

$$f(\lambda) = U(\mu_\lambda) = \int u d\mu_\lambda.$$

Proposition 4.6 *Assume that S is an interval, $X \geq a$ for some $a \in \text{Int}S$, $\mathbb{E}X < \infty$ and $c \in \text{Int}S$.*

- (i) *The function $f : [0, 1] \rightarrow \mathbb{R}$ is strictly concave and hence its maximal value is assumed for some unique $\lambda^* \in [0, 1]$.*
- (ii) *We have $\lambda^* = 1$ if $m(\mu) = \mathbb{E}X \leq c$ and $\lambda^* > 0$, if $c > c(\mu)$.*

(iii) If moreover u is differentiable, then we even have $\lambda^* = 1 \Leftrightarrow \mathbb{E} X \leq c$ and $\lambda^* = 0 \Leftrightarrow c \leq \frac{\mathbb{E} X u'(X)}{\mathbb{E} u'(X)}$.

Proof (i) Since $f(\lambda) = \mathbb{E} u(X_\lambda)$, strict concavity of f follows from strict concavity of u .

(ii) Jensen's inequality yields

$$f(\lambda) \leq u(\mathbb{E} X_\lambda) = u(\mathbb{E} X + \lambda(c - \mathbb{E} X)),$$

with equality iff $\lambda = 1$. Since u is increasing, the right hand side is non-decreasing in λ if $c \geq \mathbb{E} X$. Under this condition, $\lambda^* = 1$.

Concavity of u yields $u(X_\lambda) \geq (1 - \lambda)u(X) + \lambda u(c)$, hence

$$f(\lambda) \geq (1 - \lambda)u(c) + \lambda u(X),$$

with equality iff $\lambda = 0, 1$. The right hand side is non-decreasing under the condition $c \geq c(\mu)$, in which case $\lambda^* > 0$.

(iii) Assume that u is differentiable. Because f is concave, $\lambda^* = 0$ can only happen if f is decreasing in a neighborhood of zero, so when the right derivative $f'_+(0) \leq 0$. Let us compute this derivative. We have

$$(4.1) \quad \frac{u(X_\lambda) - u(X)}{\lambda} = \frac{u(X_\lambda) - u(X)}{X_\lambda - X}(c - X).$$

The nonnegative difference quotient in (4.1) is bounded by the derivative of u in the left endpoint of the involved interval, which is either $X_0 = X > a$ or $X_1 = c$. Hence the absolute value of (4.1) is bounded by $\min\{u'(X_0), u'(X_1)\}|c - X| \leq u'_+(c \wedge a)|c - X|$, which has finite expectation, since $u'_+(c \wedge a) < \infty$ because $c, a \in \text{Int}S$ and $\mathbb{E}|X| < \infty$. Taking expectations in (4.1) and letting $\lambda \downarrow 0$, we get by the Dominated Convergence Theorem that the limit is $f'_+(0) = \mathbb{E} u'(X)(c - X)$. Hence $f'_+(0) \leq 0$ iff $c \leq \frac{\mathbb{E} X u'(X)}{\mathbb{E} u'(X)}$.

In much the same way, $\lambda^* = 1$ iff f is non-decreasing in a neighborhood of $\lambda = 1$, $f'_-(1) \geq 0$, Exercise 4.7. Working with a difference quotient like (4.1) for $\lambda \uparrow 1$ and using that $X_1 = c$, we get $f'_-(1) = u'(c)(c - \mathbb{E} X)$. The last assertion now also follows. \square

Example 4.7 Consider a risky asset S_1 with price π_1 , and a riskless asset with interest rate r ($S_0 = 1 + r$). Suppose that an agent has a C^1 utility function u and a capital (initial wealth) w . Suppose that he builds a portfolio by investing a fraction λ of his capital in the riskless asset and the rest in the risky asset. The value of the portfolio ("at time $t = 1$ ") is then $\lambda w(1 + r) + (1 - \lambda)wS_1/\pi_1$, and the discounted net gain is

$$(1 - \lambda) \frac{w}{\pi_1} \left(\frac{S_1}{1 + r} - \pi_1 \right).$$

The previous proposition shows that $\lambda^* = 1$ (all capital invested in the riskless asset) iff $\frac{\mathbb{E} S_1}{1 + r} \leq \pi_1$. Hence such an agent is only willing to invest in the risky

asset, when the price is below the expected discounted value. Note that this holds for any risk averse investor, regardless the special form of the utility function u . Compare this with what happens under the risk-neutral measure of Section 1.1.

4.2 Arrow-Pratt coefficient

Suppose that one considers a probability measure μ that has finite variance and that is concentrated on a small interval around its mean m . Let u be a C^2 utility function on a neighborhood of this interval and let U be the associated expected utility representation. Look at the following heuristic. A Taylor expansion of u around m gives

$$u(x) \approx u(m) + (x - m)u'(m) + \frac{1}{2}(x - m)^2u''(m).$$

Hence $u(c(\mu)) = U(\mu) = \int u d\mu \approx u(m) + \frac{1}{2}\text{Var}(\mu)u''(m)$. On the other hand, the same Taylor expansion gives $u(c(\mu)) \approx u(m) + (c(\mu) - m)u'(m) + \frac{1}{2}(c(\mu) - m)^2u''(m)$. Since $c(\mu)$ will be close to m , we have $u(c(\mu)) \approx u(m) + (c(\mu) - m)u'(m)$. Hence, for the risk premium $\rho(\mu) = m - c(\mu)$ we have the approximation

$$(4.2) \quad \rho(\mu) \approx -\frac{1}{2} \frac{u''(m)}{u'(m)} \text{Var}(\mu).$$

Definition 4.8 Let u be a twice differentiable utility function on some (open) interval S . Then the quantity

$$\alpha(x) := -\frac{u''(x)}{u'(x)}$$

is called the *Arrow-Pratt coefficient of absolute risk aversion of u at the level x* .

We conclude from Equation (4.2), that for probability measures μ that are concentrated around the mean m , the risk premium $\rho(\mu)$ approximately factors as a product of the Arrow-Pratt coefficient at the level m (solely determined by the utility function u) and half the variance (which is an intrinsic quantity of μ only). Notice that by concavity, $\alpha(x) \geq 0$ for every utility function u .

Arrow-Pratt coefficients have the attractive feature that they are invariant under affine transformations. Since in Von Neumann-Morgenstern representations of preference orders, the function u is unique up to affine transformations, this means that the Arrow-Pratt coefficient in such a situation is an intrinsic feature of the preference order, not of its numerical representation (of course modulo the fact that we have to assume that u is C^2 , and that u is not constant, which would lead anyway to an uninteresting preference order).

We now give some examples of widely used utility functions.

Example 4.9 Let u be such that the Arrow-Pratt function $\alpha(\cdot)$ is a (positive) constant, also denoted by α . Then, by solving a second order linear differential equation, one finds, for some constants $a \in \mathbb{R}$ and $b > 0$,

$$u_{a,b}(x) = a - be^{-\alpha x},$$

which is an affine transformation of $u(x) = 1 - \exp(-\alpha x)$. Note that u is defined on all of \mathbb{R} . The functions $u_{a,b}$ are called CARA functions (from Constant Absolute Risk Aversion).

Example 4.10 Here we introduce the HARA (from Hyperbolic Absolute Risk Aversion) utility functions. For these functions we have that $\alpha(x) = \frac{c}{x}$, for $c, x > 0$. For convenience we write $c = 1 - \gamma$, and hence $\gamma < 1$. Solving the corresponding differential equation for u yields

$$u_{a,b}(x) = \frac{a}{\gamma}x^\gamma + b,$$

for $\gamma \neq 0$ and $u_{a,b}(x) = a \log x + b$ for $\gamma = 0$. Note that $\gamma \geq 1$ is excluded by requiring that u is strictly concave. The above functions $u_{a,b}$ are affine transformations of $u_{1,0}$.

Remark 4.11 HARA utility functions with $\gamma > 0$ are examples of utility functions $u : [0, \infty) \rightarrow \mathbb{R}$ satisfying the *Inada conditions*, i.e. $u \in C^1(0, \infty)$, with $\lim_{x \rightarrow 0} u'(x) = \infty$ and $\lim_{x \rightarrow \infty} u'(x) = 0$.

There are close connections between utility functions, risk premia and Arrow-Pratt coefficients for different preference orders.

Proposition 4.12 Suppose $u_1, u_2 : S \rightarrow \mathbb{R}$ are two C^2 utility functions, with corresponding risk premia $\rho_1(\cdot), \rho_2(\cdot)$ and Arrow-Pratt coefficients $\alpha_1(\cdot)$ and $\alpha_2(\cdot)$. The following are equivalent.

- (i) $\alpha_1(x) \geq \alpha_2(x), \forall x \in S$.
- (ii) There exist a strictly increasing concave function F , defined on the range of u_2 , such that $u_1 = F \circ u_2$.
- (iii) $\rho_1(\mu) \geq \rho_2(\mu), \forall \mu \in \mathcal{M}$.

Proof (i) \Rightarrow (ii): The obvious choice of F is $F(x) = u_1(u_2^{-1}(x))$. Clearly, F is well defined, since u_2 is strictly increasing, and since u_2^{-1} and u_1 are strictly increasing, so is F . To show that F is concave, we compute its second derivative and use that (i) is assumed. Notice that it sufficient to show that $F''(u_2(x)) \leq 0$, for all $x \in S$. We start with $u_1(x) = F(u_2(x))$ and get

$$\begin{aligned} u_1'(x) &= F'(u_2(x))u_2'(x) \\ u_1''(x) &= F''(u_2(x))u_2'(x)^2 + F'(u_2(x))u_2''(x). \end{aligned}$$

Solving the second of these two equations for $F''(u_2(x))$ and using the first one yields

$$\begin{aligned}
 (4.3) \quad F''(u_2(x)) &= \frac{u_1''(x) - \frac{u_1'(x)}{u_2'(x)}u_2''(x)}{u_2'(x)^2} \\
 &= \frac{u_1'(x)}{u_2'(x)^2} \left(\frac{u_1''(x)}{u_1'(x)} - \frac{u_2''(x)}{u_2'(x)} \right) \\
 &= \frac{u_1'(x)}{u_2'(x)^2} (\alpha_2(x) - \alpha_1(x)),
 \end{aligned}$$

by definition of the Arrow-Pratt coefficients. By assumption (i) and the fact that u_1 is increasing, we have $F''(u_2(x)) \leq 0$.

(ii) \Rightarrow (iii): By Jensen's inequality, applied to the concave function F , it holds that

$$\begin{aligned}
 (4.4) \quad u_1(c_1(\mu)) &= \int u_1 \, d\mu = \int F \circ u_2 \, d\mu \\
 &\leq F\left(\int u_2 \, d\mu\right) = F(u_2(c_2(\mu))) = u_1(c_2(\mu)).
 \end{aligned}$$

Since u_1 is increasing, we must have $c_1(\mu) \leq c_2(\mu)$, from which the result follows.

(iii) \Rightarrow (i): Suppose that (i) doesn't hold. Then for some x one has $\alpha_1(x) < \alpha_2(x)$, and by continuity of α_1 and α_2 , this equality extends to an open neighborhood O of x . By (4.3), which is also valid without assumptions (i) or (ii), we then have $F''(u_2(x)) > 0$ on O . Take now a nondegenerate probability measure μ such that $\mu(O) = 1$. Then strict convexity of $F \circ u_2$ leads to a strict equality in the opposite direction as compared to (4.4), $u_1(c_1(\mu)) > u_1(c_2(\mu))$, from which it follows that $c_1(\mu) > c_2(\mu)$, contradicting assumption (iii). \square

4.3 Exercises

4.1 Show that for a utility function $u \in C^1(\mathbb{R})$ it holds that $m(\mu) > c(\mu) > \frac{\mathbb{E} X u'(X)}{\mathbb{E} u'(X)}$, where X has nondegenerate distribution μ .

4.2 Let $u(x) = 1 - \exp(-x)$, a CARA function. Consider an investor with utility function u who wants to invest an initial capital. There is one riskless asset, having value 1 and interest rate $r = 0$, and one risky assets with random pay-off S_1 having a normal $N(m, \sigma^2)$ distribution with $\sigma^2 > 0$. Suppose he invests a fraction λ in the riskless asset and the remainder in the risky asset. The pay-off of this portfolio is thus $\lambda + (1 - \lambda)S_1$. The aim is to maximize his expected utility.

- Show that $\mathbb{E} \exp(uS_1) = \exp(um + \frac{1}{2}u^2\sigma^2)$ ($u \in \mathbb{R}$).
- Compute for each λ the certainty equivalent of the portfolio.
- Let λ^* be the optimal value of λ . Give, by direct computations, conditions on the parameters such that each of the cases $\lambda^* = 0$, $\lambda^* = 1$ or $\lambda^* \in (0, 1)$ occurs.

(d) Compare the results of (c) with the assertions of Proposition 4.6.

4.3 In Exercise 4.2 the optimization problem turns out to be of the form: maximize $\mathbb{E} Z - c \text{Var} Z$. This seems reasonable, if one thinks of Z as a random revenue. One wants to maximize the expected revenue and to keep the ‘risk’ in terms of variance low. In general such a maximization problem leads to odd results. Consider the following example. In two lotteries the random pay-off Z satisfies $\mathbb{P}(Z = h) = p_i$ and $\mathbb{P}(Z = \ell) = 1 - p_i$, $i = 1, 2$ and $h > \ell$. Find an example of values of $p_1 > p_2$ and $c > 0$ such that the second lottery is preferred to the first one.

4.4 Consider a twice differentiable utility function $u : S \rightarrow \mathbb{R}$. Define for fixed x such that $tx \in S$ the function $t \mapsto v_x(t) = u(tx)$. A way to establish the *relative risk* around x can be obtained by inspection of $v_x(t)$ in a neighborhood of $t = 1$. A measure of relative risk at x is defined by $r(x) = -v_x''(1)/v_x'(1)$. Show that $r(x) = x\alpha(x)$ ($\alpha(x)$ the Arrow-Pratt risk measure).

4.5 A utility function u is said to exhibit decreasing risk aversion if the function $x \mapsto \alpha(x)$ is decreasing. Show that this property is equivalent to saying that for every $x_1 < x_2$ there exists a concave function g such that $u(x_2 + z) = g(u(x_1 + z))$ for all z (for which the given expressions make sense).

4.6 Let u be a (continuous) strictly increasing function, $u : S \rightarrow \mathbb{R}$. Consider the fair game represented by a random variable X with values $x \pm \varepsilon$ ($\varepsilon > 0$) in S that are attained with equal probabilities $\frac{1}{2}$. Given x, ε , the *probability premium* $\pi = \pi(x, \varepsilon)$ is by definition such that the lottery with the same outcomes but with probability $\mathbb{P}(\xi = x - \varepsilon) = \frac{1}{2} - \pi$ has expected utility $u(x)$. Show that an individual who uses u for a Von-Neumann-Morgenstern representation is risk averse (in which case u is a utility function) iff $\pi(x, \varepsilon) > 0$ for all x, ε . Sketch the graph of u and construct $\pi(x, \varepsilon)$.

4.7 Show the statement concerning $\lambda^* = 1$ in Proposition 4.6(iii).

4.8 The optimal value λ^* in Proposition 4.6 should be nondecreasing and continuous as a function of c , and strictly increasing on the interval $(\frac{\mathbb{E} X u'(X)}{\mathbb{E} u'(X)}, m(\mu))$. Verify whether this is true.

4.9 Let $f : S \rightarrow \mathbb{R}$ be concave, where S is an interval. Show that f is continuous on the interior of S and give an example where f is not continuous in a boundary point of S (which is assumed to belong to S).

4.10 Let u be an exponential utility function, $u(x) = -\exp(-\alpha x)$, $x \in \mathbb{R}$, $\alpha > 0$. Find the maximizing λ for the problem in Proposition 4.6 in each of the cases (a) X assumes two values only, (b) X has an exponential distribution, (c) X has a log-normal distribution. [I have not checked whether explicit solutions exist.]

5 Stochastic dominance

Results in the previous sections were depending on the preference orders, or the utility functions, at hand. In the present section, we will look at preferences that are independent of a particular choice of a utility function belonging to a certain class. The *standing assumptions* are that we deal with the set \mathcal{M} of all probability measures on $(\mathbb{R}, \mathcal{B})$ that admit a *finite expectation*. As a consequence, for any utility function u , the integrals $\int u d\mu$ are well defined, but may take on the value $-\infty$. This holds, since every concave function has an affine function as a majorant. Indeed, since for some $a, b > 0$, one has $u(x) \leq ax + b$ for all x , it holds that $u(x)^+ \leq a|x| + b$ and hence $\int u^+ d\mu < \infty$.

5.1 Uniform order

Definition 5.1 Let $\mu, \nu \in \mathcal{M}$. One says that μ is *uniformly preferred* over ν , denoted by $\mu \succeq_{\text{uni}} \nu$, if

$$\int u d\mu \geq \int u d\nu, \text{ for all utility functions } u.$$

Remark 5.2 The uniform preference of the above definition is also called *second order stochastic dominance*. Notice that it is not a weak preference order (see Definition 2.2), since it is not complete. In Section 5.2 we will discuss first order stochastic dominance.

The next theorem gives a number of characterizations of uniform preference, there are many more.

Theorem 5.3 *There is equivalence between the following statements.*

- (i) $\mu \succeq_{\text{uni}} \nu$.
- (ii) For all increasing concave functions f , one has $\int f d\mu \geq \int f d\nu$.
- (iii) For all $c \in \mathbb{R}$, it holds that $\int (c - x)^+ \mu(dx) \leq \int (c - x)^+ \nu(dx)$.
- (iv) If F_μ and F_ν are the distribution functions of μ and ν , then $\int_{-\infty}^c F_\mu(x) dx \leq \int_{-\infty}^c F_\nu(x) dx$, for all $c \in \mathbb{R}$.

Proof (i) \Leftrightarrow (ii): Obviously (ii) \Rightarrow (i). For the converse implication we need a utility function that has finite integral under μ and ν . This can be accomplished as follows (see Figure 1 for an illustration). Take a given utility function u and an arbitrary $x_0 \in \mathbb{R}$. Modify u on $(-\infty, x_0]$ by replacing u with $x \mapsto u'_+(x_0)(2(x - x_0) - \exp(x - x_0) + 1) + u(x_0)$. Check that the modified function is still a utility function! Moreover, the modified utility function (denoted u again) has finite integral w.r.t. any probability measure with finite expectation. If f is increasing and concave, then $u_\alpha(x) := \alpha f(x) + (1 - \alpha)u(x)$ defines a strictly increasing, strictly concave continuous function, so a utility function, for every $\alpha \in [0, 1)$. The assertion follows from

$$\int f d\mu = \lim_{\alpha \uparrow 1} \int u_\alpha d\mu \geq \lim_{\alpha \uparrow 1} \int u_\alpha d\nu = \int f d\nu.$$




Figure 1: $u(x) = 1 - e^{-x}$ and $x_0 = 1$

(ii) \Leftrightarrow (iii): Clearly (ii) \Rightarrow (iii). The converse implication basically follows from the fact that every nonnegative convex decreasing function, with limit zero at infinity, is a pointwise limit of positive linear combinations of functions $x \mapsto (c - x)^+$ and that $-f$ is decreasing and convex. More formally, we have that $h = -f$ admits right derivatives $h'_+(x)$ at every point x . The function h'_+ is increasing, right continuous and on any interval $(a, b]$, *up to scaling*, it is a distribution function of a probability measure. Stated otherwise, there is a measure γ on $(\mathbb{R}, \mathcal{B})$ such that $\gamma(a, b] = h'_+(b) - h'_+(a)$, for all $a < b$. Since there exists only countably many discontinuity points of h'_+ , we have for $x < b$

$$\begin{aligned}
 h(x) &= h(b) - \int_{(x,b]} h'_+(y) \, dy \\
 (5.1) \quad &= h(b) - \int_{(x,b]} (h'_+(y) - h'_+(b)) \, dy - h'_+(b)(b - x).
 \end{aligned}$$

We first rewrite the integral in (5.1). Let $B = \{(u, y) : x < y < u \leq b\}$. We have

$$\begin{aligned}
 \int_{(x,b]} (h'_+(y) - h'_+(b)) \, dy &= - \int_{(x,b]} \gamma(y, b] \, dy \\
 &= - \int_{(x,b]} \int \mathbf{1}_{(y,b]} \, d\gamma \, dy \\
 &= - \int \int \mathbf{1}_B(u, y) \gamma(du) \, dy \\
 &= - \int \int \mathbf{1}_B(u, y) \, dy \, \gamma(du) \quad (\text{by Fubini}) \\
 &= - \int \mathbf{1}_{(x,b]}(u)(u - x) \gamma(du) \\
 &= - \int \mathbf{1}_{(-\infty, b]}(u - x)^+ \gamma(du).
 \end{aligned}$$

Hence, we can rewrite $h(x)$ as

$$h(x) = h(b) - h'_+(b)(b-x) + \int \mathbf{1}_{(-\infty, b]}(u-x)^+ \gamma(du).$$

Let μ be a probability measure on $(\mathbb{R}, \mathcal{B})$. Integration of the last expression w.r.t. μ and using Fubini's theorem again, yields

$$\begin{aligned} \int_{(-\infty, b]} h d\mu &= h(b)\mu(-\infty, b] - h'_+(b) \int (b-x)^+ \mu(dx) \\ &\quad + \int_{(-\infty, b]} \int_{(-\infty, b]} (u-x)^+ \mu(dx) \gamma(du) \\ &= h(b)\mu(-\infty, b] - h'_+(b) \int (b-x)^+ \mu(dx) \\ &\quad + \int_{(-\infty, b]} \int (u-x)^+ \mu(dx) \gamma(du). \end{aligned}$$

Using condition (iii) and the fact that $h'_+ \leq 0$, we have an upper bound for the last displayed expression by replacing μ with ν . It follows that

$$\int_{(-\infty, b]} h d\mu \leq \int_{(-\infty, b]} h d\nu + h(b)(\mu(-\infty, b] - \nu(-\infty, b]).$$

Since h is lower bounded by an affine function, we have that $\int_{(b, \infty)} h d\mu$ and $\int_{(b, \infty)} h d\nu$ are both finite. Hence we obtain

$$\begin{aligned} \int h d\mu &\leq \int h d\nu + \int_{(b, \infty)} h d\mu - \int_{(b, \infty)} h d\nu + h(b)(\mu(-\infty, b] - \nu(-\infty, b]) \\ &= \int h d\nu - \int_{(b, \infty)} (h(b) - h(x))\mu(dx) + \int_{(b, \infty)} (h(b) - h(x))\nu(dx). \end{aligned}$$

We finally show that the last two integrals vanish for $b \rightarrow \infty$. Since they are similar, we treat only the first of the two. Fix b_0 and let $b > b_0$. It holds that $0 \leq h(b) - h(x) \leq -h'_+(b_0)(x - b_0)$ for $x > b$. Hence

$$\int_{(b, \infty)} (h(b) - h(x))\mu(dx) \leq -h'_+(b_0) \int (x - b_0) \mathbf{1}_{(b, \infty)}(x) d\mu,$$

which tends to zero by the Dominated convergence theorem, since $\int |x| \mu(dx)$ is finite. Hence we obtain $\int h d\mu \leq \int h d\nu$, which is equivalent to (ii).

(iii) \Leftrightarrow (iv): This is just a matter of rewriting, using Fubini's theorem. One has

$$\int_{-\infty}^c F_\mu(y) dy = \int_{-\infty}^c \int \mathbf{1}_{(-\infty, y]}(x) \mu(dx) dy$$

$$\begin{aligned}
&= \int_{(-\infty, c]} \int_x^c dy \mu(dx) \\
&= \int_{(-\infty, c]} (c-x) \mu(dx) \\
(5.2) \quad &= \int (c-x)^+ \mu(dx).
\end{aligned}$$

□

Remark 5.4 It follows from Theorem 5.3(ii), that $\mu \succeq_{\text{uni}} \nu$ implies $m(\mu) \geq m(\nu)$. The integrals w.r.t. the measure μ in assertion (iii) of the same theorem in fact determine μ . Indeed, by the computations leading to (5.2), we see that knowing integrals of $(c-x)^+$ for all c is equivalent to knowing the integrals of F_μ up to c . Taking right derivatives w.r.t. c gives $F'_\mu(c)$ and knowing this for all c determines μ . This fact can be used to show that \succeq_{uni} defines a partial order, Exercise 5.1.

If μ is the distribution of a random variable $X \in L^1(\Omega, \mathcal{F}, \mathbb{P})$ and ν that of $Y \in L^1(\Omega, \mathcal{F}, \mathbb{P})$, such that $\mu \succeq_{\text{uni}} \nu \succeq_{\text{uni}} \mu$, then $\mu = \nu$, so X and Y have the same *distribution* (under \mathbb{P}). Yet, X and Y are in general very different as random variables. It may happen that $\mathbb{P}(X = Y) = 0$.

When two lotteries with the same mean are compared, we can develop the assertions of Theorem 5.3 a little further.

Proposition 5.5 For all probability measures $\mu, \nu \in \mathcal{M}$ the following are equivalent.

- (i) $\mu \succeq_{\text{uni}} \nu$ and $m(\mu) = m(\nu)$.
- (ii) $\int f d\mu \geq \int f d\nu$, for all concave functions f .
- (iii) $m(\mu) \geq m(\nu)$ and $\int (x-c)^+ \mu(dx) \leq \int (x-c)^+ \nu(dx)$, for all $c \in \mathbb{R}$.

Proof (i) \Rightarrow (ii): First we show that the assertion holds true for decreasing concave functions. Such a function is $x \mapsto -(c-x)^-$, for arbitrary $c \in \mathbb{R}$. Since $-(c-x)^- = c-x - (c-x)^+$, the assertion for such a function follows from Theorem 5.3 and the assumptions that $m(\mu) = m(\nu)$ and $\mu \succeq_{\text{uni}} \nu$, because $x \mapsto -(c-x)^+$ is concave and increasing. The proof for arbitrary decreasing functions is then similar to the proof of (iii) \Rightarrow (ii) of Theorem 5.3.

If f is concave, but not monotone, then there exists a $x_0 \in \mathbb{R}$, such that $f(x) \leq f(x_0)$, for all $x \in \mathbb{R}$. Let

$$f_1(x) = \begin{cases} f(x) & \text{if } x \leq x_0 \\ f(x_0) & \text{if } x > x_0 \end{cases}$$

and

$$f_2(x) = \begin{cases} f(x_0) & \text{if } x \leq x_0 \\ f(x) & \text{if } x > x_0. \end{cases}$$

Then f_1 is concave and increasing and f_2 is concave and decreasing. Knowing that the assertions hold true for f_1 and f_2 , we obtain the same result for f , because $f(x) = f_1(x) + f_2(x) - f(x_0)$.

(ii) \Rightarrow (iii): Take first $f(x) \equiv x$ to get the first assertion, and then $f(x) \equiv -(x - c)^+$, which is concave, to get the second one from (ii).

(iii) \Rightarrow (i): Rewrite the inequality between the integrals in (iii) as

$$\int_{(c, \infty)} x \mu(dx) - c + c \mu(-\infty, c] \leq \int_{(c, \infty)} x \nu(dx) - c + c \nu(-\infty, c].$$

Let $c \rightarrow -\infty$ and use that both measures have a finite first moment to arrive at $\int x \mu(dx) \leq \int x \nu(dx)$, or $m(\mu) \leq m(\nu)$. Together with the assumption, this gives $m(\mu) = m(\nu)$. Use the identity $y^+ = y + (-y)^+$ ($y \in \mathbb{R}$) to get

$$\int (c - x)^+ \mu(dx) = c - m(\mu) + \int (x - c)^+ \mu(dx).$$

A similar equality holds for ν . Using the assumption and $m(\mu) = m(\nu)$, we arrive at $\int (c - x)^+ \mu(dx) \leq \int (c - x)^+ \nu(dx)$, condition (iii) in Theorem 5.3 to get $\mu \succeq_{\text{uni}} \nu$. \square

Remark 5.6 Assume that $\mu_1 \succeq_{\text{uni}} \mu_2$ and $m(\mu_1) = m(\mu_2)$. Then it follows from Proposition 5.5 that $\text{Var} \mu_1 \leq \text{Var} \mu_2$. For normal distributions there is a converse relationship, see Exercise 5.2.

5.2 Monotone order

We turn to another concept of stochastic dominance, also called *first order stochastic dominance*. There are more of these concepts conceivable.

Definition 5.7 Let μ, ν be two probability measures on $(\mathbb{R}, \mathcal{B})$. One says that μ *stochastically dominates* ν , if for all bounded increasing continuous functions $f : \mathbb{R} \rightarrow \mathbb{R}$ it holds that

$$(5.3) \quad \int f d\mu \geq \int f d\nu.$$

In this case one writes $\mu \succeq_{\text{mon}} \nu$.

It is almost trivial to check that \succeq_{mon} defines a partial order on the space of probability distributions on $(\mathbb{R}, \mathcal{B})$. Below we give an easy characterization of $\mu \succeq_{\text{mon}} \nu$.

Proposition 5.8 Let μ, ν be two probability measures on $(\mathbb{R}, \mathcal{B})$ and let F_μ and F_ν be their distribution functions. The following are equivalent.

- (i) It holds that $\mu \succeq_{\text{mon}} \nu$.
- (ii) For all $x \in \mathbb{R}$ one has $F_\mu(x) \leq F_\nu(x)$.

Proof (i) \Rightarrow (ii): We'd like to apply the definition of stochastic dominance to the function $u \mapsto \mathbf{1}_{(x, \infty)}(u)$, which is bounded and increasing. The result would then follow. However this function is not continuous. Therefore one first uses the functions $u \mapsto (\min\{n(u-x), 1\})^+$ and let $n \rightarrow \infty$.

(ii) \Rightarrow (i): Let f be continuous, bounded and increasing. We can obtain f (which is measurable) as the pointwise limit of an increasing sequence of simple functions f_n , that are increasing themselves. To see this, we assume for simplicity that $0 \leq f \leq 1$ and we follow the usual approximation scheme, known from measure theory.

Let $n \in \mathbb{N}$ and define $E_{ni} = \{(i-1)2^{-n} < f \leq i2^{-n}\}$ for $i = 1, \dots, 2^n$ and $E_{n0} = \{f = 0\}$. Put

$$f_n = 2^{-n} \sum_{i=1}^{2^n} (i-1) \mathbf{1}_{E_{ni}} = 2^{-n} \sum_{i=2}^{2^n} (i-1) \mathbf{1}_{E_{ni}}.$$

Then we know that $f_n \uparrow f$. Using that the E_{ni} are disjoint for each n , $\bigcup_{i \geq j+1} E_{ni} = \{f > j2^{-n}\}$ and $\{f > 1\} = \emptyset$, we rewrite

$$f_n = 2^{-n} \sum_{i=2}^{2^n} \left(\sum_{j=1}^{i-1} 1 \right) \mathbf{1}_{E_{ni}} = 2^{-n} \sum_{j=1}^{2^n-1} \sum_{i=j+1}^{2^n} \mathbf{1}_{E_{ni}} = 2^{-n} \sum_{j=1}^{2^n} \mathbf{1}_{\{f > j2^{-n}\}}.$$

Note that the f_n are also increasing functions. Since f is continuous, the sets $\{f > j2^{-n}\}$ are open and since f is increasing, there are real numbers a_{nj} such that $\{f > j2^{-n}\} = (a_{nj}, \infty)$. Hence,

$$\int f_n d\mu = 2^{-n} \sum_{j=1}^{2^n} \mu((a_{nj}, \infty)) = 2^{-n} \sum_{j=1}^{2^n} (1 - F_\mu(a_{nj})).$$

It follows from the assumption that $\int f_n d\mu \geq \int f_n d\nu$. The assertion follows by application of the Monotone Convergence Theorem. \square

Remark 5.9 It follows from Theorem 5.3 and Proposition 5.8 that $\mu \succeq_{\text{mon}} \nu$ implies $\mu \succeq_{\text{uni}} \nu$.

5.3 Exercises

5.1 Show that \succeq_{uni} defines a partial order on the set of probability measures with finite mean (see also Remark 5.4).

5.2 Consider two normal distributions, $\mu_1 = N(m_1, \sigma_1^2)$ and $\mu_2 = N(m_2, \sigma_2^2)$.

- Compute $\int_{\mathbb{R}} \exp(-ax) \mu_i(dx)$ and show that $\mu_1 \succeq_{\text{uni}} \mu_2$ implies both $m_1 \geq m_2$ and $\sigma_1^2 \leq \sigma_2^2$.
- Assume that $m_1 = m_2$. Show (use Theorem 5.3(iv)) that $\sigma_1^2 \leq \sigma_2^2$ implies $\mu_1 \succeq_{\text{uni}} \mu_2$.

- (c) Let u be a utility function and assume $m_1 \geq m_2$. Put $\tilde{u}(x) = u(x + m_2)$. Verify that $\mathbb{E} u(N(m_1, \sigma_1^2)) \geq \mathbb{E} \tilde{u}(N(0, \sigma_1^2))$ (the notation should be obvious).
- (d) Let $m_1 \geq m_2$ and $\sigma_1^2 \leq \sigma_2^2$. Show that $\mu_1 \succeq_{\text{uni}} \mu_2$.

5.3 Let $\mu, \nu \in \mathcal{M}$ and f an increasing function such that $\int |f| d\mu$ and $\int |f| d\nu$ are both finite. Show that $\mu \succeq_{\text{mon}} \nu$ implies $\int f d\mu \geq \int f d\nu$ and thus $\mu \succeq_{\text{uni}} \nu$.

5.4 Let $\mu \succeq_{\text{mon}} \nu$ and $m(\mu) = m(\nu)$. Show that $\mu = \nu$. *Hint:* compute $0 \leq \int_a^b (F_\nu(x) - F_\mu(x)) dx$ for any $a < b$. Use integration by parts and let $a \rightarrow -\infty, b \rightarrow \infty$.

5.5 A random variable X has a log-normal distribution with parameters α and σ , if $X = \exp(\alpha + \sigma Z)$, where $\sigma \geq 0$ and Z has a standard normal distribution.

- (a) Compute $\mathbb{E} X^p$ for $p > 0$. In particular, one has $\mathbb{E} X = \exp(\alpha + \frac{1}{2}\sigma^2)$.
- (b) Let μ_i be log-normal distributions ($i = 1, 2$) with parameters α_i, σ_i . Show that $\mu_1 \succeq_{\text{uni}} \mu_2$ implies $m(\mu_1) \geq m(\mu_2)$ and $\sigma_1 \leq \sigma_2$.
- (c) Conversely, if $m(\mu_1) \geq m(\mu_2)$ and $\sigma_1 \leq \sigma_2$, then $\mu_1 \succeq_{\text{uni}} \mu_2$. To prove this, proceed as follows. Let $X_1 = \exp(\alpha_1 + \sigma Z_1)$ and $X_2 = \exp(\alpha_2 + \sigma Z_2)$ (in obvious notation). Let further $X_3 = \exp(\alpha_2 - \alpha_1 + \sqrt{\sigma_2^2 - \sigma_1^2} Z_3)$, where Z_3 is standard normal, independent of Z_1 . Verify that $X_1 X_3$ has the same distribution as X_2 and that $\mathbb{E} X_3 = \frac{m(\mu_2)}{m(\mu_1)}$. Use then Jensen's inequality for conditional expectations to show that $\mathbb{E} u(X_2) \leq \mathbb{E} u(X_1)$.

6 Portfolio optimization

In this section we return to the setting of Section 1 and combine it with the expected utility setting of Section 4. We consider an investor, whose preferences are determined by a utility function \tilde{u} , and who wants to invest a capital w (w from wealth). On the market there are d risky assets having a price (at $t = 0$) given by the vector π and whose future, at time $t = 1$, random pay-off is described by the random vector S , defined on a underlying probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Both vectors are assumed to have strictly positive entries. Next to the risky assets, there is a riskless asset, with price $\pi_0 = 1$ and future value $S_0 = 1 + r > 0$. Let $\bar{S} = (S_0, S)$. A portfolio is given by $\bar{\xi} \in \mathbb{R}^{d+1}$ and we also write $\bar{\xi} = (\xi_0, \xi) \in \mathbb{R} \times \mathbb{R}^d$. The future value of the portfolio is then $\bar{\xi} \cdot \bar{S}$, and it has expected utility

$$(6.1) \quad \mathbb{E} \tilde{u}(\bar{\xi} \cdot \bar{S}).$$

In order for the investor to purchase the portfolio $\bar{\xi}$, the price of it should at most be equal to the initial capital. Thus we have the *budget constraint*

$$(6.2) \quad \bar{\xi} \cdot \bar{\pi} \leq w.$$

We will study the problem of maximizing (6.1) over portfolios $\bar{\xi}$, satisfying the constraint (6.2).

6.1 Optimization and absence of arbitrage

We start this section by casting the above problem in a different, but equivalent form. Our first observation is that it can never be optimal to use only a fraction of the initial capital w . Indeed, suppose one has a portfolio $\bar{\xi}$ with $\bar{\xi} \cdot \bar{\pi} < w$. Change the investment ξ_0 into $\xi'_0 = \xi_0 + w - \bar{\xi} \cdot \bar{\pi}$. Then we have $(\xi'_0, \xi) \cdot \bar{S} = \bar{\xi} \cdot \bar{S} + (w - \bar{\xi} \cdot \bar{\pi})(1 + r) > \bar{\xi} \cdot \bar{S}$. But then, since \tilde{u} is strictly increasing, also $\mathbb{E} \tilde{u}((\xi'_0, \xi) \cdot \bar{S}) > \mathbb{E} \tilde{u}(\bar{\xi} \cdot \bar{S})$. Therefore, we assume from now on that equality holds in (6.2), and so we work with

$$(6.3) \quad \bar{\xi} \cdot \bar{\pi} = w.$$

Recall that we denoted by Y the d -dimensional random vector of discounted net gains,

$$Y = \frac{S}{1+r} - \pi.$$

Hence we have, assuming (6.3), $\bar{\xi} \cdot \bar{S} = (1+r)(\xi \cdot Y + w)$. Define a new utility function u by $u(x) = \tilde{u}((1+r)(w+x))$. Then we have $\tilde{u}(\bar{\xi} \cdot \bar{S}) = u(\xi \cdot Y)$. Of course, this expression only makes sense if $\xi \cdot Y \in D$, where D is the domain of u . Given a risky portfolio ξ , by (6.3) one can always choose a non-risky investment ξ_0 such that the total portfolio has initial price w . This makes the constraint (6.3) redundant, if one considers only ξ as the free variable. All these arguments motivate to study the following *unconstrained* optimization problem, equivalent to the original one.

Problem 6.1 Let $u : D \rightarrow \mathbb{R}$ be a utility function. Maximize

$$\mathbb{E} u(\xi \cdot Y)$$

over all risky portfolios ξ that satisfy $\xi \cdot Y \in D$.

We will study this problem under each of the two cases in the assumption below.

Assumption 6.2 Let $u : D \rightarrow \mathbb{R}$ be a utility function and Y the vector of discounted net gains. Assume either of the following.

- (i) $D = \mathbb{R}$ and u is bounded from above, or
- (ii) $D = [a, \infty)$ for some $a < 0$, and we optimize over the set of ξ such that $\xi \cdot Y \geq a$ a.s. In this case, we also assume that for those ξ the expected utility $\mathbb{E} u(\xi \cdot Y)$ is finite.

For both of these case we write $\Xi = \{\xi \in \mathbb{R}^d : \xi \cdot Y \in D \text{ a.s.}\}$.

Theorem 6.5 below shows that the maximization problem 6.1 only makes sense in an arbitrage-free market, just as pricing of portfolios and derivatives. In the proof of it we use the following two lemmas. Recall the definition of a upper semicontinuous function h (often abbreviated as u.s.c. function), it is such that $\limsup h(x_n) \leq h(x)$, whenever $x_n \rightarrow x$.

Lemma 6.3 Let $h : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{-\infty\}$ be a concave and upper semicontinuous function with $h(0) > -\infty$. Then h attains its supremum, if for all $\xi \neq 0$

$$(6.4) \quad \lim_{\alpha \uparrow \infty} h(\alpha \xi) = -\infty.$$

Proof Let $c < \sup h$. We will see that the non-empty (!) level set $\{h \geq c\}$ is compact. By the fact that h is u.s.c., this set is closed (See Exercise 6.1). So, by the Heine-Borel theorem we only have to show that it is bounded. Suppose that it is unbounded, then there exists a sequence (x_n) such that $|x_n| \rightarrow \infty$ and $h(x_n) \geq c$, for all n . We may assume that the normalized vectors $x_n/|x_n|$ converge to some limit ξ . Let $\alpha > 0$ and consider $h(\alpha \xi)$. One has for all n large enough $\alpha/|x_n| \in (0, 1)$, which will be used below, when concavity comes into play.

$$\begin{aligned} h(\alpha \xi) &= h(\lim \alpha \frac{x_n}{|x_n|}) \geq \limsup h(\alpha \frac{x_n}{|x_n|}) \\ &= \limsup h(\frac{\alpha}{|x_n|} x_n + (1 - \frac{\alpha}{|x_n|}) \cdot 0) \\ &\geq \limsup (\frac{\alpha}{|x_n|} h(x_n) + (1 - \frac{\alpha}{|x_n|}) h(0)) \\ &\geq \limsup (\frac{\alpha}{|x_n|} c + (1 - \frac{\alpha}{|x_n|}) h(0)) \\ &= h(0), \end{aligned}$$

which contradicts the assumption on h .

Knowing that $\{h \geq c\}$ is compact for all $c < \sup h$, we have

$$\{h = \sup h\} = \bigcap_{c < \sup h} \{h \geq c\},$$

which is an infinite intersection of nested non-empty compact sets. By compactness, this intersection is non-empty. \square

Lemma 6.4 *Let $u : D \rightarrow \mathbb{R}$ be a utility function, where $D = [a, \infty)$, $a < 0$. Let $0 \leq b < -a$. Let $X \geq 0$ be a random variable. Then for all $\alpha \in (0, 1]$ the implication*

$$\mathbb{E} u(\alpha X - b) < \infty \Rightarrow \mathbb{E} u(X) < \infty$$

holds.

Proof From concavity of u and $X \geq 0$ one obtains for $X > 0$ the following relations between slopes

$$\frac{u(X) - u(0)}{X} \leq \frac{u(\alpha X) - u(0)}{\alpha X} \leq \frac{u(\alpha X - b) - u(-b)}{\alpha X}.$$

Hence $u(X) - u(0) \leq (u(\alpha X - b) - u(-b))/\alpha$ (also valid if $X = 0$), from which the result follows. \square

Theorem 6.5 *Let $u : D \rightarrow \mathbb{R}$ be a utility function and Y the vector of discounted net gains. Let Assumption 6.2 be satisfied. A maximizer in Problem 6.1 exists if and only if the market is free of arbitrage. In this case the maximizer is unique if the market is non-redundant (see Definition 1.13).*

Proof Throughout the proof we suppose that the market is non-redundant, since any market can be reduced to such one in view of Proposition 1.14. First we consider uniqueness. Proposition 1.14(ii) tells us that in a non-redundant market the a.s. equality $\xi \cdot Y = \xi' \cdot Y$ implies that $\xi = \xi'$. Hence the function $\xi \mapsto u(\xi \cdot Y)$ is a.s. strictly concave, and then also $\xi \mapsto \mathbb{E} u(\xi \cdot Y)$ is strictly concave. Suppose that two maximizers ξ^* and ξ' exist, then by strict concavity $\mathbb{E} u(\frac{1}{2}(\xi^* + \xi') \cdot Y) > \frac{1}{2}(\mathbb{E} u(\xi^* \cdot Y) + \mathbb{E} u(\xi' \cdot Y)) = \mathbb{E} u(\xi^* \cdot Y)$, unless $\xi^* = \xi'$. Hence there can be at most one maximizer in this case.

We turn to existence. Suppose that the market admits an arbitrage opportunity. Let ξ be any risky portfolio. By Corollary 1.4, there exists a portfolio ξ' such that $\xi' \cdot Y \geq 0$ a.s. and $\mathbb{P}(\xi' \cdot Y > 0) > 0$. In this case one has $\mathbb{E} u((\xi + \xi') \cdot Y) > \mathbb{E} u(\xi \cdot Y)$ and therefore a maximizing portfolio cannot exist.

Assume that the market is free of arbitrage. Consider first the case where $D = \mathbb{R}$ and u has an upper bound. We will invoke Lemma 6.3 applied to the function $h(\xi) = \mathbb{E} u(\xi \cdot Y)$. We first show that h is u.s.c. Since u has an upper bound, we can apply the lim sup version of Fatou's lemma and obtain for every sequence (ξ^n) with limit ξ

$$\limsup h(\xi^n) = \limsup \mathbb{E} u(\xi^n \cdot Y) \leq \mathbb{E} \limsup u(\xi^n \cdot Y) = h(\xi),$$

by continuity of u . Next we check assumption (6.4). In view of Corollary 1.4, absence of arbitrage is equivalent to saying that for every $\xi \in \mathbb{R}^d \setminus \{0\}$ one has $\mathbb{P}(\xi \cdot Y < 0) > 0$. Indeed, $\mathbb{P}(\xi \cdot Y \geq 0) = 1$ implies $\mathbb{P}(\xi \cdot Y = 0) = 1$, which in turn implies $\xi = 0$ by non-redundancy. Hence if $\xi \neq 0$, then $\mathbb{P}(\xi \cdot Y < 0) > 0$.

Since u is concave and increasing we have that $\{\xi \cdot Y < 0\} = \{\lim_{\alpha \uparrow \infty} u(\alpha \xi \cdot Y) = -\infty\}$. From the fact that the latter set has positive probability, it follows by the Monotone convergence theorem (use also that u has an upper bound) that for all $\xi \neq 0$

$$\lim_{\alpha \rightarrow \infty} \mathbb{E} u(\alpha \xi \cdot Y) = -\infty.$$

We have shown that for the present case, absence of arbitrage leads to condition (6.4), which is sufficient for existence of a maximum of $\mathbb{E} u(\xi \cdot Y)$.

We turn to the case, where all $\xi \cdot Y$ involved have a lower bound $a < 0$. We show that $\Xi = \{\xi \in \mathbb{R}^d : \xi \cdot Y \geq a \text{ a.s.}\}$ is compact. We follow a familiar way of reasoning (see also the proof of Theorem 1.15). Supposing that this set is unbounded, we can take a sequence (ξ^n) in this set such that $|\xi^n| \rightarrow \infty$ and $\xi^n / |\xi^n| \rightarrow \eta$, for some vector $\eta >$, with $|\eta| = 1$. Then

$$\eta \cdot Y \geq \lim_{|\xi^n| \rightarrow \infty} \frac{a}{|\xi^n|} = 0 \text{ a.s.}$$

By absence of arbitrage and non-redundancy we conclude that $\eta = 0$, a contradiction. Hence we optimize over a compact set. To show that an optimizer exists, it is now sufficient to show that h is continuous on Ξ . This follows by application of the Dominated convergence theorem as soon as there is a random variable X such that $\sup_{\xi \in \Xi} u(\xi \cdot Y) \leq u(X)$ a.s. and $\mathbb{E} u(X) < \infty$, since u is lower bounded by $u(a)$.

Define $\eta \in \mathbb{R}_+^d$ by its elements $\eta_i = 0 \vee m_i$, where $m_i = \max\{\xi_i : \xi \in \Xi\}$. The m_i are finite by compactness of Ξ . By positivity of S , we have $\eta \cdot S \geq \xi \cdot S$ for $\xi \in \Xi$ and hence

$$\xi \cdot Y \leq \frac{\eta \cdot S}{1+r} - \xi \cdot \pi \leq \frac{\eta \cdot S}{1+r} + M =: X,$$

where $M = \max\{-\xi \cdot \pi : \xi \in \Xi\} \vee 0$ (also a finite number). We also have $\eta \cdot Y \geq -\eta \cdot \pi$ and, because $\eta \cdot \pi \geq 0$, there is $\alpha \in (0, 1]$ such that $\alpha \eta \cdot \pi < -a$, which implies that $\alpha \eta \cdot Y > a$. Then $\alpha \eta \in \Xi$ and, by assumption 6.2, $\mathbb{E} u(\alpha \eta \cdot Y) < \infty$. One has $a \leq \xi \cdot Y \leq X$ and $\alpha \eta \cdot Y = \alpha X - \alpha(\eta \cdot \pi + M)$. With $b = \alpha(\eta \cdot \pi + M)$ we wish to apply Lemma 6.4 to get $\mathbb{E} u(X) < \infty$, as desired. One then has to verify that $0 \leq \alpha(\eta \cdot \pi + M) < -a$, which can be accomplished by taking α small enough, meanwhile maintaining $\mathbb{E} u(\alpha \eta \cdot Y) < \infty$. This finishes the proof. \square

Knowing that under Assumption 6.2, the maximization problem has a solution, we now turn to a characterization of it under additional assumptions.

Theorem 6.6 *Let $u : D \rightarrow \mathbb{R}$ be a continuously differentiable utility function. Let Assumption 6.2 hold and assume, additionally, $\mathbb{E} |u(\xi \cdot Y)| < \infty$ for all*

$\xi \in \Xi$. Let the Problem 6.1 maximizing ξ^* be an interior point of Ξ . Then $Yu'(\xi^* \cdot Y) \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbb{P})$ and

$$(6.5) \quad \mathbb{E}Yu'(\xi^* \cdot Y) = 0.$$

Proof If differentiation and expectation commute, one has

$$\nabla_{\xi} \mathbb{E}u(\xi \cdot Y) = \mathbb{E}u'(\xi \cdot Y)Y^{\top},$$

and the result follows by taking $\xi = \xi^*$. We directly show that the right hand side is zero at $\xi = \xi^*$. Take $\eta \in \mathbb{R}^d$ and $\varepsilon \in (0, 1]$. Put $\xi_{\varepsilon} = \xi^* + \varepsilon\eta$, then $\xi_{\varepsilon} \in D$ for all ε sufficiently small, $\varepsilon < \varepsilon_0$ say. For those ε we put

$$\Delta_{\varepsilon} = \frac{u(\xi_{\varepsilon} \cdot Y) - u(\xi^* \cdot Y)}{\varepsilon},$$

and note that $\mathbb{E}\Delta_{\varepsilon} \leq 0$. Concavity of u gives that $f(\varepsilon) \equiv u(\xi_{\varepsilon} \cdot Y)$ is concave too. Hence Δ_{ε} is increasing for $\varepsilon \downarrow 0$, with limit $\eta \cdot Yu'(\xi^* \cdot Y)$. The assumption that $u(\xi \cdot Y) \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbb{P})$ for all $\xi \in \Xi$ implies that $\Delta_{\varepsilon_0} \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbb{P})$. Hence $\Delta_{\varepsilon} - \Delta_{\varepsilon_0}$ is nonnegative and increasing for $\varepsilon \downarrow 0$, which enables us to apply the Monotone convergence theorem to get

$$0 \geq \mathbb{E}\Delta_{\varepsilon} \uparrow \eta \cdot \mathbb{E}Yu'(\xi^* \cdot Y),$$

where the expectation on the right hand side is finite, since $\Delta_{\varepsilon_0} \leq \Delta_{\varepsilon} \leq 0$. We conclude that $\eta \cdot \mathbb{E}Yu'(\xi^* \cdot Y) \leq 0$ for all $\eta \in \mathbb{R}^d$. So we can replace η with $-\eta$ in the last inequality and we conclude that the linear map $\eta \mapsto \eta \cdot \mathbb{E}Yu'(\xi^* \cdot Y) = 0$ identically. But then we must have $\mathbb{E}Yu'(\xi^* \cdot Y) = 0$. \square

Proposition 6.7 *Let the assumptions of Theorem 6.6 hold and let the market be arbitrage-free. Let ξ^* be the maximizer of Problem 6.1. Then $\mathbb{E}u'(\xi^* \cdot Y) < \infty$ and*

$$(6.6) \quad \frac{d\mathbb{P}^*}{d\mathbb{P}} = \frac{u'(\xi^* \cdot Y)}{\mathbb{E}u'(\xi^* \cdot Y)}$$

defines a risk-neutral measure on (Ω, \mathcal{F}) .

Proof First we show that $\mathbb{E}u'(\xi^* \cdot Y) < \infty$, so that \mathbb{P}^* is well defined. Define

$$c := \sup\{u'(x) : x \in D \text{ and } x \in [-|\xi^*|, |\xi^*|]\}.$$

Consider first the case in which $D = \mathbb{R}$. Then, because u' is decreasing, we have $c = u'(-|\xi^*|)$. If $D = [a, \infty)$, then $c \leq u'(a)$. In both cases we have $c < \infty$. By the Cauchy-Schwartz inequality, we have $|\xi^* \cdot Y| \leq |\xi^*| \cdot |Y|$. Hence, if $|\xi^* \cdot Y| > |\xi^*|$, then $|Y| > 1$. From this it follows that

$$0 \leq u'(\xi^* \cdot Y) \leq c + u'(\xi^* \cdot Y)|Y|\mathbf{1}_{\{|Y|>1\}} \leq c + u'(\xi^* \cdot Y)|Y|,$$

where the expression on the right hand side has finite expectation, by Theorem 6.6.

By definition, a risk-neutral measure satisfies $\mathbb{E}^*Y = 0$. This is indeed the case, since

$$\mathbb{E}^*Y = \mathbb{E}Y \frac{d\mathbb{P}^*}{d\mathbb{P}} = 0,$$

because of Equation (6.5). \square

Remark 6.8 If Y is \mathbb{P} -a.s. bounded, then the Radon-Nikodym derivative in (6.6) is bounded and we have constructed a risk neutral measure with bounded density as mentioned in Theorem 1.6. If Y is not bounded under \mathbb{P} , one may change the optimization problem, by considering $\tilde{Y} = Y/(1 + |Y|)$, which is bounded, instead of Y . Indeed, along with Y , also \tilde{Y} satisfies the no arbitrage condition $\mathbb{P}(\xi \cdot \tilde{Y} \geq 0) = 1 \Rightarrow \mathbb{P}(\xi \cdot \tilde{Y} = 0) = 1$ and vice versa. If $\tilde{\xi}$ is the corresponding maximizer of $\xi \mapsto \mathbb{E}u(\tilde{\xi} \cdot Y)$, then $\frac{d\tilde{\mathbb{P}}}{d\mathbb{P}} = \frac{1}{c} \frac{u'(\tilde{\xi} \cdot \tilde{Y})}{1 + |Y|}$ defines a risk-neutral measure $\tilde{\mathbb{P}}$ equivalent to \mathbb{P} for $c = \mathbb{E} \frac{u'(\tilde{\xi} \cdot \tilde{Y})}{1 + |Y|}$.

6.2 Exponential utility and relative entropy

In the present section we fix the utility function to be given by $u(x) = 1 - \exp(-\alpha x)$, $x \in \mathbb{R}$ and $\alpha > 0$. The optimization problem 6.1 is in this case equivalent to the *minimization* of the function $Z : \mathbb{R}^d \rightarrow (0, \infty)$ defined by

$$Z(\lambda) = \mathbb{E} e^{\lambda \cdot Y}.$$

This optimization problem will be studied under the assumption that $Z(\lambda) < \infty$ for all $\lambda \in \mathbb{R}^d$, for which we have the following equivalent formulation, all exponential moments of $|Y|$ are finite.

Lemma 6.9 *It holds that $Z(\lambda) < \infty$, for all $\lambda \in \mathbb{R}^d$ iff $\mathbb{E} \exp(\alpha|Y|) < \infty$ for all $\alpha \in \mathbb{R}$.*

Proof The condition in terms of α is certainly sufficient. To prove necessity we proceed as follows. We use that $|Y| \leq \sqrt{d} \sum_i |Y_i|$ to get for $\alpha > 0$ (which is sufficient to consider) by Hölder's inequality for d random variables

$$\mathbb{E} e^{\alpha|Y|} \leq \mathbb{E} e^{\alpha\sqrt{d} \sum_i |Y_i|} \leq \prod_i (\mathbb{E} e^{\alpha d \sqrt{d} |Y_i|})^{1/d}.$$

Since $\exp(\alpha d \sqrt{d} |Y_i|) \leq \exp(\alpha d \sqrt{d} Y_i) + \exp(-\alpha d \sqrt{d} Y_i)$, each of the factors in the product is finite. \square

In the remainder of this section we assume that the condition of Lemma 6.9 holds. Before we proceed with the optimization problem, we introduce some terminology.

Definition 6.10 *The exponential family of \mathbb{P} with respect to Y is the family of probability measures \mathbb{P}_λ with $\lambda \in \mathbb{R}^d$ given by*

$$\frac{d\mathbb{P}_\lambda}{d\mathbb{P}} = \frac{e^{\lambda \cdot Y}}{Z(\lambda)}.$$

Expectation w.r.t. \mathbb{P}_λ is denoted by \mathbb{E}_λ and $m(\mathbb{P}_\lambda) := \mathbb{E}_\lambda Y$. Note that all \mathbb{P}_λ are mutually equivalent probability measures, and equivalent to \mathbb{P} .

We restate Theorem 6.5 in the present context.

Proposition 6.11 *The function $\lambda \mapsto Z(\lambda)$ takes on its minimum iff the market is arbitrage free. If this happens, any minimizer λ^* also solves the equation*

$$m(\mathbb{P}_{\lambda^*}) = 0.$$

If the market is non-redundant, then the minimizer is unique.

Proof We apply Theorem 6.5, and so a minimizer exists iff the market is free of arbitrage. From Theorem 6.6 we obtain for this case that $m(\mathbb{P}_{\lambda^*}) = 0$. \square

Below we will see a converse to this proposition, if $m(\mathbb{P}_{\lambda^*}) = 0$, then λ^* minimizes $\lambda \mapsto Z(\lambda)$.

Definition 6.12 Let \mathbb{P} and \mathbb{Q} be two probability measures on a measurable space (Ω, \mathcal{F}) . Denote by $\mathbb{E}_\mathbb{Q}$ expectation under \mathbb{Q} . If $\mathbb{Q} \ll \mathbb{P}$, then the *relative entropy*, or *Kullback-Leibler information* of \mathbb{Q} w.r.t. \mathbb{P} is defined as

$$H(\mathbb{Q}|\mathbb{P}) = \mathbb{E}_\mathbb{Q} \log \frac{d\mathbb{Q}}{d\mathbb{P}} \leq \infty.$$

If \mathbb{Q} is not absolutely continuous w.r.t. \mathbb{P} , then $H(\mathbb{Q}|\mathbb{P}) := \infty$.

Since \log is strictly concave, it follows from Jensen's inequality that always $H(\mathbb{Q}|\mathbb{P}) \geq 0$ and that $H(\mathbb{Q}|\mathbb{P}) = 0$ iff $\mathbb{Q} = \mathbb{P}$.

Proposition 6.13 *Assume there is $\lambda_0 \in \mathbb{R}^d$ such that $m(\mathbb{P}_{\lambda_0}) = 0$.*

- (i) *For all $\lambda \in \mathbb{R}^d$ it holds that $H(\mathbb{P}_\lambda|\mathbb{P}) = \lambda \cdot m(\mathbb{P}_\lambda) - \log Z(\lambda)$.*
- (ii) *If \mathbb{Q} is a probability measure on (Ω, \mathcal{F}) with $\mathbb{E}_\mathbb{Q} Y = 0$, then $H(\mathbb{Q}|\mathbb{P}) = H(\mathbb{Q}|\mathbb{P}_{\lambda_0}) + H(\mathbb{P}_{\lambda_0}|\mathbb{P})$.*
- (iii) *Let \mathcal{Q}_0 be the set of probability measures \mathbb{Q} with $\mathbb{E}_\mathbb{Q} Y = 0$. Then the mapping $\mathbb{Q} \mapsto H(\mathbb{Q}|\mathbb{P})$ assumes on the set \mathcal{Q}_0 a unique minimum for $\mathbb{Q} = \mathbb{P}_{\lambda_0}$.*
- (iv) *λ_0 is the minimizer of $\lambda \mapsto Z(\lambda)$.*

Proof (i) By definition of \mathbb{P}_λ one has $\log \frac{d\mathbb{P}_\lambda}{d\mathbb{P}} = \lambda \cdot Y - \log Z(\lambda)$. The result then follows, because $\mathbb{E}_\lambda Y = m(\mathbb{P}_\lambda)$.

(ii) Clearly, there is only something to prove if all quantities involved are finite. So we assume $\mathbb{Q} \ll \mathbb{P}$, and then we also have $\mathbb{Q} \ll \mathbb{P}_\lambda$. From the product rule

$$\frac{d\mathbb{Q}}{d\mathbb{P}} = \frac{d\mathbb{Q}}{d\mathbb{P}_\lambda} \frac{d\mathbb{P}_\lambda}{d\mathbb{P}} = \frac{d\mathbb{Q}}{d\mathbb{P}_\lambda} \frac{e^{\lambda \cdot Y}}{Z(\lambda)}$$

one obtains

$$\log \frac{d\mathbb{Q}}{d\mathbb{P}} = \log \frac{d\mathbb{Q}}{d\mathbb{P}_\lambda} + \lambda \cdot Y - \log Z(\lambda).$$

Take expectation under \mathbb{Q} to get

$$(6.7) \quad \begin{aligned} H(\mathbb{Q}|\mathbb{P}) &= H(\mathbb{Q}|\mathbb{P}_\lambda) + \lambda \cdot \mathbb{E}_{\mathbb{Q}} Y - \log Z(\lambda) \\ &= H(\mathbb{Q}|\mathbb{P}_\lambda) - \log Z(\lambda), \end{aligned}$$

since $\mathbb{E}_{\mathbb{Q}} Y = 0$. The result now follows from (i) if we take $\lambda = \lambda_0$.

(iii) Note that $\mathbb{P}_{\lambda_0} \in \mathcal{Q}_0$. It follows from (ii) that $H(\mathbb{Q}|\mathbb{P}) \geq H(\mathbb{P}_{\lambda_0}|\mathbb{P})$ for all $\mathbb{Q} \in \mathcal{Q}_0$. Equality holds iff $H(\mathbb{Q}|\mathbb{P}_{\lambda_0}) = 0$, which happens iff $\mathbb{Q} = \mathbb{P}_{\lambda_0}$.

(iv) Take in (6.7) $\mathbb{Q} = \mathbb{P}_{\lambda_0}$ to obtain

$$H(\mathbb{P}_{\lambda_0}|\mathbb{P}_\lambda) = H(\mathbb{P}_{\lambda_0}|\mathbb{P}) + \log Z(\lambda).$$

Then minimizing $Z(\lambda)$ over λ is equivalent to minimizing $H(\mathbb{P}_{\lambda_0}|\mathbb{P}_\lambda)$. But a minimizer of the latter is λ_0 . \square

We close this section by connecting the preceding results for portfolio optimization to the construction of a special risk neutral measure.

Corollary 6.14 *Suppose that the market is arbitrage-free under the probability measure \mathbb{P} . Then there exists a unique risk-neutral measure \mathbb{P}^* that minimizes the relative entropy $H(\mathbb{P}'|\mathbb{P})$ over all equivalent risk-neutral measures $\mathbb{P}' \in \mathcal{P}$. Specifically, if λ^* is the minimizer of $Z(\lambda)$, then*

$$(6.8) \quad \frac{d\mathbb{P}^*}{d\mathbb{P}} = \frac{e^{\lambda^* \cdot Y}}{\mathbb{E} e^{\lambda^* \cdot Y}}.$$

Proof We apply Proposition 6.13 together with Proposition 6.11 to obtain the result. \square

The assertion of Corollary 6.14 can be restated by saying that the optimal portfolio ξ^* of an optimization problem in terms of a CARA utility function can be characterized by a *relative entropy minimizing probability measure* \mathbb{P}^* . This measure, as presented in this corollary, is sometimes called an *exponentially tilted transformation* of \mathbb{P} , or an *Esscher transform* of \mathbb{P} .

6.3 Exercises

6.1 Show that a function $h : \mathbb{R}^d \rightarrow \mathbb{R}$, defined on some domain is upper semi-continuous iff the sets $h^{-1}(-\infty, a)$ are open for all $a \in \mathbb{R}$.

6.2 The assertion of Theorem 6.5 should also be true for a utility function $u : (a, \infty) \rightarrow \mathbb{R}$ with $a < 0$, $\lim_{x \downarrow a} u(x) = -\infty$ and u bounded from above. Investigate whether this conjecture is correct. In particular one should check whether h is upper semicontinuous and adapt the proof here and there. It could be useful to extend the definition of u by defining $u(x) = -\infty$ for $x \leq a$, since otherwise certain desirable properties of $h(\xi) = \mathbb{E} u(\xi \cdot Y)$ are hard to establish.

6.3 Consider the CARA utility function $u(x) = 1 - \exp(-\alpha x)$, $x \in \mathbb{R}$, with $\alpha > 0$, the constant Arrow-Pratt coefficient.

- (a) Show that the condition $\mathbb{E}|u(\xi \cdot Y)| < \infty$ for $\xi \in \Xi$ of Theorem 6.6 is equivalent to $\mathbb{E} \exp(\xi \cdot Y) < \infty$, for all $\xi \in \mathbb{R}^d$.
- (b) Show that the risk-neutral measure \mathbb{P}^* of Proposition 6.7 is the same for all $\alpha > 0$.
- (c) Suppose that Y has a d -dimensional multivariate normal distribution with mean vector m and invertible covariance matrix Σ . Compute the optimal $\xi^* \in \mathbb{R}^d$.

6.4 Let \mathbb{P} be a probability measure on some measurable space (Ω, \mathcal{F}) . Show that the mapping $\mathbb{Q} \mapsto H(\mathbb{Q}|\mathbb{P})$ is strictly convex on the set of all probability measures on this space such that $H(\mathbb{Q}|\mathbb{P}) < \infty$.

6.5 Consider a market with one risky asset only ($d = 1$). Let the assumptions of Theorem 6.6 hold. Let ξ^* be the optimal investment in the risky asset. Show, by inspecting the objective function $\xi \mapsto \mathbb{E} u(\xi \cdot Y)$ (u defined on \mathbb{R} and differentiable) near $\xi = 0$, that $\xi^* > 0$ iff $\mathbb{E} Y > 0$. (This yields an alternative to Example 4.7).

6.6 Consider a market with $r = 0$ and one risky good S_1 , that can take on arbitrarily large and arbitrarily small numbers, $\mathbb{P}(S_1 \leq \varepsilon) > 0$ and $\mathbb{P}(S_1 \geq 1/\varepsilon) > 0$ for any $\varepsilon > 0$ and assume $\mathbb{E} S_1 < \pi_1$. Let $D = [-\pi_1, \infty)$ and u a sufficiently smooth utility function with domain D . Note that $Y \geq -\pi_1$, hence $u(Y)$ is well defined.

- (a) Show that $\Xi = [0, 1]$.
- (b) From Example 4.7 one deduces $\xi^* = 0$, a boundary point of Ξ . Show that $\mathbb{E} u'(\xi^* \cdot Y) Y < 0$, violating Condition 6.5.

6.7 Given a utility function \tilde{u} as at the beginning of this section, the transformed utility function u depends on the initial capital w . In general, an optimal portfolio will also depend on w . We study this a bit for the case $d = 1$ and (for simplicity) $r = 0$. We avoid redundancy of the market, Y is non-degenerate. Assume that \tilde{u} is a C^2 function and that everywhere below interchanging of expectation and differentiation is allowed. Put $f(w, \xi) = \mathbb{E} \tilde{u}'(\xi Y + w) Y$.

- (a) Show that $\frac{\partial f}{\partial \xi}(w, x) < 0$.
- (b) Conclude that (locally) for every $w > 0$, there is a C^1 function $w \mapsto \xi^*(w)$ such that $f(w, \xi^*(w)) = 0$.
- (c) Show that

$$\frac{d\xi^*(w)}{dw} = - \frac{\mathbb{E} \tilde{u}''(\xi^*(w) Y + w) Y}{\mathbb{E} \tilde{u}''(\xi^*(w) Y + w) Y^2}.$$

- (d) Assume that $\mathbb{E} Y > 0$ and that Arrow-Pratt coefficient $\tilde{\alpha}(\cdot)$ of \tilde{u} is a decreasing function. Show that $Y \tilde{\alpha}(\xi^*(w) Y + w) \leq Y \tilde{\alpha}(w)$.
- (e) Conclude, under the assumptions in (d), that $\xi^*(\cdot)$ is an increasing function of w . (In Micro-economics, assets with the latter property are called *normal goods*. Assets with *decreasing demand* ξ^* are called *inferior goods*.)

6.8 Lemma 6.3 also has a converse. If h is strictly concave and upper semi-continuous, then existence of a minimizer implies (6.4). Show this and give an example that shows that the *strict* concavity of h cannot be missed here.

6.9 Consider a market with one risky good, its value at $t = 1$ is S and price π (at $t = 0$). Assume that S has under \mathbb{P} a Poisson distribution with parameter $\alpha > 0$. Consider the exponential family of Definition 6.10.

- (a) Show that $Z(\lambda) < \infty$ for all $\lambda \in \mathbb{R}$
- (b) Show that S has a Poisson distribution with parameter αe^λ under \mathbb{P}_λ .
- (c) Compute the minimizer of $\lambda \mapsto Z(\lambda)$ directly.
- (d) Verify that the minimizer is in agreement with Proposition 6.13.

7 Optimal contingent claims

In the previous sections we studied the problem of portfolio optimization. In a complete market, every contingent claim (recall Definition 1.10) has the same pay-off as some portfolio, but in an incomplete market this is no longer true. Therefore, in the latter case, it makes sense to study the maximization of the expected utility $\mathbb{E}u(X)$, where u is a utility function and X some contingent claim, belonging to some suitable convex set \mathcal{X} . The specification of \mathcal{X} will depend on the context.

7.1 An expected utility optimization problem

Let w be the initial capital of some investor. Let \mathbb{P}^* be a probability measure, equivalent to \mathbb{P} with Radon-Nikodym derivative $\frac{d\mathbb{P}^*}{d\mathbb{P}} = \phi$. Given a contingent claim with *discounted pay-off* X , define a *pricing rule* by $\mathbb{E}^*X = \mathbb{E}\phi X$.

In this context, ϕ is also called a *pricing kernel*. The *budget constraint* on claims X in the present context is given by $\mathbb{E}^*X \leq w$. We introduce the *budget set*, also called the set of *admissible pay-offs*,

$$\mathcal{B} = \{X \in \mathcal{X} \cap L^1(\Omega, \mathcal{F}, \mathbb{P}^*) : \mathbb{E}^*X \leq w\}.$$

Note that \mathcal{B} is a convex set. We will study the following

Problem 7.1 Maximize $\mathbb{E}u(X)$ over the set \mathcal{B} .

To have this problem well-defined, we need that $\mathbb{P}^* \sim \mathbb{P}$, not just $\mathbb{P}^* \ll \mathbb{P}$ or $\mathbb{P} \ll \mathbb{P}^*$, see Exercise 7.1. Note that this relates to Theorem 6.5, where a similar situation has been encountered and where we required the market to be arbitrage-free, which is equivalent to the existence of a Risk neutral measure. We will also assume that $\mathbb{P}(X \in D) = 1$, for all $X \in \mathcal{B}$, where D is the domain of u .

If Problem 7.1 has a solution, it is necessarily unique. This is due to the fact that u is strictly concave, see the proof of Theorem 6.5. Another fact that we encountered before is, that it can never be optimal to invest less than the initial capital w . Indeed, if a given claim $X \in \mathcal{B}$ is such that $\mathbb{E}^*X < w$, then $X' = X + w - \mathbb{E}^*X > X$ and so $\mathbb{E}u(X') > \mathbb{E}u(X)$, whereas $X' \in \mathcal{B}$, since $\mathbb{E}^*X' = w$.

For the time being, we drop the budget constraint and let \mathcal{X} be the set of all random variables. Suppose that X^* is the optimal claim. Let X be any bounded random variable and consider the ‘perturbed’ claims, belonging to \mathcal{B} as well,

$$X_\lambda = X^* + \lambda(X - \mathbb{E}^*X), \lambda \in \mathbb{R}.$$

Among the pay-offs X_λ , the optimal one is found for $\lambda = 0$. Hence

$$\frac{d}{d\lambda} \mathbb{E}u(X_\lambda) = 0, \text{ for } \lambda = 0.$$

Scrupulously interchanging differentiation and expectation in the above equation yields

$$\begin{aligned} 0 &= \mathbb{E}(u'(X^*)(X - \mathbb{E}^* X)) \\ &= \mathbb{E} X(u'(X^*) - \phi \mathbb{E} u'(X^*)). \end{aligned}$$

Let $c = \mathbb{E} u'(X^*)$, then the above identity yields

$$\mathbb{E} X u'(X^*) = c \mathbb{E}(X \phi),$$

valid for all bounded X , in particular for $X = \mathbf{1}_F$, $F \in \mathcal{F}$. This means that both sides of the above equality define the same measures on \mathcal{F} , absolutely continuous w.r.t. \mathbb{P} and hence the Radon-Nikodym derivatives are the same. In other words,

$$u'(X^*) = c \phi,$$

yielding $X^* = (u')^{-1}(c \phi)$. Hence we have found a candidate solution. The heuristics above are justified by the following theorem.

Theorem 7.2 *Suppose that $u \in C^1(\mathbb{R})$, $\lim_{x \rightarrow -\infty} u'(x) = \infty$ and u bounded from above. Let I be the inverse of the function u' , defined on the range of u' . Let $c > 0$ and $X^* := I(c \phi)$. Then X^* is well defined a.s. Moreover, assume that $X^* \in L^1(\Omega, \mathcal{F}, \mathbb{P}^*)$ and let $w = \mathbb{E}^* X^*$. Then X^* is the unique maximizer of Problem 7.1 for $\mathcal{X} = L^0(\Omega, \mathcal{F}, \mathbb{P})$.*

Proof We have already discussed uniqueness and so we turn to existence. It follows from the assumptions that $u'(x) \rightarrow 0$ for $x \rightarrow \infty$. Hence every positive number is in the range of u' . Since $\mathbb{P}^* \sim \mathbb{P}$, we have that $\mathbb{P}(\phi > 0) = 1$. Hence $I(c \phi)$ is \mathbb{P} -a.s. well-defined.

Concavity of u yields for any $X \in L^1(\Omega, \mathcal{F}, \mathbb{P}^*)$ that

$$u(X) \leq u(X^*) + u'(X^*)(X - X^*) = u(X^*) + c \phi(X - X^*).$$

Taking expectations in this inequality yields

$$\begin{aligned} \mathbb{E} u(X) &\leq \mathbb{E} u(X^*) + c \mathbb{E} \phi(X - X^*) \\ &= \mathbb{E} u(X^*) + c \mathbb{E}^*(X - X^*) \\ &= \mathbb{E} u(X^*) + c(\mathbb{E}^* X - w) \\ &\leq \mathbb{E} u(X^*), \end{aligned}$$

which shows that X^* is the maximizer. □

Let W be a nonnegative random variable with values in $[0, \infty]$. Until further notice we assume that the set \mathcal{X} is that of random variables X satisfying $0 \leq X \leq W$ a.s. In this case we assume that $u : [0, \infty) \rightarrow \mathbb{R}$.

Remark 7.3 One can show that for any utility function u , there exists a maximizer $X^* \in \mathcal{B}$ of $\mathbb{E} u(X)$ under the conditions stipulated above. For a constructive result we will narrow the class of utility functions under consideration by assuming differentiability.

Let $u \in C^1(0, \infty)$ be a utility function. We can extend the domain of u to $[0, \infty)$, by setting $u(0) = \lim_{x \rightarrow 0} u(x) \geq -\infty$. Since u' is decreasing, the limits

$$a = \lim_{x \rightarrow \infty} u'(x)$$

and

$$b = \lim_{x \rightarrow 0} u'(x)$$

exist. Moreover, we have $0 \leq a < b \leq \infty$. On the open interval (a, b) , the function u' has a well-defined continuous and decreasing inverse I . We extend I to a function $I^+ : [0, \infty] \rightarrow [0, \infty]$, by setting

$$I^+(y) = \begin{cases} +\infty & \text{if } 0 \leq y \leq a \\ I(y) & \text{if } a < y < b \\ 0 & \text{if } y \geq b. \end{cases}$$

It is obvious that I^+ is decreasing and continuous on $[0, \infty]$.

Theorem 7.4 *Consider the optimization problem under the restriction $0 \leq X \leq W \leq \infty$. Let $X^* = I^+(c\phi) \wedge W$ and let $w = \mathbb{E}^* X^* < \infty$. If $\mathbb{E} u(X^*) < \infty$, then X^* is the unique maximizer of $\mathbb{E} u(X)$ over $X \in \mathcal{B}$.*

Proof Consider the function $v : \mathbb{R} \times \Omega \rightarrow \mathbb{R}$ (the Legendre-Fenchel transform of u , a common tool in convex analysis) defined by

$$(7.1) \quad v(y, \omega) = \sup\{u(x) - xy : 0 \leq x \leq W(\omega)\}.$$

Suppose that $W(\omega) < \infty$. By continuity of u , for each y and ω the supremum will be attained at some $x^*(y, \omega) \in [0, W(\omega)]$, which is unique by strict concavity of u . We discern three cases.

Suppose $x^*(y, \omega) = 0$. Then for all $x \in (0, W(\omega)]$ we have $u(x) - xy < u(0)$. Consider $\Delta_h u(x) := \frac{u(x+h) - u(x)}{h}$ for $x > 0$ and $x + h > 0$. By concavity of u , $\Delta_h u$ is decreasing for every fixed h , in particular $\Delta_h u(x) \leq \Delta_h u(0) = \frac{u(h) - u(0)}{h} \leq y$. It follows that $\sup_{x > 0} u'(x) \leq y$ and hence $y \geq b$. Conversely, if $y \geq b$, then $u'(x) < y$ for all $x \in (0, W(\omega))$, then $x \mapsto u(x) - xy$ is decreasing and $x^*(y, \omega) = 0$. One similarly shows that $x^*(y, \omega) = W(\omega)$ iff $y \leq a$. If $x^*(y, \omega) \in (0, W(\omega))$ is an interior maximizer, then $y = u'(x^*(y, \omega))$ holds and $y \in (a, b)$, so $x^*(y, \omega) = I(y)$. Taking all three cases into account, one arrives at

$$x^*(y, \omega) = I^+(y) \wedge W(\omega),$$

which is a measurable function, jointly in (y, ω) , since I^+ is continuous. Hence, by definition of X^* , one has $X^*(\omega) = x^*(c\phi(\omega), \omega)$ on $\{\omega : W(\omega) < \infty\}$, measurable in ω . Suppressing the dependence on ω , we thus found $X^* = x^*(c\phi)$ on $\{W < \infty\}$.

Consider now the case $W(\omega) = \infty$. The case where a finite maximizer $x^*(y, \omega)$ exists can be treated as before and one gets $x^*(y, \omega) = I^+(y) = I^+(y) \wedge W(\omega)$. The supremum in (7.1) is not attained for a finite argument, iff $u'(x) > y$

for all $x > 0$, and then $y \leq a$. By definition of I^+ , we can put $x^*(y, \omega) = I^+(y) = I^+(y) \wedge W(\omega)$. On the other hand, the assumption on X^* implies $w \geq \mathbb{E}^* X^* \mathbf{1}_{\{W=\infty\}} = \mathbb{E}^* I^+(c\phi) \mathbf{1}_{\{W=\infty\}}$. It follows that $X^* = x^*(c\phi)$ is finite a.s. on $\{W = \infty\}$. Combining this with the previous case, we obtain $X^* = x^*(c\phi)$, which is an a.s. finite random variable. But then, using the definition of x^* as the maximizer of (7.1), we get for arbitrary $X \in \mathcal{B}$

$$u(X^*) - c\phi X^* \geq u(X) - c\phi X \text{ a.s.}$$

Take expectations to get

$$\mathbb{E} u(X^*) - c\mathbb{E}\phi X^* \geq \mathbb{E} u(X) - c\mathbb{E}\phi X,$$

equivalent to

$$\mathbb{E} u(X^*) - \mathbb{E} u(X) \geq c(\mathbb{E}^* X^* - \mathbb{E}^* X),$$

which is nonnegative by $X \in \mathcal{B}$. The uniqueness issue has already been addressed before. \square

The previous theorems dealt with the existence of an optimizer for problems where the initial capital w was defined in terms of a property of the candidate optimizer, involving the constant c that was also depending on the candidate optimizer. In a practical situation, w is given before hand and so we can apply the previous theorems only if c is such that the assumptions in these theorems are met. The next corollary gives a simple sufficient condition for this.

Corollary 7.5 *Let $w > 0$ be given and assume $0 < w < \mathbb{E}^* W < \infty$ and $\mathbb{E} u(W) < \infty$. Then there exists a unique constant $c^* > 0$ such that $X^* := I^+(c^* \phi) \wedge W$ satisfies $\mathbb{E}^* X^* = w$. Hence X^* is the maximizer of $\mathbb{E} u(X)$ over $X \in \mathcal{B}$.*

Proof Let $\beta > 0$ and define f_β by $f_\beta(y) = I^+(y) \wedge \beta$. Then f_β is bounded, continuous and decreasing. Moreover, $\lim_{y \uparrow \beta} f_\beta(y) = 0$ and $f_\beta(y) = \beta$ for $y \leq u'(\beta)$. Put $g(c) = \mathbb{E}^* f_W(c\phi) = \mathbb{E}^* I^+(c\phi) \wedge W \leq \mathbb{E}^* W$. By dominated convergence, g is continuous. Furthermore, $\lim_{c \rightarrow \infty} g(c) = 0$, $\lim_{c \downarrow 0} g(c) = \mathbb{E}^* W$ and g is *strictly* decreasing (Exercise 7.5) on the interval $g^{-1}[(0, \mathbb{E}^* W)]$, which contains w by assumption. We thus obtain that there exists a unique c^* such that $w = g(c^*)$. Theorem 7.4 yields that X^* is the expected utility maximizer, as $\mathbb{E} u(X^*) < \infty$ is guaranteed by $\mathbb{E} u(W) < \infty$. \square

7.2 Optimization under uniform order restrictions

In this section we study an optimization problem, that involves the uniform order \succeq_{uni} , recall Definition 5.1. We transplant this order to the space of random variables, by saying that $X \succeq_{\text{uni}} Y$ iff $\mu_X \succeq_{\text{uni}} \mu_Y$, where μ_X and μ_Y denote the laws of X and Y respectively. The problem we are going to address is the following.

Problem 7.6 Let $\mathbb{P}^* \sim \mathbb{P}$ and $X_0 \in \mathcal{X} = L_+^1(\Omega, \mathcal{F}, \mathbb{P})$ be given. Note that $X_0 \geq 0$ a.s. under \mathbb{P} and \mathbb{P}^* and assume that $\mathbb{E}^* X_0 < \infty$. The objective is to minimize $\mathbb{E}^* X$ over all random variables $X \in \mathcal{X}$ satisfying $X \succeq_{\text{uni}} X_0$.

The interpretation of this problem is that one wants to find the minimal budget needed among all X that are at least as attractive as X_0 , in the sense that $X \succeq_{\text{uni}} X_0$. Note that the latter requirement is stated in terms of \mathbb{P} , whereas we want to find a minimal expectation under \mathbb{P}^* .

Before we state a theorem with the solution to this problem, we need some additional properties of the \succeq_{uni} order in terms of *quantile functions*.

Definition 7.7 If F is a distribution function, then $q : (0, 1) \rightarrow \mathbb{R}$ is called a *quantile function* for F if for all $t \in (0, 1)$ it holds that $F(q(t)-) \leq t \leq F(q(t))$. If X is a random variable with distribution function F , we also say that q is a quantile function for X if q is a quantile function of F . Such a quantile function is also denoted q_F and q_X .

Recall that there are two ‘extremal’ quantile functions, q^- and q^+ , defined by $q^-(t) = \sup\{x \in \mathbb{R} : F(x) < t\}$ and $q^+(t) = \sup\{x \in \mathbb{R} : F(x) \leq t\}$. Recall also the fundamental equivalences $q^-(t) \leq x \Leftrightarrow t \leq F(x)$ and $q^+(t) < x \Leftrightarrow t < F(x)$. Moreover, $q^+ = q^-$ a.e. w.r.t. Lebesgue measure. Since any quantile function q satisfies $q^- \leq q \leq q^+$, we also have $q = q^- = q^+$ a.e. w.r.t. Lebesgue measure, and hence integrals of these functions w.r.t. Lebesgue measure have the same value. In particular, if U has a uniform distribution on $(0, 1)$, $q(U)$ has distribution function F for any q that is a quantile function for F .

Lemma 7.8 Let F be a distribution function of a distribution with finite mean and q an associated quantile function. Then for all $x \in \mathbb{R}$ it holds that

$$xF(x) = \int_{-\infty}^x F(u) \, du + \int_0^{F(x)} q(u) \, du.$$

Moreover, for arbitrary $x \in \mathbb{R}$ and $t \in (0, 1)$, one has

$$(7.2) \quad xt \leq \int_{-\infty}^x F(u) \, du + \int_0^t q(u) \, du.$$

Proof Both relations follow from maximizing $x \mapsto xt - \int_{-\infty}^x F(u) \, du$ (Exercise 7.7). \square

Remark 7.9 Equation (7.2) is also valid for $t = 0, 1$ and if the distribution doesn’t have a finite mean, in which case the right hand may be infinite.

Lemma 7.10 Let μ, ν be probability measures on \mathbb{R} and let q_μ and q_ν be corresponding quantile functions. The following statements are equivalent.

- (i) $\mu \succeq_{\text{uni}} \nu$.
- (ii) For all $t \in (0, 1)$, it holds that $\int_0^t q_\mu(s) \, ds \geq \int_0^t q_\nu(s) \, ds$.

(iii) For all decreasing functions $h : (0, 1) \rightarrow [0, \infty)$ it holds that

$$(7.3) \quad \int_0^1 h(s)q_\mu(s) ds \geq \int_0^1 h(s)q_\nu(s) ds.$$

(iv) For all bounded decreasing functions $h : (0, 1) \rightarrow [0, \infty)$ inequality (7.3) holds true.

Proof (i) \Leftrightarrow (ii) follows from Lemma 7.8 and Theorem 5.3 (Exercise 7.8).

(ii) \Rightarrow (iii): Since h is decreasing, it has at most countably many discontinuities, so the integrals in (7.3) don't change if we replace h with its right-continuous modification. Then, up to an additive positive constant, h can be seen as the 'complement of a distribution function' of a measure η on $(0, 1)$, $h(t) = \eta(t, 1)$. We apply Fubini's theorem as in the proof of Theorem 5.3. We have

$$\begin{aligned} \int_0^1 h(t)q_\mu(t) dt &= \int_0^1 \int_{(t,1)} \eta(ds) q_\mu(t) dt \\ &= \int_{(0,1)} \int_0^s q_\mu(t) dt \eta(ds) \\ &\geq \int_{(0,1)} \int_0^s q_\nu(t) dt \eta(ds) \\ &= \int_0^1 h(s)q_\nu(s) ds. \end{aligned}$$

(iii) \Rightarrow (iv): trivial.

(iv) \Rightarrow (ii): Take $h = \mathbf{1}_{(0,t]}$. □

Lemma 7.11 Let X, Y be nonnegative random variables. Then

$$\mathbb{E}XY \geq \int_0^1 q_X(1-t)q_Y(t) dt,$$

where q_X and q_Y are quantile functions for X and Y respectively.

Proof First we note that by Fubini it holds that

$$\begin{aligned} \mathbb{E}XY &= \mathbb{E} \int_0^\infty \int_0^\infty \mathbf{1}_{\{x < X, y < Y\}} dx dy \\ &= \int_0^\infty \int_0^\infty \mathbb{P}(X > x, Y > y) dx dy. \end{aligned}$$

Next we have the trivial relations

$$\begin{aligned} \mathbb{P}(X > x, Y > y) &= \mathbb{P}(X > x) - \mathbb{P}(X > x, Y \leq y) \\ &\geq \mathbb{P}(X > x) - \mathbb{P}(Y \leq y). \end{aligned}$$

Since the extreme term on the left is nonnegative, we also have

$$\mathbb{P}(X > x, Y > y) \geq (\mathbb{P}(X > x) - \mathbb{P}(Y \leq y))^+.$$

For $F_Y(y) \leq 1 - F_X(x)$ we have

$$\begin{aligned} \mathbb{P}(X > x) - \mathbb{P}(Y \leq y) &= \int_0^1 \mathbf{1}_{\{F_Y(y) \leq t \leq 1 - F_X(x)\}} dt \\ &= \int_0^1 \mathbf{1}_{\{y \leq q_Y^+(t), x \leq q_X^+(1-t)\}} dt. \end{aligned}$$

In case $F_Y(y) > 1 - F_X(x)$, the integrand on the right hand side is zero, hence we can replace the left hand side with $(\mathbb{P}(X > x) - \mathbb{P}(Y \leq y))^+$, whatever x and y , meanwhile maintaining the integral expression. Integrating the right hand side with respect to x and y yields by Fubini's theorem

$$\int_0^1 \int_0^\infty \int_0^\infty \mathbf{1}_{\{y \leq q_Y^+(t), x \leq q_X^+(1-t)\}} dx dy dt = \int_0^1 q_X^+(1-t) q_Y^+(t) dt,$$

which proves the assertion, since all quantile function are Lebesgue-a.e. the same. \square

Theorem 7.12 *Let ϕ denote the Radon-Nikodym derivative $\frac{d\mathbb{P}^*}{d\mathbb{P}}$. Consider problem 7.6. If $X \in \mathcal{X}$ satisfies $X \succeq_{\text{uni}} X_0$, then*

$$(7.4) \quad \mathbb{E}^* X \geq \int_0^1 q_\phi(1-s) q_{X_0}(s) ds.$$

Let μ_ϕ be the law of ϕ under \mathbb{P} and F_ϕ its distribution function. If ν is the measure on $(\mathbb{R}, \mathcal{B})$ characterized by $\nu(-\infty, x] = \int_0^{F_\phi(x)} q_{X_0}(1-t) dt$, then $\nu \ll \mu_\phi$ and equality in (7.4) holds for $X = X^* := f(\phi)$ with $f = \frac{d\nu}{d\mu_\phi}$. Moreover, $X^* \succeq_{\text{uni}} X_0$ and hence X^* is the minimizer sought for. The function f has the following explicit expression.

$$(7.5) \quad f(x) = \begin{cases} q_{X_0}(1 - F_\phi(x)) & \text{if } x \text{ is a continuity point of } F_\phi \\ \frac{\int_{F_\phi(x-)}^{F_\phi(x)} q_{X_0}(1-t) dt}{F_\phi(x) - F_\phi(x-)} & \text{else.} \end{cases}$$

Proof We will use the auxiliary probability space $((0, 1), \mathcal{B}(0, 1), \lambda)$, where λ denotes Lebesgue measure. Expectations and conditional expectations w.r.t. λ will be denoted by expressions like $\mathbb{E}_\lambda(X)$, $\mathbb{E}_\lambda[X|Y]$. Furthermore let U be the identity mapping on $(0, 1)$; it has the uniform distribution on $(0, 1)$. Then $\tilde{\phi} = q_\phi(U)$ has the same distribution as ϕ . A property of conditional expectation yields that there exists a Borel-measurable function f such that

$$(7.6) \quad \mathbb{E}_\lambda[q_{X_0}(1-U)|\tilde{\phi}] = f(\tilde{\phi}), \lambda\text{-a.s.}$$

Put $X^* = f(\phi)$, then $X^* \stackrel{d}{=} f(\tilde{\phi})$. Hence we have, using Jensen's inequality for conditional expectations,

$$\begin{aligned}\mathbb{E} u(X^*) &= \mathbb{E}_\lambda(u(f(\tilde{\phi}))) \\ &= \mathbb{E}_\lambda(u(\mathbb{E}_\lambda[q_{X_0}(1-U)|\tilde{\phi}])) \\ &\geq \mathbb{E}_\lambda(\mathbb{E}_\lambda[u(q_{X_0}(1-U))|\tilde{\phi}]) \\ &= \mathbb{E}_\lambda(u(q_{X_0}(1-U))) \\ &= \mathbb{E} u(X_0),\end{aligned}$$

which shows that $X^* \succeq_{\text{uni}} X_0$. Likewise we compute

$$\begin{aligned}\mathbb{E}^* X^* &= \mathbb{E}(X^* \phi) = \mathbb{E}(f(\phi)\phi) = \mathbb{E}_\lambda(f(\tilde{\phi})\tilde{\phi}) \\ &= \mathbb{E}_\lambda(\mathbb{E}_\lambda[q_{X_0}(1-U)|\tilde{\phi}]\tilde{\phi}) \\ &= \mathbb{E}_\lambda(\mathbb{E}_\lambda[q_{X_0}(1-U)\tilde{\phi}|\tilde{\phi}]) \\ &= \mathbb{E}_\lambda(q_{X_0}(1-U)\tilde{\phi}) \\ &= \mathbb{E}_\lambda(q_{X_0}(1-U)q_\phi(U)) \\ &= \int_0^1 q_{X_0}(1-t)q_\phi(t) dt.\end{aligned}$$

For any $X \geq 0$, one has $\mathbb{E}^* X = \mathbb{E}(X\phi) \geq \int_0^1 q_X(t)q_\phi(1-t) dt$ by virtue of Lemma 7.11. If moreover $X \succeq_{\text{uni}} X_0$, we obtain from Lemma 7.10 (applied by choosing $h(t) = q_\phi(1-t)$), that $\mathbb{E}^* X \geq \int_0^1 q_{X_0}(t)q_\phi(1-t) dt = \mathbb{E}^* X^*$. We conclude that X^* is indeed the optimizer.

It remains to identify the function f as the Radon-Nikodym derivative $\frac{d\nu}{d\mu_\phi}$ and in terms of the quantile function q_{X_0} and the distribution function F_ϕ . This is the content of Exercise 7.10. \square

7.3 Exercises

7.1 Let X be a bounded random variable. Suppose that \mathbb{P} is not absolutely continuous w.r.t \mathbb{P}^* . Then there exists $F \in \mathcal{F}$ such that $\mathbb{P}^*(F) = 0$ and $\mathbb{P}(F) > 0$. Put $X_1 = X + c \mathbf{1}_F$. Show that X_1 'performs better than X '. Find also an example for the case where \mathbb{P}^* is not absolutely continuous w.r.t \mathbb{P} .

7.2 Let $u(x) = 1 - e^{-\alpha x}$, $\alpha > 0$ and assume that $H(\mathbb{P}^*|\mathbb{P}) < \infty$.

- Determine I and show that $\mathbb{E}^* I(c\phi) = -\frac{1}{\alpha}(\log \frac{c}{\alpha} + H(\mathbb{P}^*|\mathbb{P}))$.
- Compute X^* for problem 7.1 for a given initial capital w .
- Let \mathbb{P}^* be the probability measure of Corollary 6.14 (and let $\lambda^* = -\alpha\xi^*$). Show that in this case $X^* = \frac{\tilde{\xi}^* \cdot \bar{S}}{1+r}$, where $\tilde{\xi} = (\xi_0, \xi)$ for some ξ_0 (which one?).

7.3 This exercise concerns the case where $W = \infty$ (see Theorem 7.4). Consider the CARA utility function $u(x) = -e^{-\alpha x}$.

(a) Show that

$$I^+(y) = \left(-\frac{1}{\alpha} \log \frac{y}{\alpha}\right)^+$$

for $y \in [0, \infty]$.

(b) Show that the function $g : (0, \infty) \rightarrow (0, \infty]$ defined by $g(y) = \mathbb{E}^* I^+(y \phi)$ is decreasing and continuous on the set where it is finite and that $\lim_{y \downarrow 0} g(y) = +\infty$, $\lim_{y \rightarrow \infty} g(y) = 0$.

(c) Let \mathbb{P}^* be the risk-neutral measure of Proposition 6.7 and consider the optimization problem addressed in that proposition. Show that the optimal X^* is now of the form $X^* = (\xi^* \cdot Y - K)^+$ (a kind of European call option), where

$$K = \frac{1}{\alpha} \log \frac{c}{\alpha} + \frac{1}{\alpha} H(\mathbb{P}^* | \mathbb{P}).$$

7.4 In the setting of Theorem 7.4, let $W = \infty$ and let $u = u_{1,0}$ be a HARA utility function with index $\gamma \in [0, 1)$, see Example 4.10.

(a) Let $\gamma = 0$, $u(x) = \log x$. Show that for given $w > 0$ the optimal X^* is given by $X^* = w \frac{d\mathbb{P}}{d\mathbb{P}^*}$ and that the maximal expected utility equals $\log w + H(\mathbb{P} | \mathbb{P}^*)$ (assume that this is finite).

(b) Let $\gamma \in (0, 1)$. Compute the optimal X^* for this case.

7.5 Show that the function g in the proof of Corollary 7.5 is strictly decreasing on $g^{-1}[(0, \mathbb{E}^* W)]$.

7.6 Investigate whether the assertion of Corollary 7.5 continues to hold for the case where $W = \infty$ and $0 < w < \infty$. Impose additional assumptions (on u for instance as in Theorem 7.2), if needed.

7.7 Prove Lemma 7.8. (*Depending on the proof, it may be convenient to distinguish between $x \geq 0$ and $x < 0$. It is a good idea to interpret integrals as areas*).

7.8 Show the equivalence (i) \Leftrightarrow (ii) of Lemma 7.10.

7.9 Let X^* be the optimal random variable of Theorem 7.12. Show that $\mathbb{E} X^* = \mathbb{E} X_0$. Are the laws of X^* and X_0 the same under \mathbb{P} ? What is X^* if it happens that $\mathbb{P}^* = \mathbb{P}$? Is there an intuitive explanation for this?

7.10 Let $\nu(B) = \lambda[\mathbf{1}_B(q_\phi(U))q_{X_0}(1 - U)]$, $B \in \mathcal{B}(\mathbb{R})$ and let μ_ϕ be the distribution of ϕ .

(a) Show that $\nu \ll \mu_\phi$ and that $\nu(\mathbb{R}) = \mathbb{E} X_0$.

(b) Let f be as in (7.6). Show that (up to sets of Lebesgue measure zero) it holds that $f = \frac{d\nu}{d\mu_\phi}$.

(c) Identify f as given in Equation (7.5).

7.11 Give a concrete example where the X^* in Theorem 7.12 is different from X_0 .

7.12 Let F be a distribution function and q any of its quantile functions. Let q^- and q^+ be the extremal quantile functions and note that $q^- \leq q^+$.

- (a) Show that $\{q^- = q = q^+\}$ has Lebesgue measure one. You may use Theorem 3.10 of the MTP lecture notes.
- (b) If U is a random variable with the uniform distribution on $(0, 1)$, show that $q(U)$ has distribution function F .

8 Dynamic arbitrage theory

We return to the setting of Section 1 in the sense that we will work with a market of $d+1$ assets, of which one is often taken to be non-risky. The crucial difference though, is that we will work with *dynamic models*. That is, prices will be given by *stochastic processes* with a non-trivial time set. So, instead of working only with times $t = 0$, where all random quantities involved are deterministic (known) and a time $t = 1$, where prices of risky assets are understood as random variables, we will consider processes with a time index $t \in \{0, 1, \dots, T\}$, where T is some fixed integer greater than (or equal to) one.

We denote by S_t the d -dimensional random vector representing the nonnegative prices of the risky assets at time t . The quantities S_t^0 will be the prices of the non-risky asset at times t . Usually we take the S_t^0 *non-random* and $S_0^0 = 1$. By \bar{S}_t we denote the vector (S_t^0, S_t) . Similar notation is used for the portfolio and we have $\bar{\xi}_t = (\xi_t^0, \xi_t)$ with the obvious interpretation. The value of a portfolio at time t will be denoted by W_t , so we have $W_t = \bar{\xi}_t \cdot \bar{S}_t$.

The reader is supposed to be familiar with the notions of *filtration*, *adapted* and *predictable* processes, *martingales* and other concepts that are standard within this context.

8.1 Self-financing trading strategies

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be the probability space on which all random variables below are defined. We assume that we are given a filtration $\mathbb{F} = \{\mathcal{F}_0, \dots, \mathcal{F}_T\}$, where \mathcal{F}_0 is trivial, $\mathcal{F}_0 = \{\emptyset, \Omega\}$. Since we fix the time horizon to be T , we assume that $\mathcal{F}_T = \mathcal{F}$. The price process $S = (S_t)_{t=0}^T$ is assumed to be adapted to the filtration \mathbb{F} .

Definition 8.1 A *trading strategy* $\bar{\xi} = \{\bar{\xi}_1, \dots, \bar{\xi}_T\}$ is a $d+1$ -dimensional predictable process, i.e. for every $t > 0$, the random vector ξ_t is \mathcal{F}_{t-1} -measurable.

The interpretation of a trading strategy is that at time $t-1$ ($t \geq 1$) an investor composes a portfolio $\bar{\xi}_t$, for which (s)he then has to pay $\bar{\xi}_t \cdot \bar{S}_{t-1}$, where $t \geq 1$. This portfolio is held until time t , when the value of the portfolio changes into $W_t = \bar{\xi}_t \cdot \bar{S}_t$. At that time (s)he can re-balance the portfolio to ξ_{t+1} , for which (s)he has to pay $\bar{\xi}_{t+1} \cdot \bar{S}_t$. This re-balancing may happen without infusion or withdrawing of money and will then only be financed by the current value. The requirement of a trading strategy to be predictable is of course reasonable, an investor is not supposed to know future price movements of the stocks (s)he invests in. By definition, a predictable process is formally only defined for $t \geq 1$, but for notational convenience, we will also use $\bar{\xi}_0 := \bar{\xi}_1$.

Definition 8.2 A trading strategy is called *self-financing*, if one has

$$\bar{\xi}_t \cdot \bar{S}_t = \bar{\xi}_{t+1} \cdot \bar{S}_t,$$

for all $t \in \{0, \dots, T-1\}$.

For any stochastic process X we denote by ΔX the process with $\Delta X_t = X_t - X_{t-1}$, for $t \geq 1$ and $\Delta X_0 = X_0$.

Proposition 8.3 *A trading strategy is self-financing iff for all $t \in \{1, \dots, T\}$ one has*

$$\Delta W_t = \bar{\xi}_t \cdot \Delta \bar{S}_t.$$

Proof Exercise 8.1. □

We will take the process S^0 as a *numéraire*. For this we need and *assume* that S^0 is strictly positive (occasionally strictly positive a.s.). The *discounted* processes X^i ($i = 0, \dots, d$) are defined by

$$X_t^i = \frac{S_t^i}{S_t^0}.$$

Of course $X_t^0 = 1$ for all t . Write $X_t = (X_t^1, \dots, X_t^d)$ and $\bar{X}_t = (X_t^0, X_t)$. The (*discounted*) *value process* V is defined by

$$V_t = \frac{W_t}{S_t^0}, \quad t = 1, \dots, T$$

or, equivalently,

$$V_t = \bar{\xi}_t \cdot \bar{X}_t.$$

Note that $V_0 = W_0$. We also need the (*discounted*) *gains process* G , defined by

$$G_t = \sum_{k=1}^t \xi_k \cdot \Delta X_k.$$

We now characterize a self-financing strategy in terms of the discounted gains process.

Proposition 8.4 *Let $\bar{\xi}$ be a trading strategy. The following are equivalent.*

- (i) $\bar{\xi}$ is self-financing.
- (ii) $V_t = V_0 + G_t$, $t = 0, \dots, T$.

Proof By Definition 8.2, the strategy $\bar{\xi}$ is self-financing iff $\bar{\xi}_t \cdot \bar{X}_t = \bar{\xi}_{t+1} \cdot \bar{X}_t$ for $t = 0, \dots, T-1$, which is in turn equivalent to $\Delta V_t = \Delta G_t$, $t = 1, \dots, T$. □

We see that a strategy is self-financing iff changes in the discounted net gains process are completely due to changes in the (discounted) value process, $\Delta V_t = \xi_t \Delta X_t$.

Remark 8.5 As before, we will concentrate on the risky part ξ of the strategy $\bar{\xi}$, if $\bar{\xi}$ is self-financing. This, together with the initial investment V_0 completely determines the trading strategy. Indeed, for a self-financing strategy one has

$$\xi_{t+1}^0 = \xi_t^0 - (\xi_{t+1} - \xi_t) \cdot X_t,$$

and

$$\xi_1^0 = V_0 - \xi_1 \cdot X_0.$$

Conversely, knowing the risky part of a strategy, the above two equations yield a self-financing strategy $\bar{\xi} = (\xi^0, \xi)$.

8.2 Arbitrage

As before, the intuitive meaning of arbitrage is that it is possible to make a (positive) profit, whereas losses are impossible, also called absence of *downside risk*. The formal definition is as follows.

Definition 8.6 A self-financing trading strategy is called an *arbitrage opportunity* if its value process V satisfies $V_0 \leq 0$, $\mathbb{P}(V_T \geq 0) = 1$ and $\mathbb{P}(V_T > 0) > 0$. A market is called arbitrage free, if no arbitrage opportunities exist.

As in Section 1, absence of arbitrage in the market is necessary to obtain a fair and sensible pricing system. We first give a characterization of existence of arbitrage. Later on we alternatively characterize absence of arbitrage.

Proposition 8.7 *An arbitrage opportunity exists iff there is a $t \in \{1, \dots, T\}$ and a \mathcal{F}_{t-1} -measurable random vector η_t such that $\mathbb{P}(\eta_t \cdot \Delta X_t \geq 0) = 1$ and $\mathbb{P}(\eta_t \cdot \Delta X_t > 0) > 0$.*

Proof Let $\bar{\xi}$ be an arbitrage opportunity and V the corresponding discounted value process. Put

$$t = \min\{k \geq 1 : \mathbb{P}(V_k \geq 0) = 1 \text{ and } \mathbb{P}(V_k > 0) > 0\}.$$

Then $t \leq T$ and $\mathbb{P}(V_{t-1} \geq 0) < 1$ or $\mathbb{P}(V_{t-1} > 0) = 0$. In the first case, let $\eta_t = \xi_t \mathbf{1}_{\{V_{t-1} < 0\}}$. Then η_t is \mathcal{F}_{t-1} -measurable and

$$\eta_t \cdot \Delta X_t = \Delta V_t \mathbf{1}_{\{V_{t-1} < 0\}} \geq -V_{t-1} \mathbf{1}_{\{V_{t-1} < 0\}},$$

and the requirements are met. In the other case, we take $\eta_t = \xi_t$ and then $\xi_t \cdot \Delta X_t = \Delta V_t \geq V_t$ a.s. and again the requirements are met, by definition of t .

Conversely, assume that η_t with the stipulated properties exists. Define the trading strategy ξ by $\xi_s = \eta_t \mathbf{1}_{\{t\}}(s)$ and complete it by choosing $V_0 = 0$ and ξ^0 as in Remark 8.5 such that $\bar{\xi}$ is self-financing. Then $V_T = \eta_t \cdot \Delta X_t$ and we have an arbitrage property. \square

We have seen in Section 1 that absence of arbitrage was equivalent with the existence of a risk-neutral measure \mathbb{P}^* , that by definition had the property, using the current notation, that $\mathbb{E}^* X_1 = X_0$, which is in fact the martingale property of the pair (X_0, X_1) , since \mathcal{F}_0 is trivial. This makes the next definition understandable.

Definition 8.8 A probability measure \mathbb{Q} on (Ω, \mathcal{F}_T) is called a *martingale measure* or a *risk-neutral measure* if the process X is a martingale under \mathbb{Q} . If a martingale measure \mathbb{P}^* is equivalent to \mathbb{P} on \mathcal{F}_T , then it is called an *equivalent martingale measure*. The set of all equivalent martingale measures is denoted by \mathcal{P} .

There are various ways to characterize martingale measures. We use the following

Theorem 8.9 Let \mathbb{Q} be a probability measure on (Ω, \mathcal{F}_T) . Equivalent are

- (i) \mathbb{Q} is a martingale measure.
- (ii) If $\bar{\xi}$ is self-financing and bounded, then its value process V is a \mathbb{Q} -martingale.
- (iii) If $\bar{\xi}$ is self-financing and $\mathbb{E}_{\mathbb{Q}} V_T^- < \infty$, then V is a \mathbb{Q} -martingale.
- (iv) If $\bar{\xi}$ is self-financing and $\mathbb{Q}(V_T \geq 0) = 1$, then $\mathbb{E}_{\mathbb{Q}} V_T = V_0$.

Proof (i) \Rightarrow (ii): It follows that V_t is \mathbb{Q} -integrable for each t , since ξ is bounded. From Proposition 8.4 and ξ being predictable, we have for $t \geq 1$

$$\begin{aligned}\mathbb{E}_{\mathbb{Q}} [\Delta V_t | \mathcal{F}_{t-1}] &= \xi_t \cdot \mathbb{E}_{\mathbb{Q}} [\Delta X_t | \mathcal{F}_{t-1}] \\ &= 0,\end{aligned}$$

since X is a \mathbb{Q} -martingale.

(ii) \Rightarrow (iii): As a first step in the proof, we show

$$(8.1) \quad \mathbb{E}_{\mathbb{Q}} V_t^- < \infty \Rightarrow \mathbb{E}_{\mathbb{Q}} [V_t | \mathcal{F}_{t-1}] = V_{t-1} \mathbb{Q}\text{-a.s.}$$

Since $\mathbb{E}_{\mathbb{Q}} V_t^- < \infty$, the (generalized) conditional expectation $\mathbb{E}_{\mathbb{Q}} [V_t | \mathcal{F}_{t-1}]$ is well defined. Fix $a > 0$ and put $\xi_t^a = \xi_t \mathbf{1}_{\{|\xi_t| \leq a\}}$. Then $\xi_t^a \cdot \Delta X_t$ is the increment of a martingale, since ξ_t^a is bounded, so $\mathbb{E}_{\mathbb{Q}} [\xi_t^a \cdot \Delta X_t | \mathcal{F}_{t-1}] = 0$. Hence

$$\begin{aligned}\mathbb{E}_{\mathbb{Q}} [V_t | \mathcal{F}_{t-1}] \mathbf{1}_{\{|\xi_t| \leq a\}} &= \mathbb{E}_{\mathbb{Q}} [V_t \mathbf{1}_{\{|\xi_t| \leq a\}} | \mathcal{F}_{t-1}] \\ &= \mathbb{E}_{\mathbb{Q}} [V_t \mathbf{1}_{\{|\xi_t| \leq a\}} | \mathcal{F}_{t-1}] - \mathbb{E}_{\mathbb{Q}} [\xi_t^a \cdot \Delta X_t | \mathcal{F}_{t-1}] \\ &= \mathbb{E}_{\mathbb{Q}} [(V_{t-1} + \xi_t \cdot \Delta X_t) \mathbf{1}_{\{|\xi_t| \leq a\}} - \xi_t^a \cdot \Delta X_t | \mathcal{F}_{t-1}] \\ &= V_{t-1} \mathbf{1}_{\{|\xi_t| \leq a\}}.\end{aligned}$$

By letting $a \rightarrow \infty$, one obtains (8.1). Use this equation for $t = T$ to get

$$\mathbb{E}_{\mathbb{Q}} V_{T-1}^- = \mathbb{E}_{\mathbb{Q}} (\mathbb{E}_{\mathbb{Q}} [V_T | \mathcal{F}_{T-1}])^- \leq \mathbb{E}_{\mathbb{Q}} V_T^- ,$$

by Jensen's inequality for $x \mapsto x^-$. From the assumption, we get that $\mathbb{E}_{\mathbb{Q}} V_{T-1}^- < \infty$. Iterating this procedure, one obtains $\mathbb{E}_{\mathbb{Q}} V_t^- < \infty$ for all t and by (8.1) also $\mathbb{E}_{\mathbb{Q}} V_t = \mathbb{E}_{\mathbb{Q}} V_0 = V_0 < \infty$. It follows that all V_t are integrable and combined with (8.1) this makes that V is a martingale.

(iii) \Rightarrow (iv): We clearly have $\mathbb{E}_{\mathbb{Q}} V_T^- = 0$ and by the fact that V is then a martingale, $\mathbb{E}_{\mathbb{Q}} V_T = \mathbb{E}_{\mathbb{Q}} V_0 = V_0$.

(iv) \Rightarrow (i): We have to show that X is a \mathbb{Q} -martingale. First we show that every element X_t^i of X_t is \mathbb{Q} -integrable, if $t \leq T$ by selecting an appropriate trading strategy. Let $\xi_s^i = \mathbf{1}_{\{s \leq t\}}$ and $\xi^j = 0$ if $1 \leq j \neq i$. Let $V_0 = X_0^i$ and choose ξ^0 such that $\bar{\xi}$ is self-financing. It follows that now $V_T = X_T^i \geq 0$. Using the assumption, we get that X_t^i has finite expectation, in fact

$$(8.2) \quad \mathbb{E}_{\mathbb{Q}} X_t^i = X_0^i.$$

Next we show that $\mathbb{E}_{\mathbb{Q}} [\Delta X_t^i | \mathcal{F}_{t-1}] = 0$, \mathbb{Q} -a.s., equivalently, $\mathbb{E}_{\mathbb{Q}} [\mathbf{1}_A \Delta X_t^i] = 0$ for every $A \in \mathcal{F}_{t-1}$, by selecting another appropriate trading strategy. Given such

A , we define $\xi_s^i = \mathbf{1}_{s \leq t} - \mathbf{1}_A \mathbf{1}_{\{s=t\}}$ and $\xi_s^j = 0$ if $1 \leq j \neq i$. Let $V_0 = X_0^i \geq 0$ and complement ξ to obtain a self-financing strategy (note that it is indeed predictable). A simple computation gives

$$V_T = X_T^i - \mathbf{1}_A \Delta X_t^i = \mathbf{1}_{A^c} X_t^i + \mathbf{1}_A X_{t-1}^i \geq 0.$$

The assumption $\mathbb{E}_{\mathbb{Q}} V_T = V_0$ now reads $\mathbb{E}_{\mathbb{Q}} (X_t^i - \mathbf{1}_A \Delta X_t^i) = \mathbb{E}_{\mathbb{Q}} X_t^i$, in view of (8.2). It follows that $\mathbb{E}_{\mathbb{Q}} [\mathbf{1}_A \Delta X_t^i] = 0$. \square

Remark 8.10 Suppose that \mathbb{P} itself is a martingale measure and that a risk averse investor uses the same probability measure to decide whether or not to invest in products with a certain expected pay-off. According to Example 4.7, he will invest all his capital in a riskless product. Moreover, the market is arbitrage free. Indeed, if $\bar{\xi}$ is a self-financing strategy and $V_0 \leq 0$, then we obtain that $V_T \geq 0$ \mathbb{P} -a.s. implies that $\mathbb{E} V_T = 0$ and hence $V_T = 0$ \mathbb{P} -a.s.

The main theorem of this section is Theorem 8.12 below, the dynamic version of Theorem 1.6. To prove it, we need a lemma that concerns a static one period model as in Section 1, but now with *random* initial prices. We single out one time step of the dynamic model, the one from $t-1$ to t . Below we write $L^0(\Omega, \mathcal{G}, \mathbb{P})$ for the set of \mathcal{G} -measurable random variables, with the identification of \mathbb{P} -a.s. equal random variables to be the same.

The *space* generated by the discounted net gains from $t-1$ to t with $t \in \{1, \dots, T\}$ is

$$(8.3) \quad \mathcal{K}_t = \{\xi \cdot \Delta X_t : \xi^i \in L^0(\Omega, \mathcal{F}_{t-1}, \mathbb{P}), i = 1, \dots, d\}.$$

Hence, by Proposition 8.7 an arbitrage-free market can be characterized by the relation

$$\mathcal{K}_t \cap L_+^0(\Omega, \mathcal{F}_t, \mathbb{P}) = \{0\}, \forall t \in \{1, \dots, T\}.$$

The following is of *fundamental* importance, a cornerstone in the proof of Theorem 8.12.

Lemma 8.11 *Let $t \in \{1, \dots, T\}$. The following statements are equivalent.*

- (i) *The intersection $\mathcal{K}_t \cap L_+^0(\Omega, \mathcal{F}_t, \mathbb{P}) = \{0\}$.*
- (ii) *There exists a probability measure \mathbb{P}_t^* on \mathcal{F}_t , equivalent to \mathbb{P} , with a bounded \mathcal{F}_t -measurable density $Z_t = \frac{d\mathbb{P}_t^*}{d\mathbb{P}}$, such that $\mathbb{E}_{\mathbb{P}_t^*}[\Delta X_t | \mathcal{F}_{t-1}] = 0$. The measure \mathbb{P}_t^* can be trivially extended to a probability measure on \mathcal{F} by putting $\mathbb{P}_t^*(F) = \mathbb{E}_{\mathbb{P}} \mathbf{1}_F Z_t$ for all $F \in \mathcal{F}$.*

Proof See Section 8.3, where this lemma is alternatively formulated as Corollary 8.24. \square

We return to the dynamic setting. The next theorem is the first *Fundamental Theorem of Asset Pricing* for a dynamic market in discrete time.

Theorem 8.12 *The market is free of arbitrage iff there exists a risk-neutral measure \mathbb{P}^* on \mathcal{F}_T with bounded Radon-Nikodym derivative $\frac{d\mathbb{P}^*}{d\mathbb{P}}$.*

Proof Assume that a risk-neutral measure \mathbb{P}^* exists. Let $\bar{\xi}$ be any self-financing trading strategy with $V_0 \leq 0$ and $\mathbb{P}^*(V_T \geq 0) = 1$. Theorem 8.9 yields $0 \leq \mathbb{E}^* V_T = V_0 \leq 0$, hence $\mathbb{P}^*(V_T = 0) = 1$ and an arbitrage opportunity doesn't exist under \mathbb{P}^* , and then also not under \mathbb{P} by equivalence of the two measures.

Conversely, assume that the market is free of arbitrage. Let $t \in \{1, \dots, T\}$ and let \mathcal{K}_t be as in (8.3). Recall that by Proposition 8.7 it holds that $\mathcal{K}_t \cap L_+^0(\Omega, \mathcal{F}_t, \mathbb{P}) = \{0\}$ for all t . Consider $t = T$, then Lemma 8.11 applies with $t = T$ and we conclude to the existence of a probability measure \mathbb{P}_T^* on $\mathcal{F}_T = \mathcal{F}$, with $\mathbb{P}_T^* \sim \mathbb{P}$ and $\mathbb{E}_{\mathbb{P}_T^*}[\Delta X_T | \mathcal{F}_{T-1}] = 0$. Moreover $Z_T = \frac{d\mathbb{P}_T^*}{d\mathbb{P}}$ is bounded.

We proceed by *backward* induction. Suppose that for $t < T$ a probability measure \mathbb{P}_{t+1}^* on \mathcal{F} is found such that $\mathbb{P}_{t+1}^* \sim \mathbb{P}$, with bounded density, and

$$(8.4) \quad \mathbb{E}_{\mathbb{P}_{t+1}^*}[\Delta X_k | \mathcal{F}_{k-1}] = 0, \text{ for } t+1 \leq k \leq T,$$

in other words, the process $(X_k)_{k \in \{t, \dots, T\}}$ is a martingale under \mathbb{P}_{t+1}^* . By equivalence we also have $\mathcal{K}_t \cap L_+^0(\Omega, \mathcal{F}_t, \mathbb{P}_{t+1}^*) = \{0\}$. Then, we apply Lemma 8.11 again to obtain existence of a probability measure \mathbb{P}_t^* on \mathcal{F} , equivalent to \mathbb{P}_{t+1}^* , with bounded density $Z_t = \frac{d\mathbb{P}_t^*}{d\mathbb{P}_{t+1}^*}$ which is \mathcal{F}_t -measurable, and such that

$$\mathbb{E}_{\mathbb{P}_t^*}[\Delta X_t | \mathcal{F}_{t-1}] = 0.$$

Our next aim is to show for $t+1 \leq k \leq T$ the equality $\mathbb{E}_{\mathbb{P}_t^*}[\Delta X_k | \mathcal{F}_{k-1}] = 0$, equivalently $\mathbb{E}_{\mathbb{P}_t^*}[\mathbf{1}_A \Delta X_k] = 0$, for $A \in \mathcal{F}_{k-1}$. Take such an A and compute, using \mathcal{F}_t -measurability of Z_t and (8.4),

$$\begin{aligned} \mathbb{E}_{\mathbb{P}_t^*}[\mathbf{1}_A \Delta X_k] &= \mathbb{E}_{\mathbb{P}_{t+1}^*}[\mathbf{1}_A \Delta X_k Z_t] \\ &= \mathbb{E}_{\mathbb{P}_{t+1}^*} \mathbb{E}_{\mathbb{P}_{t+1}^*}[\mathbf{1}_A \Delta X_k Z_t | \mathcal{F}_{k-1}] \\ &= \mathbb{E}_{\mathbb{P}_{t+1}^*} (Z_t \mathbf{1}_A \mathbb{E}_{\mathbb{P}_{t+1}^*}[\Delta X_k | \mathcal{F}_{k-1}]) \\ &= 0. \end{aligned}$$

Hence Equation (8.4) remains true with the substitution $t+1 \rightarrow t$. Moreover,

$$\frac{d\mathbb{P}_t^*}{d\mathbb{P}} = Z_t \frac{d\mathbb{P}_{t+1}^*}{d\mathbb{P}}$$

is bounded as well. By iteration, we conclude that the procedure yields a probability measure $\mathbb{P}^* = \mathbb{P}_1^*$ with the desired properties. \square

We close this section by studying what happens under a *change of numéraire*. Absence of arbitrage is defined as the impossibility to have a \mathbb{P} -almost sure profit. Clearly, we can replace in this statement \mathbb{P} with any risk-neutral measure \mathbb{P}^* , since these measures define the same null sets and the role of the process S^0 is not relevant to describe arbitrage. But any particular \mathbb{P}^* is such that the price processes discounted by the numéraire process S^0 are, by definition, \mathbb{P}^* -martingales. Hence, if one prefers to take the price process of another asset as a discount factor, there will be another risk-neutral measure. So, the set

of risk-neutral measures depends on the choice of numéraire and it is interesting to investigate how different risk-neutral measures resulting from different numéraires are related.

Suppose one takes the process S^1 as a numéraire. It is assumed that S^1 is \mathbb{P} -a.s. strictly positive. Put

$$\bar{Y}_t = \frac{\bar{S}_t}{S_t^1},$$

then $\bar{Y}_t X_t^1 = \bar{X}_t$, $t \in \{0, \dots, T\}$. Let $\tilde{\mathcal{P}}$ denote the set of all probability measures $\tilde{\mathbb{P}}$ equivalent to \mathbb{P} that are such that \bar{Y} is a $\tilde{\mathbb{P}}$ -martingale. Absence of arbitrage is then equivalent to $\tilde{\mathcal{P}} \neq \emptyset$, by virtue of Theorem 8.12.

Proposition 8.13 *A probability measure $\tilde{\mathbb{P}}$ belongs to $\tilde{\mathcal{P}}$ iff there exists a probability measure $\mathbb{P}^* \in \mathcal{P}$ such that $\tilde{\mathbb{P}} \sim \mathbb{P}^*$ and*

$$(8.5) \quad \frac{d\tilde{\mathbb{P}}}{d\mathbb{P}^*} = \frac{X_T^1}{X_0^1}.$$

In this case one also has

$$\frac{d\mathbb{P}^*}{d\tilde{\mathbb{P}}} = \frac{Y_T^0}{Y_0^0}.$$

Proof Let \mathbb{P}^* be given. The random variables $\frac{X_t^1}{X_0^1}$ form a martingale under \mathbb{P}^* , with $\mathbb{E}^* \frac{X_T^1}{X_0^1} = 1$. Hence, if we define $\tilde{\mathbb{P}}$ by (8.5), then it is a probability measure, equivalent to \mathbb{P}^* and for $t > s$ one has

$$\begin{aligned} \mathbb{E}_{\tilde{\mathbb{P}}}[\bar{Y}_t | \mathcal{F}_s] &= \frac{\mathbb{E}^*[\bar{Y}_t \frac{X_t^1}{X_0^1} | \mathcal{F}_s]}{\mathbb{E}^*[\frac{X_t^1}{X_0^1} | \mathcal{F}_s]} \\ &= \frac{\mathbb{E}^*[\bar{Y}_t X_t^1 | \mathcal{F}_s]}{X_s^1} \\ &= \frac{\mathbb{E}^*[\bar{X}_t | \mathcal{F}_s]}{X_s^1} \\ &= \frac{\bar{X}_s}{X_s^1} \\ &= \bar{Y}_s. \end{aligned}$$

Hence $\tilde{\mathbb{P}}$ is a martingale measure for \bar{Y} , or $\tilde{\mathbb{P}} \in \tilde{\mathcal{P}}$. To prove the other implication, one just swaps the roles of X and Y in the previous part. \square

Proposition 8.14 *Suppose that X_T^1 is not degenerate under \mathbb{P} . Then the sets \mathcal{P} and $\tilde{\mathcal{P}}$ have empty intersection.*

Proof Exercise 8.2. \square

8.3 Proof of Lemma 8.11

This section extends the proof of existence of an equivalent martingale measure (FTAP, Theorem 1.6) to the situation of a non-trivial initial history, \mathcal{F}_0 is not necessarily the trivial σ -algebra. The background is a multi-period model as in Section 8 in which *we single out one arbitrary time step*, from $t-1$ to t , for some $t \in \{1, \dots, T\}$. The prices S_t^i are \mathcal{F}_t -measurable nonnegative random variables and the portfolio choices ξ_t^i are \mathcal{F}_{t-1} -measurable. Note that \mathcal{F}_{t-1} is in general not a trivial σ -algebra. By a time shift, we may as well consider a one period model with $t = 0, 1$ as in Section 1, but with the generalization that \mathcal{F}_0 is no longer trivial. Having done so we can extend the results below to an arbitrary step in a multi-period setting, which eventually leads to Theorem 8.12.

Here is some notation. We use L^p as an abbreviation of $L^p(\Omega, \mathcal{F}_1, \mathbb{P})$, for $p \geq 0$. For $p = 0$, we make L^0 a metric space by using the metric d defined by $d(X, Y) = \mathbb{E}|X - Y| \wedge 1$. It then holds that $d(X_n, X) \rightarrow 0$ iff $X_n \xrightarrow{\mathbb{P}} X$. By L_+^p we denote the nonnegative elements of L^p .

We adopt the following standing assumption throughout this section. *The $(d + 1)$ -dimensional price process S is assumed to be adapted and such that the discounted prices $X_t^i = S_t^i/S_t^0$ have finite expectation for all $1 \leq i \leq d$ and $t = 0, 1$. The integrability assumption can be circumvented more or less as in Exercise 1.3. Furthermore, we require (non-random) $S_0^0 > 0$ and $S_1^0 > 0$ to have the X_t^i well defined.*

A portfolio is a $(d + 1)$ -dimensional random vector that is \mathcal{F}_0 -measurable and thus not necessarily constant. As usual we denote by ξ the investments in the risky assets, now a d -dimensional \mathcal{F}_0 -measurable random vector. The vector of net gains Y is also defined as usual, but adapted to the current situation we have

$$Y = X_1 - X_0.$$

The random vector Y is \mathcal{F}_1 -measurable and (component wise) integrable under the standing assumption. A market is arbitrage free if $\xi \cdot Y \geq 0$ a.s. implies $\xi \cdot Y = 0$ a.s. The characterization of an arbitrage free market now becomes $\mathcal{K} \cap L_+^0 = \{0\}$, where $\mathcal{K} = \{\xi \cdot Y : \xi^i \in \mathcal{F}_0, i = 1, \dots, d\}$. The concept of martingale measure, adapted to the present situation, is as follows.

Definition 8.15 A probability measure \mathbb{Q} on (Ω, \mathcal{F}_1) is called a martingale measure, or risk-neutral measure, if $\mathbb{E}_{\mathbb{Q}}[Y|\mathcal{F}_0] = 0$ \mathbb{Q} -a.s. It is called *equivalent martingale measure*, if moreover $\mathbb{Q} \sim \mathbb{P}$. The set of equivalent martingale measures is denoted \mathcal{P} .

Lemma 8.16 *There is equivalence between $\mathcal{K} \cap L_+^0 = \{0\}$ and $(\mathcal{K} - L_+^0) \cap L_+^0 = \{0\}$.*

Proof Assume $\mathcal{K} \cap L_+^0 = \{0\}$ and let $Z \in \mathcal{K} - L_+^0$, $Z = \xi \cdot Y - U$ say. If $Z \geq 0$, then also $\xi \cdot Y \geq 0$ and by the hypothesis $\xi \cdot Y = 0$, which yields $Z = -U \leq 0$. So $Z = 0$. The converse implication follows from $\mathcal{K} \subset \mathcal{K} - L_+^0$. \square

In all what follows we let

$$\mathcal{C} = (\mathcal{K} - L_+^0) \cap L^1.$$

Note that \mathcal{C} is a cone, $W \in \mathcal{C}$ implies $\lambda W \in \mathcal{C}$ for all $\lambda \geq 0$. In the proof of the next lemma we need the formula for conditional expectations under an absolutely continuous change of measure. Let $\mathbb{Q} \ll \mathbb{P}$ with Radon-Nikodym derivative Z . If $\mathbb{E}|XZ| < \infty$ and \mathcal{G} a sub- σ -algebra of \mathcal{F} , then

$$(8.6) \quad \mathbb{E}_{\mathbb{Q}}[X|\mathcal{G}] = \frac{\mathbb{E}[XZ|\mathcal{G}]}{\mathbb{E}[Z|\mathcal{G}]} \quad \mathbb{Q}\text{-a.s.}$$

Lemma 8.17 *Suppose there is $Z \in L^\infty$ such that $\mathbb{E}(ZW) \leq 0$ for all $W \in \mathcal{C}$. Then $Z \geq 0$ a.s. and if $\mathbb{E}Z = 1$, then $d\mathbb{Q} = Z d\mathbb{P}$ defines a martingale measure \mathbb{Q} .*

Proof Note that $W = -\mathbf{1}_{\{Z < 0\}} \in \mathcal{C}$. Hence $\mathbb{E}(-Z\mathbf{1}_{\{Z < 0\}}) \leq 0$ and it follows that $Z\mathbf{1}_{\{Z < 0\}} = 0$ a.s., hence $Z \geq 0$ a.s. Under the condition $\mathbb{E}Z = 1$, \mathbb{Q} is a probability measure, absolutely continuous w.r.t. \mathbb{P} .

Let ξ be bounded, \mathcal{F}_0 -measurable and $\lambda \in \{-1, 1\}$. Since $\xi \cdot Y \in \mathcal{K}$, also $\lambda\xi \cdot Y \in \mathcal{K} \subset \mathcal{K} - L_+^0$. Because ξ is bounded, we also have $(\lambda\xi) \cdot Y \in L^1$, hence $(\lambda\xi) \cdot Y \in \mathcal{C}$ and therefore $\lambda\mathbb{E}(\xi \cdot Y)Z \leq 0$. But since $\lambda \in \{-1, 1\}$ is arbitrary, we must have $\mathbb{E}(\xi \cdot Y)Z = 0$. One then has $0 = \mathbb{E}(\xi \cdot Y)Z = \mathbb{E}[\mathbb{E}[(\xi \cdot Y)Z|\mathcal{F}_0]] = \mathbb{E}[\xi \cdot \mathbb{E}[YZ|\mathcal{F}_0]]$ for all bounded ξ . But then $\mathbb{E}[YZ|\mathcal{F}_0] = 0$ a.s. Equation (8.6) yields

$$\mathbb{E}_{\mathbb{Q}}[Y|\mathcal{F}_0] = \frac{\mathbb{E}[YZ|\mathcal{F}_0]}{\mathbb{E}[Z|\mathcal{F}_0]} = 0,$$

whence \mathbb{Q} is a martingale measure. □

Remark 8.18 The use of Equation (8.6) to prove that \mathbb{Q} is a martingale measure can be circumvented by computing for every $F \in \mathcal{F}_0$

$$\mathbb{E}_{\mathbb{Q}}[Y\mathbf{1}_F] = \mathbb{E}[Y\mathbf{1}_F Z] = \mathbb{E}[\mathbf{1}_F \mathbb{E}[YZ|\mathcal{F}_0]] = 0.$$

Let

$$(8.7) \quad \mathcal{Z} = \{Z \in \mathcal{F}_1 : 0 \leq Z \leq 1, \mathbb{E}Z > 0 \text{ and } \mathbb{E}(ZW) \leq 0, \forall W \in \mathcal{C}\}.$$

If the set \mathcal{Z} is nonempty, the normalization $\zeta = Z/\mathbb{E}Z$ can then serve as a Radon-Nikodym derivative of a martingale measure w.r.t. \mathbb{P} . We shall see that under the additional condition that the market is arbitrage free, the set \mathcal{Z} is indeed non-empty and one can even select a Z^* from it satisfying $\mathbb{P}(Z^* > 0) = 1$, which yields the existence of an *equivalent* martingale measure. The technical property that we need is that the set \mathcal{C} is *closed in L^1* , which will be proved first. This is quite some work.

To accomplish this we need two technical results, a decomposition of L^0 into suitable ‘orthogonal’ subspaces and a version of the Bolzano-Weierstraß theorem for sequences of random variables. Note that a random variable X can be viewed

as a collection of real numbers $X(\omega)$ for $\omega \in \Omega$ and is thus in general an infinite dimensional object. So a straightforward application of the classical Bolzano-Weierstraß theorem for sequences in a finite-dimensional Euclidean space is not possible. Here we go.

Lemma 8.19 *Let (ξ_n) be a sequence of d -dimensional random vectors defined on some $(\Omega, \mathcal{F}, \mathbb{P})$ such that $\liminf |\xi_n| < \infty$ a.s. Then exists a sequence of strictly increasing random variables σ_m and an a.s. finite random vector ξ such that $\xi_{\sigma_m} \xrightarrow{\text{a.s.}} \xi$.*

Proof Let $L = \liminf |\xi_n|$. Then $\mathbb{P}(L = \infty) = 0$ and for definiteness we define $\sigma_m = m$ on $\{L = \infty\}$. From now on we work on the set $F = \{L < \infty\} \in \mathcal{F}$. Put $\sigma_1^0 = 1$ and define recursively for $m \geq 1$

$$\sigma_{m+1}^0(\omega) = \inf A_m(\omega),$$

where

$$A_m(\omega) = \{n > \sigma_m^0(\omega) : ||\xi_n(\omega)| - L(\omega)| < \frac{1}{m}\}.$$

Note that $A_m(\omega)$ contains infinite many elements for every m and $\omega \in F$ by the hypothesis and that all σ_m^0 are \mathcal{F} -measurable. It follows that also the $\xi_{\sigma_m^0}$ are \mathcal{F} -measurable. The details are left as Exercise 8.11. Write ξ_n^1 for the first element of ξ_n and define $\xi^1 = \liminf \xi_{\sigma_m^0}^1$. Since the $\xi_{\sigma_m^0}^1(\omega)$ converge along a subsequence it makes sense to define $\sigma_1^1 = 1$ and recursively

$$\sigma_{m+1}^1(\omega) = \inf \{\sigma_n^0(\omega) > \sigma_m^1(\omega) : |\xi_{\sigma_n^0(\omega)}^1(\omega) - \xi^1(\omega)| < \frac{1}{m}\}.$$

We conclude that $\xi_{\sigma_m^1}^1 \rightarrow \xi^1$ on F , which is the desired behavior for the first component of the ξ_n . The further idea is to thin the sequence of σ_m^1 in order to obtain a subsequence for which also the second components converge. Thereto one first defines the candidate limit $\xi^2 = \liminf \xi_{\sigma_m^1}^2$ and finds a sequence (σ_m^2) by mimicking the above procedure. Go on like this with subsequent thinning until also the last component converges. \square

We proceed with the announced ‘orthogonal’ decomposition of L^0 . Recall the one-period setting, in particular ξ and η below are always d -dimensional \mathcal{F}_0 -measurable random vectors.

Lemma 8.20 *Let $N = \{\eta \in L^0(\Omega, \mathcal{F}_0, \mathbb{P})^d : \eta \cdot Y = 0 \text{ a.s.}\}$ and $N^\perp = \{\xi \in L^0(\Omega, \mathcal{F}_0, \mathbb{P})^d : \xi \cdot \eta = 0 \text{ a.s., } \forall \eta \in N\}$. Then N and N^\perp are closed subsets of $L^0(\Omega, \mathcal{F}_0, \mathbb{P})^d$, $N \cap N^\perp = \{0\}$ and $L^0(\Omega, \mathcal{F}_0, \mathbb{P})^d = N + N^\perp$.*

Proof Let $(\eta_n) \subset N$ such that $\eta_n \xrightarrow{\mathbb{P}} \eta$. Since almost sure convergence holds along a subsequence we must also have $\eta \cdot Y = 0$ a.s. Closedness of N^\perp is proved similarly. If $\eta \in N \cap N^\perp$, then $\eta \cdot \eta = 0$ a.s. and hence $\eta = 0$ a.s.

The final assertion we first prove for the non-random standard basis vectors e_i of \mathbb{R}^d by a projection argument. Note that every e_i belongs to the Hilbert

space $H = L^2(\Omega, \mathcal{F}_0, \mathbb{P})^d$. Moreover, $N \cap H$ and $N^\perp \cap H$ are both closed subspaces of H (why?) and have trivial intersection. By using the orthogonal projections on these subspaces we should have $e_i = \eta_i + \xi_i$, with $\eta_i \in N \cap H$ and $\xi_i \in N^\perp \cap H$. Note that this is not immediately guaranteed, since we don't know yet that $(N^\perp \cap H) + (N^\perp \cap H) = H$. We proceed as follows. Let η_i be the orthogonal projection of e_i onto $N \cap H$ and define $\xi_i = e_i - \eta_i$, the projection error, which is orthogonal to N by construction and has $\mathbb{E}|\xi_i|^2 < \infty$. Suppose that $\xi_i \notin N^\perp$. Then there must be $\eta \in N$ such that $\xi_i \cdot \eta \neq 0$ a.s., say $\mathbb{P}(\xi_i \cdot \eta > 0) > 0$. The truncated random vector $\tilde{\eta} := \eta \mathbf{1}_{\{\xi_i \cdot \eta > 0, |\eta| \leq c\}}$ also belongs to N , as well as to H for every $c > 0$. Now $\tilde{\eta} \cdot \xi_i = \eta \cdot \xi_i \mathbf{1}_{\{\xi_i \cdot \eta > 0, |\eta| \leq c\}}$ is positive with positive probability for c large enough and it follows that $\mathbb{E}(\tilde{\eta} \cdot \xi_i) > 0$ contradicting that ξ_i is orthogonal to N .

Having established the decomposition for the basis vectors e_i , we now turn to the general case. Every \mathcal{F}_0 -measurable random vector V can be written as $V = \sum_{i=1}^d V_i e_i$, with \mathcal{F}_0 -measurable random variables V_i . Since $e_i = \xi_i + \eta_i$ with $\eta_i \in N$ and $\xi_i \in N^\perp$, we have $V = \sum_{i=1}^d V_i \xi_i + \sum_{i=1}^d V_i \eta_i$. One verifies that along with the $\eta_i \in N$ also the $V_i \eta_i \in N$, since the V_i are \mathcal{F}_0 -measurable. Likewise the $V_i \xi_i$ belong to N^\perp . Since both spaces N and N^\perp are closed under addition, we have established a decomposition of V . Uniqueness follows from $N \cap N^\perp = \{0\}$. \square

Having done all these preparations, we can show the closedness property of \mathcal{C} .

Proposition 8.21 *Under the no arbitrage condition $\mathcal{K} \cap L_+^0 = \{0\}$ it holds that $\mathcal{K} - L_+^0$ is closed in L^0 and hence \mathcal{C} is closed in L^1 .*

Proof It is sufficient to show the first assertion, the latter being its direct consequence (why?). Let (W_n) be a sequence in $\mathcal{K} - L_+^0$ with W as its limit in probability. Along a subsequence (again denoted (W_n)) we have a.s. convergence to W . Since $W_n \in \mathcal{K} - L_+^0$, we can write $W_n = \xi_n \cdot Y - U_n$, with $U_n \geq 0$ a.s. Moreover, we may even assume $\xi_n \in N^\perp$. Indeed, by virtue of Lemma 8.20, every \mathcal{F}_0 -measurable ξ_n can be decomposed as $\xi_n = \xi'_n + \eta_n$ with $\xi'_n \in N^\perp$ and $\eta_n \in N$. But then $\xi_n \cdot Y = \xi'_n \cdot Y$.

In order to apply Lemma 8.19, we first show that $\liminf |\xi_n| < \infty$ a.s. Consider the $\zeta_n := \xi_n / |\xi_n|$ (well defined if $|\xi_n| > 0$, which is w.l.o.g. true on the set I below), these form a bounded sequence. Invoking Lemma 8.19, we can choose an increasing sequence of \mathcal{F}_0 -measurable random integers τ_n such that $\zeta_{\tau_n} \xrightarrow{\text{a.s.}} \zeta$ for some \mathcal{F}_0 -measurable random vector ζ with norm one. Since the W_n converge a.s. to a finite limit, we have *on the set* $I = \{\liminf |\xi_n| = \infty\}$

$$0 \leq \frac{U_{\tau_n}}{\xi_{\tau_n}} = \zeta_{\tau_n} \cdot Y - \frac{W_{\tau_n}}{|\xi_{\tau_n}|} \xrightarrow{\text{a.s.}} \zeta \cdot Y.$$

Since $\mathcal{K} \cap L_+^0 = \{0\}$, we conclude that $\zeta \cdot Y = 0$ a.s. on $\{\liminf |\xi_n| = \infty\}$, so $\mathbf{1}_I \zeta \cdot Y = 0$ a.s. Furthermore, since the $\xi_n \in N^\perp$, we have for every $\eta \in N$ that also $\zeta_{\tau_n} \cdot \eta = 0$ a.s. Because N^\perp is closed under a.s. convergence, it follows that $\zeta \in N^\perp$, but then also $\mathbf{1}_I \zeta \in N^\perp$, because $\mathbf{1}_I$ is \mathcal{F}_0 -measurable. Together with

the previously established fact $\mathbf{1}_I \zeta \cdot Y = 0$ (so $\mathbf{1}_I \zeta \in N$), we conclude $\mathbf{1}_I \zeta = 0$ a.s. Since $|\zeta| = 1$ a.s., this can only happen if $\mathbb{P}(I) = 0$.

Having established $\liminf |\xi_n| < \infty$ a.s., we invoke Lemma 8.19 again to obtain an a.s. finite random vector ξ and a sequence of strictly increasing \mathcal{F}_0 -measurable integer valued random variables σ_n such that $\xi_{\sigma_n} \rightarrow \xi$ a.s. Note (verify) that also $W_{\sigma_n} \rightarrow W$ a.s. Hence

$$0 \leq U_{\sigma_n} = \xi_{\sigma_n} \cdot Y - W_{\sigma_n} \rightarrow \xi \cdot Y - W =: U \text{ a.s.}$$

Hence we have $W = \xi \cdot Y - U$ with $U \in L_+^0$, i.e. W belongs to $\mathcal{K} - L_+^0$. \square

Having proved that \mathcal{C} is closed in L^1 , we shall show the existence of a $Z^* \in \mathcal{Z}$ (recall that \mathcal{Z} is defined in (8.7)) that is strictly positive \mathbb{P} -a.s, Theorem 8.23 below. First we need an auxiliary result.

Lemma 8.22 *Assume that $(\mathcal{K} - L_+^0) \cap L_+^1 = \{0\}$. If U is a non-negative element of L^1 and $\mathbb{P}(U > 0) > 0$, then there exists $Z \in \mathcal{Z}$ such that $\mathbb{E}(UZ) > 0$.*

Proof We use the Hahn-Banach theorem in the version of Corollary A.9. By the hypothesis we have that U is not an element of the nonempty convex set \mathcal{C} , which is closed in L^1 by Proposition 8.21. Hence there exists a $Z' \in L^\infty$ with $\sup\{\mathbb{E}(WZ') : W \in \mathcal{C}\} < \mathbb{E}(UZ') < \infty$. Since $0 \in \mathcal{C}$, it follows that $\mathbb{E}(UZ') > 0$. Moreover we also have $\beta := \sup\{\mathbb{E}(WZ') : W \in \mathcal{C}\} < \infty$ and then even $\beta \leq 0$. Indeed, for $W \in \mathcal{C}$ we have $\mathbb{E}(WZ') \leq \beta$ and since $\lambda W \in \mathcal{C}$ for every $\lambda > 0$, also $\lambda \mathbb{E}(WZ') \leq \beta$, hence $\mathbb{E}(WZ') \leq \beta/\lambda$, for every $\lambda > 0$. Hence $\mathbb{E}(WZ') \leq 0$ and $\beta \leq 0$ follows since $W \in \mathcal{C}$ was arbitrary. It now follows from Lemma 8.17 that $Z' \geq 0$ a.s. We conclude that $Z := Z'/\|Z'\|_\infty$ belongs to \mathcal{Z} and has the property $\mathbb{E}(UZ) > 0$. \square

Theorem 8.23 *Assume that $(\mathcal{K} - L_+^0) \cap L_+^1 = \{0\}$. Then there exists a $Z^* \in \mathcal{Z}$ with $\mathbb{P}(Z^* > 0) = 1$.*

Proof Let $\alpha := \sup\{\mathbb{P}(Z > 0) : Z \in \mathcal{Z}\} \leq 1$. By definition of α , there exists a sequence $(Z_n) \subset \mathcal{Z}$ such that $\mathbb{P}(Z_n > 0) \uparrow \alpha$. Let $Z^* := \sum_{n \geq 1} 2^{-n} Z_n$. Check, use the dominated convergence theorem, that this infinite sum belongs to \mathcal{Z} as well. Since for every n it holds that $\mathbb{P}(Z^n > 0) \leq \mathbb{P}(Z^* > 0)$, it follows that $\mathbb{P}(Z^* > 0) = \alpha$.

To show that $\alpha = 1$, we assume the contrary, $\mathbb{P}(Z^* = 0) > 0$ and construct a $Z' \in \mathcal{Z}$ with $\mathbb{P}(Z' > 0) > \alpha$. So let $\mathbb{P}(Z^* = 0) > 0$, then $U = \mathbf{1}_{\{Z^*=0\}}$ is nonnegative and $\mathbb{P}(U > 0) > 0$. Lemma 8.22 yields the existence of $Z \in \mathcal{Z}$ such that $\mathbb{E}(\mathbf{1}_{\{Z^*=0\}}Z) > 0$ and we must have $\mathbb{P}(\mathbf{1}_{\{Z^*=0\}}Z > 0) > 0$, so $\mathbb{P}(Z^* = 0, Z > 0) > 0$. Let now $Z' = \frac{1}{2}(Z + Z^*)$. One verifies that $Z' \in \mathcal{Z}$ and

$$\begin{aligned} \mathbb{P}(Z' > 0) &= \mathbb{P}(Z + Z^* > 0, Z^* > 0) + \mathbb{P}(Z + Z^* > 0, Z^* = 0) \\ &= \mathbb{P}(Z^* > 0) + \mathbb{P}(Z > 0, Z^* = 0) > \alpha, \end{aligned}$$

a contradiction. \square

Here is Lemma 8.11, formulated in agreement with the terminology and notation of the present section.

Corollary 8.24 *If the one-period market is arbitrage free, $\mathcal{K} \cap L_+^0 = \{0\}$, there exists an equivalent martingale measure \mathbb{P}^* such that $\frac{d\mathbb{P}^*}{d\mathbb{P}}$ is bounded. Conversely, if there exists an equivalent martingale measure \mathbb{P}^* , the market is arbitrage free.*

Proof Assume that the market is arbitrage free. In view of Lemma 8.16, we have $(\mathcal{K} - L_+^0) \cap L_+^0 = \{0\}$. But then also $\mathcal{C} \cap L_+^1 = (\mathcal{K} - L_+^0) \cap L_+^1 = \{0\}$. Theorem 8.23 yields the existence of $Z^* \in \mathcal{Z}$ such that $\mathbb{P}(Z^* > 0) = 1$. Then \mathbb{P}^* defined by

$$\frac{d\mathbb{P}^*}{d\mathbb{P}} = \frac{Z^*}{\mathbb{E} Z^*}$$

is a probability measure equivalent to \mathbb{P} and a martingale measure in view of Lemma 8.17. Since $Z^* \in \mathcal{Z}$, it is bounded.

Conversely, take a risk-neutral measure \mathbb{P}^* and a \mathcal{F}_0 -measurable ξ such that $\xi \cdot Y \in \mathcal{K} \cap L_+^0(\Omega, \mathcal{F}, \mathbb{P})$. Then for all $m > 0$ the random variable $\mathbf{1}_{\{|\xi| \leq m\}} \xi \cdot Y$ belongs to $\mathcal{K} \cap L_+^1(\Omega, \mathcal{F}, \mathbb{P}^*)$, because its expectation $\mathbb{E}^* \mathbf{1}_{\{|\xi| \leq m\}} \xi \cdot Y$ is well defined. Hence we have we have

$$\mathbb{E}^* \mathbf{1}_{\{|\xi| \leq m\}} \xi \cdot Y = \mathbb{E}^*(\mathbf{1}_{\{|\xi| \leq m\}} \xi \cdot \mathbb{E}^*[Y|\mathcal{F}_0]) = 0.$$

It follows that $\mathbf{1}_{\{|\xi| \leq m\}} \xi \cdot Y = 0$, \mathbb{P}^* -a.s. for all m and therefore $\xi \cdot Y = 0$, \mathbb{P}^* -a.s. By equivalence, $\xi \cdot Y = 0$, \mathbb{P} -a.s. too. Hence ξ does not yield an arbitrage opportunity. \square

8.4 European contingent claims

In this section we study the valuation problem for European contingent claims. *The standing assumption is that the market is arbitrage-free.*

Definition 8.25 A *contingent claim* C is a nonnegative \mathcal{F}_T -measurable random variable. It is called a *derivative* of the underlying assets, if C is measurable w.r.t. the σ -algebra $\sigma(S_0, \dots, S_T)$.

Occasionally we will extend the definition of a contingent claim to include random variables that allow negative values as well, although we will always impose that they are lower bounded. If a claim C is a derivative, then there exists a Borel function $f : (\mathbb{R}^{d+1})^{T+1} \rightarrow \mathbb{R}$ such that $C = f(S_0, \dots, S_T)$.

We give some examples of contingent claims. The first one is $C = (S_T^i - K)^+$, the European call option on S^i with maturity T and strike price K . An Asian option is for instance the claim $C = (\frac{1}{T+1} \sum_{t=0}^T S_t^i - K)^+$. A knock-in option is for instance $C = \mathbf{1}_{\{\max_{0 \leq t \leq T} S_t^i \geq B\}}$, where B is a (nonnegative) constant.

Definition 8.26 A contingent claim C is called attainable if there exists a self-financing trading strategy $\bar{\xi}$ such that $C = \bar{\xi}_T \cdot \bar{S}_T$. Such a strategy is called a *replicating* or *hedge* strategy for C .

The discounted value of a claim C is given by

$$H = \frac{C}{S_T^0}.$$

If the claim is attainable, then we also have

$$H = \bar{\xi}_T \cdot \bar{X}_T = V_T = V_0 + \sum_{t=1}^T \xi_t \cdot \Delta X_t,$$

and we will also say that H is attainable. Notice that $H \geq 0$ a.s.

Proposition 8.27 *Let \mathbb{P}^* be any equivalent martingale measure and H an attainable claim. Then $\mathbb{E}^* H < \infty$. If $\bar{\xi}$ is a replicating strategy, then*

$$(8.8) \quad V_t = \mathbb{E}^*[H|\mathcal{F}_t] \text{ a.s.},$$

for all $t = 0, \dots, T$, hence V is a nonnegative martingale under \mathbb{P}^* .

Proof This follows from Theorem 8.9, since $H = V_T \geq 0$. □

Notice that this proposition concerns the discounted value of the claim. Of course, if S_T^0 is deterministic, also C has finite expectation under each equivalent martingale measure. Moreover, it has two important consequences. The first one is that V_t , although it can be viewed as a conditional expectation, is the same for every equivalent martingale measure and the second one is that every replicating strategy for H has the same value process. Considering Equation (8.8) for $t = 0$, we obtain $V_0 = \mathbb{E}^*[H]$, which has the interpretation as the unique arbitrage free price of the discounted attainable claim H . Any other price would result in an arbitrage opportunity, see the arguments for the corresponding statement in Section 1.

Equation (8.8) can be rewritten as

$$\bar{\xi}_t \cdot \bar{S}_t = S_t^0 \mathbb{E}^*\left[\frac{C}{S_T^0} \middle| \mathcal{F}_t\right],$$

which for $t = 0$ yields the initial investment to purchase the replicating strategy,

$$V_0 = S_0^0 \mathbb{E}^* \frac{C}{S_T^0} = \mathbb{E}^* \frac{C}{S_T^0}.$$

This number can be interpreted as the fair price (at $t = 0$) of the undiscounted claim C . For non-attainable claims we have the following formal definition (compare also to Definition 1.11).

Definition 8.28 A nonnegative real number π^H is called an arbitrage-free price (at $t = 0$) of a discounted contingent claim H , if there exists an adapted process X^{d+1} such that a.s.

$$X_0^{d+1} = \pi^H,$$

$$\begin{aligned} X_t^{d+1} &\geq 0, \text{ for } t = 1, \dots, T-1, \\ X_T^{d+1} &= H, \end{aligned}$$

and if the extended market with (discounted) price process (X^1, \dots, X^{d+1}) is arbitrage-free. The set of all arbitrage-free prices is denoted by $\Pi(H)$.

It is mathematically more convenient to define a price for the discounted claim H . But of course, this is equivalent to a similar definition of an arbitrage price π^C for the undiscounted claim C . One has $\pi^C = S_0^0 \pi^H$. Since one usually takes $S_0^0 = 1$, it follows that $\pi^C = \pi^H$.

Definition 8.28 is the dynamic counterpart of Definition 1.11. Note that for an attainable discounted claim H , one can take $X_t^{d+1} = \mathbb{E}^*[H|\mathcal{F}_t]$, which is equal to the value V_t of a replication strategy, to see that the fair price of H is equal to V_0 . This is in agreement with Proposition 8.27 and the discussion after it.

Our first result in the valuation of claims is presented below, see Theorem 1.12 and its proof for the static version.

Theorem 8.29 *The set $\Pi(H)$ is non-empty and one has*

$$(8.9) \quad \Pi(H) = \{\mathbb{E}^*H : \mathbb{P}^* \in \mathcal{P}, \mathbb{E}^*H < \infty\}.$$

*Moreover, the upper and lower bounds of $\Pi(H)$ are given by $\sup\{\mathbb{E}^*H : \mathbb{P}^* \in \mathcal{P}\}$ and $\inf\{\mathbb{E}^*H : \mathbb{P}^* \in \mathcal{P}\}$ respectively.*

Proof First we show that the set on the right hand side of (8.9) is non-empty. Define a probability measure \mathbb{P}' on \mathcal{F}_T by

$$\frac{d\mathbb{P}'}{d\mathbb{P}} = \frac{c}{H+1},$$

where c is the normalization constant. Then $\mathbb{P}' \sim \mathbb{P}$, hence under \mathbb{P}' the market is arbitrage-free, and $\mathbb{E}_{\mathbb{P}'}H < \infty$. According to Theorem 8.12, there exists a risk-neutral measure \mathbb{P}^* such that $\frac{d\mathbb{P}^*}{d\mathbb{P}'}$ is bounded. But then $\mathbb{E}^*H < \infty$, and thus belongs to $\{\mathbb{E}^*H : \mathbb{P}^* \in \mathcal{P}, \mathbb{E}^*H < \infty\}$.

Next we prove (8.9). Take $\pi^H \in \Pi(H)$, recall Definition 8.28 and apply Theorem 8.12 to the extended market. This yields the existence of a probability measure \mathbb{P}^* on \mathcal{F}_T such that the X^i become martingales for $i = 1, \dots, d+1$. But this implies that $\mathbb{P}^* \in \mathcal{P}$ and $\pi^H = X_0^{d+1} = \mathbb{E}^*X_T^{d+1} = \mathbb{E}^*H$. So $\pi^H \in \{\mathbb{E}^*H : \mathbb{P}^* \in \mathcal{P}, \mathbb{E}^*H < \infty\}$.

Conversely, take $\mathbb{P}^* \in \mathcal{P}$ such that $\mathbb{E}^*H < \infty$. Define $X_t^{d+1} = \mathbb{E}^*[H|\mathcal{F}_t]$. Then \mathbb{P}^* is an equivalent martingale measure for the extended market, which is thus arbitrage free, and the requirements of Definition 8.28 are met with $\pi^H = \mathbb{E}^*H$. By the first part of the proof we now also know that $\Pi(H) \neq \emptyset$.

That $\inf \Pi(H) = \inf\{\mathbb{E}^*H : \mathbb{P}^* \in \mathcal{P}\}$ is trivial. To show the companion statement, we note that we only have to consider the case in which $\{\mathbb{E}^*H : \mathbb{P}^* \in \mathcal{P}\}$ differs from $\Pi(H)$, which happens if there exists some $\mathbb{P}_\infty \in \mathcal{P}$ such that

$\mathbb{E}_{\mathbb{P}_\infty} H = \infty$. The desired equality follows, as soon as we can show that for all $c > 0$, there exists a $\mathbb{P}_c \in \mathcal{P}$ such that $\infty > \mathbb{E}_{\mathbb{P}_c} H > c$. Indeed, in this case we have by the first part of the theorem that $\mathbb{E}_{\mathbb{P}_c} H \in \Pi(H)$ and it follows that $\sup \Pi(H) = \infty$.

First we note that for given $c > 0$, by monotone convergence, there exists n such that $\pi_n := \mathbb{E}_{\mathbb{P}_\infty} H \wedge n > c$. Put $X_t^{d+1} = \mathbb{E}_{\mathbb{P}_\infty} [H \wedge n | \mathcal{F}_t]$. Then \mathbb{P}_∞ becomes an equivalent martingale measure in the extended market with the additional asset $H \wedge n$, which is free of arbitrage, when the price vector is extended with π_n . Application of the first part of the theorem to the extended market then shows that for any contingent claim, in particular H , there exists a \mathbb{P}_c equivalent to \mathbb{P}_∞ such that $\mathbb{E}_{\mathbb{P}_c} H < \infty$. But then this \mathbb{P}_c is also an equivalent martingale measure for the original market and thus $\mathbb{E}_{\mathbb{P}_c} H \in \Pi(H)$. On the other hand, the price process X^{d+1} is a martingale under \mathbb{P}_c as well, and so $\mathbb{E}_{\mathbb{P}_c} X_T^{d+1} = X_0^{d+1}$. Using this fact, we have

$$\mathbb{E}_{\mathbb{P}_c} H \geq \mathbb{E}_{\mathbb{P}_c} H \wedge n = \mathbb{E}_{\mathbb{P}_c} X_T^{d+1} = X_0^{d+1} = \pi_n > c,$$

which finishes the proof for the supremum. \square

We extend results of Section 1 to a dynamic setting. Recall Proposition 1.18 and consider its dynamic version.

Theorem 8.30 *Assume the market to be arbitrage free. Let H be a discounted claim. If H is attainable, $\Pi(H)$ consists of one element, the value at $t = 0$ of any replicating portfolio. If H is not attainable, then $\Pi(H)$ is an open interval.*

Proof If H is attainable, then the assertion follows from Theorem 8.29 combined with the discussion after Proposition 8.27.

The proof of the other case is much more involved. As in the proof of Proposition 1.18 we observe that $\Pi(H)$ is convex and thus an interval. We will show that it is open. To that end, let $\pi \in \Pi(H)$. It is sufficient to show that there are $\pi_0, \pi_1 \in \Pi(H)$ such that $\pi_0 < \pi < \pi_1$. We first construct π_1 .

Take $\mathbb{P}^* \in \mathcal{P}$ such that $\mathbb{E}^* H = \pi$ and let $M_t = \mathbb{E}^*[H | \mathcal{F}_t]$. Then

$$H = M_0 + \sum_{t=1}^T \Delta M_t.$$

Since H is not attainable, there must be some $t \in \{1, \dots, T\}$ such that ΔM_t can not be written as $\xi_t \cdot \Delta X_t$, for some \mathcal{F}_{t-1} -measurable ξ_t with $\xi_t \cdot \Delta X_t \in L^1(\Omega, \mathcal{F}_t, \mathbb{P}^*)$. It follows from the proof of Proposition 8.21 (take all $U_n = 0$ there) that the collection \mathcal{C}_t of all random variables that are a.s. equal to such a $\xi_t \cdot \Delta X_t$ is a closed linear subspace of $L^1(\Omega, \mathcal{F}_t, \mathbb{P}^*)$ (and thus convex as well). We apply the infinite dimensional version of the separating hyperplane theorem of Corollary A.9 to conclude that there exists a $Z \in L^\infty(\Omega, \mathcal{F}_t, \mathbb{P}^*)$ such that

$$\sup_{W \in \mathcal{C}_t} \mathbb{E}^* W Z < \mathbb{E}^* \Delta M_t Z < \infty.$$

If we replace in the above inequality W with αW for arbitrary $\alpha \in \mathbb{R}$, then by linearity the inequality can only be preserved if

$$(8.10) \quad \mathbb{E}^* W Z = 0, \forall W \in \mathcal{C}_t.$$

We conclude that

$$(8.11) \quad \mathbb{E}^* \Delta M_t Z > 0.$$

By multiplying Z by a sufficiently small positive number, if necessary, we may assume that (8.11) is true for a random variable Z with $\mathbb{P}^*(|Z| < \frac{1}{2}) = 1$. Let

$$Z_t = 1 + Z - \mathbb{E}^*[Z|\mathcal{F}_{t-1}].$$

Then $\mathbb{P}^*(0 < Z_t < 2) = 1$, $\mathbb{E}^* Z_t = 1$ and $\frac{d\mathbb{P}_t}{d\mathbb{P}^*} = Z_t$ defines a probability measure $\mathbb{P}_t \sim \mathbb{P}^*$. Notice that Z_t is \mathcal{F}_t -measurable. We compute

$$\begin{aligned} \mathbb{E}_{\mathbb{P}_t} H &= \mathbb{E}^* H Z_t \\ &= \mathbb{E}^* H + \mathbb{E}^*(\mathbb{E}^*[H|\mathcal{F}_t]Z) - \mathbb{E}^*(H\mathbb{E}^*[Z|\mathcal{F}_{t-1}]) \\ &= \mathbb{E}^* H + \mathbb{E}^* M_t Z - \mathbb{E}^*(\mathbb{E}^*[H|\mathcal{F}_{t-1}]\mathbb{E}^*[Z|\mathcal{F}_{t-1}]) \\ &= \mathbb{E}^* H + \mathbb{E}^* M_t Z - \mathbb{E}^*(M_{t-1}\mathbb{E}^*[Z|\mathcal{F}_{t-1}]) \\ &= \mathbb{E}^* H + \mathbb{E}^* M_t Z - \mathbb{E}^* M_{t-1} Z \\ &= \mathbb{E}^* H + \mathbb{E}^*(\Delta M_t Z) \\ &> \mathbb{E}^* H, \end{aligned}$$

in view of (8.11). Since $\mathbb{E}_{\mathbb{P}_t} H = \mathbb{E}^* H Z_t \leq 2\mathbb{E}^* H < \infty$, we can take $\pi_1 = \mathbb{E}_{\mathbb{P}_t} H$ and then

$$(8.12) \quad \pi_1 > \mathbb{E}^* H.$$

Hence we have reached our aim, provided that \mathbb{P}_t belongs to \mathcal{P} , which we are going to prove now.

Let $k > t$. Since Z_t is \mathcal{F}_t -measurable and hence \mathcal{F}_{k-1} -measurable, we have

$$\begin{aligned} \mathbb{E}_{\mathbb{P}_t}[\Delta X_k | \mathcal{F}_{k-1}] &= \frac{\mathbb{E}^*[\Delta X_k Z_t | \mathcal{F}_{k-1}]}{\mathbb{E}^*[Z_t | \mathcal{F}_{k-1}]} \\ &= \mathbb{E}^*[\Delta X_k | \mathcal{F}_{k-1}] = 0. \end{aligned}$$

For $k = t$, we also have $\mathbb{E}^*[\Delta X_t Z | \mathcal{F}_{t-1}] = 0$. Indeed, let $F \in \mathcal{F}_{t-1}$ arbitrary. Because of $\mathbf{1}_F \in \mathcal{F}_{t-1}$ and (8.10), it holds that $\mathbb{E}^*(\mathbf{1}_F \Delta X_t Z) = 0$. Notice also that $\mathbb{E}^*[Z_t | \mathcal{F}_{t-1}] = 1$, straight from the definition of Z_t . But then

$$\begin{aligned} \mathbb{E}_{\mathbb{P}_t}[\Delta X_t | \mathcal{F}_{t-1}] &= \mathbb{E}^*[\Delta X_t (1 - \mathbb{E}^*[Z | \mathcal{F}_{t-1}]) | \mathcal{F}_{t-1}] + \mathbb{E}^*[\Delta X_t Z | \mathcal{F}_{t-1}] \\ &= (1 - \mathbb{E}^*[Z | \mathcal{F}_{t-1}])\mathbb{E}^*[\Delta X_t | \mathcal{F}_{t-1}] + 0 = 0. \end{aligned}$$

The third case, $k < t$, is easy. Since then $\mathbb{E}^*[Z_t | \mathcal{F}_k] = 1$, one obtains $\mathbb{E}^*[\frac{d\mathbb{P}_t}{d\mathbb{P}^*} | \mathcal{F}_k] = 1$, so the measures \mathbb{P}_t and \mathbb{P}^* coincide on \mathcal{F}_k : $\mathbb{P}_t(A) = \mathbb{E}^*\mathbb{E}^*[\mathbf{1}_A Z_t | \mathcal{F}_k] = \mathbb{E}^*\mathbf{1}_A \mathbb{E}^*[Z_t | \mathcal{F}_k] = \mathbb{P}^*(A)$ for $A \in \mathcal{F}_k$. Hence

$$\mathbb{E}_{\mathbb{P}_t}[\Delta X_k | \mathcal{F}_{k-1}] = \mathbb{E}^*[\Delta X_k | \mathcal{F}_{k-1}] = 0.$$

We conclude that \mathbb{P}_t is an equivalent martingale measure and hence $\pi_1 \in \Pi(H)$.

We turn to the construction of π_0 . Let

$$\frac{d\mathbb{P}_0}{d\mathbb{P}^*} = 2 - Z_t.$$

Then $\mathbb{P}^*(0 < \frac{d\mathbb{P}_0}{d\mathbb{P}^*} < 2) = 1$ and $\mathbb{E}^* \frac{d\mathbb{P}_0}{d\mathbb{P}^*} = 1$. Hence also \mathbb{P}_0 is a probability measure, equivalent to \mathbb{P}^* , a martingale measure as well (this follows from \mathbb{P}_t being a martingale measure) and

$$\mathbb{E}_{\mathbb{P}_0} H = \mathbb{E}^*(2 - Z_t)H = 2\mathbb{E}^* H - \mathbb{E}^* Z_t H = 2\mathbb{E}^* H - \pi_1 < \mathbb{E}^* H = \pi,$$

by (8.12). Taking $\pi_0 = \mathbb{E}_{\mathbb{P}_0} H$ completes the proof. \square

8.5 Complete markets

The definition of a complete market looks the same as for the static case, Definition 1.19, but it involves the more subtle notion of attainability in the multi period setting as in Definition 8.26.

Definition 8.31 An arbitrage-free market is *complete*, if every contingent claim is attainable.

A consequence of a market being complete is that every contingent claim has a unique price, in view of Theorem 8.30. We now present what is sometimes called the *Second Fundamental Theorem of Asset Pricing*, see also Theorem 1.21.

Theorem 8.32 *An arbitrage-free market is complete iff there exists a unique equivalent martingale measure. The number of atoms of $(\Omega, \mathcal{F}, \mathbb{P})$ in case of a complete market is at most $(d + 1)^T$. Moreover, $\dim L^0(\Omega, \mathcal{F}, \mathbb{P}) \leq (d + 1)^T$.*

Proof If the market is complete, we argue as in the proof of Theorem 1.21. Every claim $\mathbf{1}_F$, with $F \in \mathcal{F} = \mathcal{F}_T$ has a unique price. Hence there is a unique \mathbb{P}^* . Conversely, if there exists only one equivalent martingale measure, the result follows from Theorem 8.30.

We turn to the number of atoms. We have seen the statement to be true for $T = 1$ in Theorem 1.21 and we proceed by induction. Suppose that the assertion is true for a time horizon $T - 1$. By completeness, every claim can be replicated. So if H is a bounded nonnegative discounted claim, there is a replicating strategy $\bar{\xi}$ with value process V such that

$$H = V_{T-1} + \xi_T \cdot \Delta X_T.$$

Since V_{T-1} and ξ_T are \mathcal{F}_{T-1} -measurable, they are constant on atoms A that belong to \mathcal{F}_{T-1} . Consider for such A the restricted probability space $(A, \mathcal{F}_T^A, \mathbb{P}^A)$, where $\mathcal{F}_T^A = \{F \cap A : F \in \mathcal{F}_T\}$, and \mathbb{P}^A the conditional probability $\mathbb{P}(\cdot | A)$ restricted to \mathcal{F}_T^A . As we just said, on this restricted probability space V_{T-1} and ξ_T are constant. Hence Theorem 1.21 applies and so the dimension of

$L^\infty(A, \mathcal{F}_T^A, \mathbb{P}^A)$ is at most $d+1$. Then Proposition 1.20 implies that $(A, \mathcal{F}_T^A, \mathbb{P}^A)$ has at most $d+1$ atoms.

Every atom of $(\Omega, \mathcal{F}_T, \mathbb{P})$ is an atom of some $(A, \mathcal{F}_T^A, \mathbb{P}^A)$. Applying the induction hypothesis, we know that there are at most $(d+1)^{T-1}$ of such restricted probability spaces. The conclusion follows by multiplication, and it implies the assertion on the dimension. \square

Consider the set \mathcal{Q} of all martingale measures as in Definition 8.8, it is a convex set. Likewise the set of equivalent martingale measures \mathcal{P} is convex. We will see below that complete markets can be characterized by extreme points of those convex sets. Recall that an extreme point of a convex set is such that it doesn't admit a non-trivial convex combination of points in the convex set.

Theorem 8.33 *Let $\mathbb{P}^* \in \mathcal{P}$. The following are equivalent.*

- (i) $\mathcal{P} = \{\mathbb{P}^*\}$ (the market is complete).
- (ii) \mathbb{P}^* is an extreme point of \mathcal{P} .
- (iii) \mathbb{P}^* is an extreme point of \mathcal{Q} .
- (iv) If M is a martingale under \mathbb{P}^* , then there exists a d -dimensional predictable process ξ , such that

$$M_t = M_0 + \sum_{k=1}^t \xi_k \cdot \Delta X_k, \quad t \in \{0, \dots, T\}.$$

Proof (i) \Rightarrow (iii): Write $\mathbb{P}^* = \alpha \mathbb{Q}_1 + (1 - \alpha) \mathbb{Q}_2$ for $\alpha \in (0, 1)$ and $\mathbb{Q}_1, \mathbb{Q}_2 \in \mathcal{Q}$. Then \mathbb{Q}_1 and \mathbb{Q}_2 are necessarily absolutely continuous w.r.t. \mathbb{P}^* . But also $\mathbb{P}_i = \frac{1}{2}(\mathbb{Q}_i + \mathbb{P}^*)$ ($i = 1, 2$), being convex combinations of martingale measures, are martingale measures too, and equivalent to \mathbb{P}^* . From the assumption it follows that $\mathbb{P}_1 = \mathbb{P}_2 = \mathbb{P}^*$ and then also $\mathbb{Q}_1 = \mathbb{Q}_2 = \mathbb{P}^*$.

(iii) \Rightarrow (ii): Trivial, since $\mathcal{P} \subset \mathcal{Q}$.

(ii) \Rightarrow (i): Let $\mathbb{P}_0^* \in \mathcal{P}$, different from \mathbb{P}^* . We first show that we can find a $\mathbb{P}_1^* \in \mathcal{P}$ different from \mathbb{P}^* such that $\frac{d\mathbb{P}_1^*}{d\mathbb{P}^*}$ is bounded. Since $\mathbb{P}_0^* \neq \mathbb{P}^*$, there must be a set $E \in \mathcal{F}_T$ such that $\mathbb{P}^*(E) \neq \mathbb{P}_0^*(E)$. Extend the market with the \mathbb{P}_0^* -martingale X^{d+1} given by $X_t^{d+1} = \mathbb{E}_{\mathbb{P}_0^*}[\mathbf{1}_E | \mathcal{F}_t]$. Choose $(\Omega, \mathcal{F}_T, \mathbb{P}^*)$ to be the underlying probability space. It follows by construction, that \mathbb{P}_0^* is an equivalent martingale measure for the extended market, and so the extended market is also arbitrage free under \mathbb{P}^* . Theorem 8.12 then provides the existence of a probability measure \mathbb{P}_1^* equivalent to \mathbb{P}^* such that $\frac{d\mathbb{P}_1^*}{d\mathbb{P}^*}$ is bounded by some $B > 0$, which is a risk-neutral measure for the extended market. Since

$$X_0^{d+1} = \mathbb{P}_0^*(E) \neq \mathbb{P}^*(E) = \mathbb{E}^* X_T^{d+1}$$

\mathbb{P}^* is not a martingale measure for the extended market, and hence \mathbb{P}_1^* must be different from \mathbb{P}^* .

Choose $\varepsilon < 1/B$ and put

$$Z = 1 + \varepsilon - \varepsilon \frac{d\mathbb{P}_1^*}{d\mathbb{P}^*}.$$

Notice that $\varepsilon \leq Z \leq 1 + \varepsilon$ and $\mathbb{E}^* Z = 1$. Hence

$$\frac{d\mathbb{P}_2^*}{d\mathbb{P}^*} = Z$$

defines a probability measure $\mathbb{P}_2^* \in \mathcal{P}$ (that \mathbb{P}_2^* is a martingale measure is Exercise 8.9) with bounded density Z . Moreover, \mathbb{P}^* turns out to be a convex combination,

$$\mathbb{P}^* = \frac{\varepsilon}{1 + \varepsilon} \mathbb{P}_1^* + \frac{1}{1 + \varepsilon} \mathbb{P}_2^*,$$

which contradicts that \mathbb{P}^* is extreme.

(i) \Rightarrow (iv): Let M be a positive \mathbb{P}^* -martingale. Then we can see M_T as a contingent claim, which is attainable by Theorem 8.32. Let $\bar{\xi}$ be a replicating strategy. Then, \mathbb{P}^* -a.s.,

$$M_T = V_0 + \sum_{k=1}^T \bar{\xi}_k \cdot \Delta X_k.$$

By Proposition 8.27, the corresponding value process V , with

$$V_t = V_0 + \sum_{k=1}^t \bar{\xi}_k \cdot \Delta X_k = \mathbb{E}^*[M_T | \mathcal{F}_t],$$

is a martingale and so $M_t = V_t$, which proves the assertion for positive martingales. The general case follows by the decomposition $M_T = M_T^+ - M_T^-$.

(iv) \Rightarrow (i): Pick $E \in \mathcal{F}_T$ and put $M_t = \mathbb{E}^*[\mathbf{1}_E | \mathcal{F}_t]$, $t \in \{0, \dots, T\}$. Then the assumption implies that $\mathbf{1}_E$ is an attainable claim. According to Theorem 8.30, it has a unique arbitrage free price. Hence for all $\mathbb{P}^* \in \mathcal{P}$, we have that $\mathbb{P}^*(E)$ is one and the same number. Since E is arbitrary, \mathcal{P} must be a singleton. But then the market is complete in view of Theorem 8.32. \square

Remark 8.34 Property (iv) of Theorem 8.33 is also called the discrete time Martingale Representation Theorem, similar to a theorem for so called Brownian martingales in continuous time.

8.6 CRR model

In this section we consider the Cox-Ross-Rubinstein (CRR) model, a popular model of a financial market in discrete time. Apart from its tractability and that of related pricing issues, it is also interesting, because pricing formulas tend to Black-Scholes related formulas under the right kind of asymptotics. We will not treat this aspect in the present course.

In the CRR model, there is a riskless asset, whose price evolves according to

$$S_t^0 = (1 + r)^t,$$

for some $r \in (-1, \infty)$, although usually $r \geq 0$. There is only one risky asset with price process $S^1 = S$, whose *returns*

$$R_t := \frac{\Delta S_t}{S_{t-1}}$$

are random variables greater than -1 and assume for each time t only two values. Often these values are taken to be the same, as we will do, for all $t \geq 1$, say a and b , with $a < b$. The simplest probability space that carries all random variables below, assuming a finite time horizon T , is $\Omega = \{a, b\}^T$. The obvious filtration is such that $\mathcal{F}_t = \sigma(R_1, \dots, R_t)$, $t \in \{0, \dots, T\}$. In this case, any sensible probability measure on \mathcal{F}_T must be such that all singletons have positive probability. The totality of all these conventions will be referred to as the CRR model. We will see that absence of arbitrage has a simple characterization in terms of the parameters a and b .

We use X to denote the discounted price process of the risky asset, so

$$X_t = \frac{S_t}{S_t^0},$$

and note that

$$\begin{aligned} X_t &= \frac{1 + R_t}{1 + r} X_{t-1}, \\ \Delta X_t &= \frac{R_t - r}{1 + r} X_{t-1}. \end{aligned}$$

Proposition 8.35 *The CRR model is arbitrage-free iff $a < r < b$. Moreover, if it is arbitrage-free, it is also complete. The unique equivalent martingale measure is such that the R_t become i.i.d. random variables, whose common distribution is determined by*

$$\mathbb{P}^*(R_t = b) = \frac{r - a}{b - a} =: p^*.$$

Proof First we show that if a martingale measure exists, it is necessarily unique. Let \mathbb{Q} be a martingale measure. Then

$$X_{t-1} = \mathbb{E}_{\mathbb{Q}}[X_t | \mathcal{F}_{t-1}] = X_{t-1} \mathbb{E}_{\mathbb{Q}}\left[\frac{1 + R_t}{1 + r} | \mathcal{F}_{t-1}\right].$$

Since X_t is positive \mathbb{Q} -a.s., we can solve this equation to get

$$(8.13) \quad \mathbb{E}_{\mathbb{Q}}[R_t | \mathcal{F}_{t-1}] = r.$$

Let $q = \mathbb{Q}(R_t = b | \mathcal{F}_{t-1}) = 1 - \mathbb{Q}(R_t = a | \mathcal{F}_{t-1})$. Then we can rewrite (8.13) as $qb + (1 - q)a = r$, which yields

$$q = \mathbb{Q}(R_t = b | \mathcal{F}_{t-1}) = \frac{r - a}{b - a}.$$

This implies that R_t is, under \mathbb{Q} , independent of \mathcal{F}_{t-1} and that its unconditional distribution is also given by $\mathbb{Q}(R_t = b) = q$. It follows that, necessarily, the R_t are i.i.d. under \mathbb{Q} , and hence \mathbb{Q} must be unique. Note that we have $q = p^*$. For \mathbb{Q} to be a probability measure, we need $p^* \in [0, 1]$, which is equivalent to $r \in [a, b]$. To have that \mathbb{Q} is in fact a risk-neutral measure, we need that $\mathbb{Q} \sim \mathbb{P}$, which means that $p^* \in \{0, 1\}$ is to be excluded. In that case $a < r < b$.

Let the market be arbitrage free. Then there exists an equivalent martingale measure \mathbb{P}^* . By the above reasoning, we necessarily have that \mathbb{P}^* is as asserted. The market is then also complete in view of Theorem 8.33.

If the condition $a < r < b$ holds true, then we can define the measure \mathbb{P}^* on Ω , by putting

$$\mathbb{P}^*(\{\omega\}) = (p^*)^{k(\omega)}(1 - p^*)^{T - k(\omega)},$$

where $k(\omega)$ denotes the number of b 's in ω . Clearly we have $\mathbb{P}^* \sim \mathbb{P}$, independence of the R_t follows and we also see that the marginal distribution of each R_t is the same as for the others. We have seen above that \mathbb{P}^* defines a martingale measure. \square

We turn to the pricing of contingent claims. Recall that they have a unique price by completeness of the market. Consider a discounted claim H . Since H is \mathcal{F}_T -measurable, there exists a function $h : \Omega \rightarrow \mathbb{R}$ such that

$$(8.14) \quad H = h(R_1, \dots, R_T).$$

The value process for H is, whatever replicating strategy (but in Exercise 8.8 it is shown to be unique) given by

$$V_t = \mathbb{E}^*[H | \mathcal{F}_t].$$

Put

$$v_t(r_1, \dots, r_t) = \mathbb{E}^*h(r_1, \dots, r_t, R_{t+1}, \dots, R_T).$$

Exploiting the independence of the R_t and using properties of conditional expectation, we get that

$$V_t = v_t(R_1, \dots, R_t).$$

Moreover, using the martingale property of V , we similarly obtain

$$v_{t-1}(r_1, \dots, r_{t-1}) = p^*v_t(r_1, \dots, r_{t-1}, b) + (1 - p^*)v_t(r_1, \dots, r_{t-1}, a).$$

If the discounted claim H only depends on the terminal price S_T , then we have $H = k(S_T)$, for some function k . The relation between k and the above h is

$$k(S_0(1 + r_1) \cdots (1 + r_T)) = h(r_1, \dots, r_T).$$

Put $w_t(s) = \mathbb{E}^*k(s(1 + R_{t+1}) \cdots (1 + R_T))$. Notice also, that we can alternatively write

$$(8.15) \quad w_t(s) = \mathbb{E}^*k\left(\frac{sS_T}{S_t}\right).$$

Between v_t and w_t one has the relation

$$v_t(r_1, \dots, r_t) = w_t(S_0(1 + r_1) \cdots (1 + r_t)),$$

and hence $V_t = w_t(S_t)$. See also Exercises 8.7 and 8.8.

8.7 Exercises

8.1 Prove Proposition 8.3.

8.2 Prove Proposition 8.14. *Hint:* Apply Jensen's inequality to $\mathbb{E}_{\mathbb{P}} \frac{X_T^0}{X_T^1}$.

8.3 Fix the time horizon at T and assume the initial σ -algebra \mathcal{F}_0 to be trivial. Let S_t^0 be identically equal to 1 and let $Z_t = \log \frac{S_t^1}{S_{t-1}^1}$. Suppose that the market that is described by the pair of processes S^0, S^1 is arbitrage-free. Suppose that \mathbb{P} is such that the Z_t are i.i.d. with a common normal distribution. Give a relation between the parameters if $\mathbb{P} \in \mathcal{P}$. Can $\exp(Z_t)$ have a Gamma distribution if $\mathbb{P} \in \mathcal{P}$?

8.4 Consider an arbitrage-free market with one risky asset. Let S^1 be its price process and S^0 the deterministic price process of the riskless asset. Consider a European call option with discounted payoff

$$H = \frac{(S_T^1 - K)^+}{S_T^0},$$

for some $K > 0$. Assume that S_T^1 has a density w.r.t. Lebesgue measure under any risk-neutral measure. Let π^* be an arbitrage-free price of the call option under some risk-neutral measure \mathbb{P}^* . Obviously π^* depends on K and S_0^1 , so we write $\pi^* = \pi^*(K, S_0^1)$. Show that

$$0 < \frac{\partial \pi^*}{\partial S_0^1} < 1$$

$$\frac{\partial \pi^*}{\partial K} = -\frac{1}{S_T^0} (1 - F^*(K)),$$

where F^* is the distribution function of S_T^1 under \mathbb{P}^* . To show the first assertion you may make additional assumptions, e.g. that S_T^1 is increasing in S_0^1 , or even more explicit, $S_T^1 = S_0^1 R_T$, with R_T a positive random variable.

8.5 Consider a market with underlying $\Omega = \{1, 2, 3, 4\}$. Assume that $T = 2$ and that $S_t^0 = 1$ for $t = 0, 1, 2$, the price of the riskless asset is constant. Let the evolution for the price S_t of the *single* risky asset be as given in the table.

ω	$S_0(\omega)$	$S_1(\omega)$	$S_2(\omega)$
1	5	8	9
2	5	8	6
3	5	4	6
4	5	4	3

- (a) Assume that \mathbb{P} gives positive probability to each singleton. Show that the market is complete and that \mathbb{P}^* as represented by the vector $(\frac{1}{6}, \frac{1}{12}, \frac{1}{4}, \frac{1}{2})$ is the unique equivalent martingale measure.

- (b) Let H be the European call option $H = (S_2 - 5)^+$. Let $\bar{\xi}$ be the replicating strategy. Show that $\bar{\xi}_2(\omega) = (-5, 1)$ if $\omega = 1, 2$ and $\bar{\xi}_2(\omega) = (-1, \frac{1}{3})$ if $\omega = 3, 4$.
- (c) Compute V_1 and show that $\bar{\xi}_1(\omega) = (-\frac{7}{3}, \frac{2}{3})$ for all ω . What is the value of the claim at $t = 0$?
- (d) As an alternative you can use the self-financing property in the form $V_t = V_{t-1} + \Delta G_t$, $t = 1, 2$. Use this to compute the replicating strategy anew.
- (e) Suppose that the riskless interest rate is r . For which possible values of r do we still have an arbitrage free market?

8.6 Consider the market of Exercise 8.5. Compute the value of the claim $H = (\frac{1}{3}(S_0 + S_1 + S_2) - 5)^+$.

8.7 Give an explicit formula for $w_t(s)$, see (8.15), as a sum involving the probabilities of the Binomial distribution with parameters $T - t$ and p^* .

8.8 Let H be a claim as in (8.14). Show that the hedge strategy is given by

$$\xi_t = (1 + r)^t \frac{v_t(R_1, \dots, R_{t-1}, b) - v_t(R_1, \dots, R_{t-1}, a)}{S_{t-1}(b - a)}.$$

Give also an expression for ξ_t , if $H = h(S_T)$. What is the explicit resulting strategy if $H = (1 + r)^{-T} S_T$?

8.9 Show that the probability measure \mathbb{P}_2^* in the proof of Theorem 8.33 is a martingale measure.

8.10 Let Y_1, \dots, Y_T be *iid* random variables on some $(\Omega, \mathcal{F}, \mathbb{P})$ with $\mathbb{E} Y_t = 0$ for all t and let $\mathcal{F}_t = \sigma(Y_1, \dots, Y_t)$. Let $X_t = \sum_{k=1}^t Y_k$ for $t \leq T$. Obviously, the X_t form a martingale. Consider an *insider trader*, a trader whose information pattern is given by the σ -algebras $\tilde{\mathcal{F}}_t := \sigma(\mathcal{F}_t \cup \sigma(X_T))$, i.e. at any time $t \leq T$ she ‘knows’ the final value X_T .

- (a) Show that the X_t don’t result in a martingale w.r.t. the enlarged filtration of the $\tilde{\mathcal{F}}_t$.
- (b) Let $\tilde{X}_t = X_t - \sum_{k=0}^{t-1} \frac{X_T - X_k}{T - k}$, $t \leq T$. Show that the \tilde{X}_t yield a martingale w.r.t. enlarged filtration. *Hint: use that $\mathbb{E}[X_t | X_T] = \frac{t}{T} X_T$ and independence of the Y_k .*
- (c) Construct a self-financing strategy of investments $\tilde{\xi}_t$ w.r.t. the enlarged filtration (so the $\tilde{\xi}_t$ are $\tilde{\mathcal{F}}_{t-1}$ -measurable) such that $\mathbb{E} \sum_{t=1}^T \xi_t (X_t - X_{t-1})$ is positive. This should follow from maximization of the expected gain $\mathbb{E} \sum_{t=1}^T \tilde{\xi}_t (X_t - X_{t-1})$ over all self-financing strategies such that $|\tilde{\xi}_t| \leq 1$.

8.11 Show that the σ_m^0 in the proof of Lemma 8.19 are \mathcal{F} -measurable, as well as the $\xi_{\sigma_m^0}$. Same question for the σ_m^1 and the $\xi_{\sigma_m^1}$. Finish the proof of Lemma 8.19.

8.12 The proof of Proposition 8.21 is a lot simpler if \mathcal{F}_0 is the trivial σ -algebra $\{\emptyset, \Omega\}$. In this case all ξ_n in the proof are just vectors in \mathbb{R}^d and the spaces N and N^\perp are closed linear subspaces of \mathbb{R}^d . Rewrite (and shorten) the proof under

this additional assumption and make clear that Lemma 8.19 and Lemma 8.20 can be circumvented by using standard analysis arguments instead.

8.13 Show that it follows from the proof of Proposition 8.21 that under the same assumption also \mathcal{K} is closed in L^0 .

8.14 If one drops the no arbitrage assumption in Proposition 8.21, the assertion is no longer true in general. This exercise contains an example. Assume that market contains only one risky asset ($d = 1$). Let in $(\Omega, \mathcal{F}, \mathbb{P})$, $\Omega = [0, 1]$, \mathcal{F} the Borel σ -algebra, and \mathbb{P} the Lebesgue measure. Assume that $Y : \Omega \rightarrow \mathbb{R}$ is given by $Y(\omega) = \omega$.

- (a) Show that the no arbitrage condition is violated.
- (b) Let $Z \geq 1$ be a constant. Show that Z cannot belong to \mathcal{C} , and conclude that \mathcal{C} is not all of L^1 .
- (c) Let $Z \in L^1$ and define $Z_n = (Z^+ \wedge n)\mathbf{1}_{[\frac{1}{n}, 1]} - Z^-$. Show that $Z_n \in \mathcal{C}$ (establish first that $(Z^+ \wedge n)\mathbf{1}_{[\frac{1}{n}, 1]} \leq c_n Y$ for some constant c_n) and that $Z_n \rightarrow Z$ in L^1 for $n \rightarrow \infty$.
- (d) Conclude that \mathcal{C} is not closed.

9 Optimization in dynamic models

In this section we study portfolio optimization over a nontrivial horizon. We thus extend the results of Section 6 to a dynamic case. We will present two methods to find an optimal portfolio, one is based on *Dynamic Programming*, the other is based on first finding an optimal random pay-off and then to construct a trading strategy that replicates this pay-off. This method is known as the *martingale method* or as the *risk neutral approach*.

9.1 Dynamic programming

Dynamic programming is a main tool in optimization for dynamic models, especially useful if the relevant underlying processes are defined by recursive models, or else have a model describing their time dependent behavior. This could mean e.g. that they are Markov processes or martingales. We first explain the two key ideas behind dynamic programming and then proceed with a more formal treatment. The proofs of the results in this section are either exercises or given in Appendix A.5

We give the ideas underlying dynamic programming in its most rudimentary form. Suppose one wants to maximize a function V of two variables, of the specific form

$$V(u_1, u_2) = V_1(u_1) + V_2(f(u_1), u_2).$$

This maximization problem can be carried out as the iterated maximization

$$\max_{u_1, u_2} V(u_1, u_2) = \max_{u_1} (V_1(u_1) + \max_{u_2} V_2(f(u_1), u_2)),$$

with the underlying idea that the functions V_i are to be interpreted as ‘rewards’ at ‘times’ $i = 1, 2$. The maximization on the right hand side over u_2 , with any u_1 fixed, yields (assuming a maximizer exists and is unique) an optimal

$$u_2^*(u_1) = g(u_1),$$

for some function g . Substitution of this relation for u_2 in $V(u_1, u_2)$ yields a function of u_1 only,

$$V(u_1, g(u_1)) = V_1(u_1) + V_2(u_1, g(u_1)),$$

and maximizing over u_1 yields an optimal solution u_1^* , that in turn yields $u_2^* = g(u_1^*)$.

So, the (first) idea is to optimize over the second variable and then over the first one, and that, given that the first one yields an optimal value, the optimum of the second step is immediately known. Hence, if one views the pair (u_1^*, u_2^*) as some kind of optimal path to reach one’s goal, then the second part of the path, u_2^* , is optimal once the ‘starting value’ u_1^* is given. This reflects the second idea behind dynamic programming, also called *Bellman’s optimality principle*.

We move on to a random dynamic setting, where all random variables and

processes are defined on some $(\Omega, \mathcal{F}, \mathbb{P})$. Suppose that one has an \mathbb{R}^d -valued stochastic process $X = (X_0, \dots, X_T)$, in fact it is going to be a family of processes as we shall soon see, that obey the recursion

$$(9.1) \quad X_{t+1} = f_t(X_t, U_t, \varepsilon_t),$$

and that start in some value X_0 .

The random variables $X_0, \varepsilon_0, \varepsilon_1, \dots, \varepsilon_{T-1}$ are given and they are assumed to be independent. The m -dimensional random variables U_t are supposed to be of the form

$$(9.2) \quad U_t = u_t(X_0, \dots, X_t),$$

for certain measurable functions $u_t : (\mathbb{R}^d)^{t+1} \rightarrow \mathbb{R}^m$. Likewise, the functions f_t are also assumed to be jointly measurable in their arguments. As a filtration we take the family of σ -algebras $\mathcal{F}_t = \sigma(X_0, \varepsilon_0, \dots, \varepsilon_{t-1})$. Then the processes X and U are adapted. Moreover, if $U_t = u_t(X_t)$ (as we shall see below, this is an important case), the resulting process X is even Markov.

Lemma 9.1 *The process X is Markov w.r.t. the filtration specified above, if U_t depends on X_t only, so $U_t = u_t(X_t)$, where the u_t are measurable functions $u_t : \mathbb{R}^d \rightarrow \mathbb{R}^m$.*

Proof Exercise 9.1. □

Remark 9.2 Under the conditions of Lemma 9.1, we have $X_{t+1} = F_t(X_t, \varepsilon_t)$ for $F_t(x, y) = f_t(x, u_t(x), y)$. But a similar structure is also valid, if e.g. $U_t = u_t(X_t, X_{t-1})$ by suitable rewriting. Indeed, let $\mathbf{X}_t = (X_t, X_{t-1})$, $x = (x_1, x_2)$. Then we obtain $\mathbf{X}_{t+1} = \mathbf{F}_t(\mathbf{X}_t, \varepsilon_t)$, where $\mathbf{F}_t(x, y) = (f_t(x_1, u_t(x_1, x_2), y), x_1)$. This shows that the setup of (9.1) and (9.2) is more general than it appears, and it includes k -step Markov processes as well.

Our aim is to solve the following problem.

Problem 9.3 Let $g_0, \dots, g_{T-1} : \mathbb{R}^d \times \mathbb{R}^m \rightarrow \mathbb{R}$ be measurable functions, as well as $g_T : \mathbb{R}^d \rightarrow \mathbb{R}$. The problem is to maximize over $U = (U_0, \dots, U_{T-1})$ the expectation

$$J(U) := \mathbb{E} \left(\sum_{t=0}^{T-1} g_t(X_t, U_t) + g_T(X_T) \right),$$

with each U_t as in (9.2). This problem is thus equivalent to the finding of measurable functions u_t such that the constraint (9.2) holds. With $u = (u_0, \dots, u_{T-1})$ we also write $J(u)$ instead of $J(U)$ to emphasize that J depends on the *functions* u_t . Usually, the functions have to satisfy certain constraints. These will be clear in the appropriate context and not always explicitly mentioned. For instance, it is tacitly assumed that all random quantities involved are such that the expectations exist.

Definition 9.4 A sequence of functions $u^* = (u_0^*, \dots, u_{T-1}^*)$ is called optimal if $J(u^*) = \sup J(u)$ holds.

Definition 9.5 Let u be a sequence of functions (u_0, \dots, u_{T-1}) and let the process X^u be defined by (9.1) and (9.2), the notation expresses the dependence of X on u . Note that $X_0^u = X_0$. Then, we define $J_T(u) = g_T(X_T^u)$, and for $t < T$,

$$\begin{aligned} J_t(u) &:= \mathbb{E} \left[\sum_{s=t}^{T-1} g_s(X_s^u, U_s) + g_T(X_T^u) \middle| \mathcal{F}_t \right] \\ &= g_t(X_t^u, U_t) + \mathbb{E} [J_{t+1} | \mathcal{F}_t]. \end{aligned}$$

Note that $\mathbb{E} J_0(u) = J(u)$. The J_t can be interpreted as expected future rewards, given the past up to time t .

Define also for certain given measurable functions $v_0, \dots, v_T : \mathbb{R}^d \rightarrow \mathbb{R}$

$$(9.3) \quad \hat{v}_{t+1}(x, y) = \mathbb{E} v_{t+1}(f_t(x, y, \varepsilon_t)), \quad t = 0, \dots, T-1,$$

so $\hat{v}_{t+1} : \mathbb{R}^d \times \mathbb{R}^m \rightarrow \mathbb{R}$. Note that these functions are measurable in x and y .

Important is the situation in which all the U_t are such that $U_t = u_t(X_t)$, for some measurable functions $u_t : \mathbb{R}^d \rightarrow \mathbb{R}^m$. Denote the class of sequences $u = (u_0, \dots, u_{T-1})$ of such functions by \mathcal{M} . The main theorem of this section is

Theorem 9.6 Define recursively the functions v_t , $t = 0, \dots, T$, by

$$(9.4) \quad \begin{aligned} v_T(x) &= g_T(x) \\ v_t(x) &= \sup_y \{g_t(x, y) + \hat{v}_{t+1}(x, y)\}, \quad t = 0, \dots, T-1. \end{aligned}$$

Assume that the v_t are measurable functions. Then the following hold true.

(i) For any sequence $u \in \mathcal{M}$ of functions u_t one has

$$v_t(X_t^u) \geq J_t(u) \text{ a.s.}$$

and $\mathbb{E} v_0(X_0) \geq J(u)$.

(ii) Let $u^* \in \mathcal{M}$. Then u^* is optimal iff the supremum in (9.4) is attained for $y = u_t^*(x)$. If this happens, then $v_t(X_t^{u^*}) = J_t(u^*)$ and $\sup_u J(u) = J(u^*) = \mathbb{E} v_0(X_0)$.

Proof See Section A.5. □

The condition in Theorem 9.6 that the functions v_t are measurable, is satisfied under the additional assumption that both $g_t(x, \cdot)$ and $\hat{v}_{t+1}(x, \cdot)$ are continuous for all t and x . Theorem 9.6 also provides an algorithm that yields the optimal functions u_t^* .

Algorithm 9.7 (Dynamic programming) Suppose that the suprema in Equation (9.4) are attained for all t . Define

$$\begin{aligned} v_T(x) &= g_T(x) \\ u_{T-1}^*(x) &= \arg \sup_y \{g_{T-1}(x, y) + \hat{v}_T(x, y)\}, \end{aligned}$$

and by backwards recursion for $t \in \{0, \dots, T-1\}$

$$\begin{aligned} v_t(x) &= \sup_y \{g_t(x, y) + \hat{v}_{t+1}(x, y)\} \\ &= g_t(x, u_t^*(x)) + \hat{v}_{t+1}(x, u_t^*(x)) \\ u_{t-1}^*(x) &= \arg \sup_y \{g_{t-1}(x, y) + \hat{v}_t(x, y)\}. \end{aligned}$$

This yields the sequence of functions $v_T, u_{T-1}^*, v_{T-1}, u_{T-2}^*, \dots, v_1, u_0^*$ in which the u_t^* constitute the optimal sequence u^* . The functions v_t are called the *(optimal) value functions*.

Proposition 9.8 (Optimality principle) Let $u^* = (u_0^*, \dots, u_{T-1}^*)$ be the optimal sequence for Problem 9.3 as obtained from Algorithm 9.7. Then the sequence $(u_t^*, \dots, u_{T-1}^*)$ is optimal for the corresponding optimization problem over the time set $\{t, \dots, T\}$, when starting in $X_t = X_t^{u^*}$. In this case the optimal value is equal to $\mathbb{E} v_t(X_t^{u^*})$.

Proof Exercise 9.2. □

Remark 9.9 The results above strongly depend on the fact that $\hat{v}_{t+1}(x, y)$ is equal to the conditional expectation $\mathbb{E}[f_t(x, y, \varepsilon_t) | \mathcal{F}_t]$, which follows from the assumed independence of X_0 and the ε_t , see Lemma A.14. In Section 9.2 however, we will come across situations, where this assumption is often violated. We proceed with giving some results for a more general setting.

From here on, we drop the assumption that the ε_t are independent. One can still define ‘functions’ \hat{v}_{t+1} , but now we alter the definition of (9.3) into

$$\hat{v}_{t+1}(x, y) = \mathbb{E}[v_{t+1}(f_t(x, y, \varepsilon_t)) | \mathcal{F}_t].$$

Then the $\hat{v}_{t+1}(x, y)$ are in general not deterministic anymore, but become \mathcal{F}_t -measurable random variables. However, many of the above results continue to hold. For instance, the technical result (A.7) is still correct. On the other hand, we need an alternative to Lemma A.16, used in the proof of Theorem 9.6, and the theorem itself. The following proposition uses the concept of *essential supremum*, see Section A.3. We will consider $\mathcal{B}(\mathbb{R}^d) \times \mathcal{F}_t$ -measurable functions v_t , meaning that the mapping $(x, \omega) \mapsto v_t(x, \omega)$ is $\mathcal{B}(\mathbb{R}^d) \times \mathcal{F}_t$ -measurable. As usual, dependence on ω is often suppressed and then we write $v_t(x)$ for the random variable $\omega \mapsto v_t(x, \omega)$.

Proposition 9.10 Suppose that $\mathcal{B}(\mathbb{R}^d) \times \mathcal{F}_t$ -measurable functions v_t (for $t = 0, \dots, T$) a.s. satisfy

$$\begin{aligned} v_T(x) &= g_T(x) \\ v_t(x) &= \text{ess sup}\{g_t(x, y) + \mathbb{E}[v_{t+1}(f_t(x, y, \varepsilon_t)) | \mathcal{F}_t] : y \in \mathcal{F}_t\}, t \leq T - 1. \end{aligned}$$

Then, for any sequence $u = (u_0, \dots, u_{T-1})$ it holds that $v_t(X_t^u) \geq J_t(u)$ for $t = 0, \dots, T$. Optimality is obtained for the random variables $u_t^*(x)$ for which the essential supremum is attained.

Proof Essentially as the proofs of Lemma A.16 and Theorem 9.6. □

There is a major difference between the above proposition and previous results. The optimizing $u_t^*(x)$ are now \mathcal{F}_t -measurable random variables, and to emphasize this, we should write $u_t^*(x) = u_t^*(x, \omega)$. The dependence on ω , is usually through X_0, \dots, X_t . Writing $u_t^*(X_t^{u^*})$ suggests explicit dependence of an optimal control through $X_t^{u^*}$ only, but this is in general not the case. Nevertheless, in many practical situations it is still possible to explicitly compute the optimal strategy, see for instance Exercise 9.4 for a problem in a financial context.

9.2 Optimal portfolios via dynamic programming

We apply the general results of the previous section to an expected utility maximization problem. We assume that the assumptions of Section 6 are in force. An investor has an initial capital w at his disposal. He can invest in shares in a market described by the model of Section 8. Without further explanation, below we use the notation of that section. The first problem we consider is the problem of maximizing the expected utility of terminal wealth, so we want to maximize

$$\mathbb{E} \tilde{u}(W_T),$$

subject to the budget constraint $W_0 \leq w$ and to the constraint that W_T results from a self-financing trading strategy, so $W_T = \bar{\xi}_T \cdot \bar{S}_T$. Writing $W_T = S_T^0 V_T$ and $V_T = V_0 + G_T$, we see that we have to maximize

$$\mathbb{E} \tilde{u}(S_T^0(w + G_T)),$$

where of course we have taken $V_0 = W_0 = w$, since again it can never be optimal to use only a fraction of the initial capital w . Let us assume that the process S^0 is deterministic. Then we can define a new utility function u by putting $u(x) = \tilde{u}(S_T^0 x)$. Hence we have to find

$$\max \mathbb{E} u(w + G_T),$$

where $G_T = \sum_{t=1}^T \xi_t \cdot \Delta X_t$ and the ξ_t are \mathcal{F}_{t-1} -measurable. Notice that the ξ_t will be our (random) decision variables. In principle we'd like to apply the dynamic programming algorithm 9.7, with the proper substitutions and change of notation. For instance, we have that the functions g_t are zero for $t \leq T - 1$

and $X_{t+1} = f_t(X_t, u_t, \varepsilon_t)$ now reads $V_{t+1} = V_t + \xi_{t+1} \cdot \Delta X_{t+1}$. Note however that the ΔX_t are in general not independent, which spoils the fact that the optimal decisions at time t not only depend on V_t , but also on past values. Therefore the version of the dynamic programming algorithm that we need below is taken from Proposition 9.10. Define

$$\tilde{v}_T(x) = \tilde{u}(x)$$

and then, recursively, for $t \in \{0, \dots, T-1\}$

$$(9.5) \quad \tilde{v}_t(x) = \text{ess sup} \{ \mathbb{E} [\tilde{v}_{t+1}(S_{t+1}^0 (\frac{x}{S_t^0} + \xi \cdot \Delta X_{t+1})) | \mathcal{F}_t] : \xi \in \mathcal{F}_t \}.$$

Assume that for every t the essential supremum is attained at some $\xi_{t+1}^* = \xi_{t+1}^*(x) \in \mathcal{F}_t$. This eventually gives rise to a self-financing strategy $\bar{\xi}$ by the by now familiar choices for the ξ_t^0 , the investments in the non-risky asset. At the final step of the algorithm ($t = 0$), we find ξ_0 and Theorem 9.6, or rather Proposition 9.10, tells us that $\mathbb{E} \tilde{v}_0(w)$ is the optimal value for our problem. Conditions for which the suprema are attained, for instance at an interior point of the domains of the \tilde{v}_t , can be derived from Theorem 6.6, although this theorem in general has to be applied ‘ ω -wise’, which involves some subtleties.

In terms of the modified utility functions u , we can recast the optimal value functions resulting from Dynamic Programming as

$$(9.6) \quad v_t(x') = \text{ess sup} \{ \mathbb{E} [v_{t+1}(x' + \xi \cdot \Delta X_{t+1}) | \mathcal{F}_t] : \xi \in \mathcal{F}_t \}.$$

One can show that this leads to the same optimum, if it exists (Exercise 9.5).

We generalize the above problem as to include *consumption*.

Definition 9.11 A consumption process $C = (C_0, \dots, C_T)$ is a nonnegative adapted process. A consumption-investment plan is a pair $(C, \bar{\xi})$, where C is a consumption process and $\bar{\xi}$ a trading strategy. Such a plan is called *self-financing*, if

$$W_t = C_t + \bar{\xi}_{t+1} \cdot \bar{S}_t, \quad t \in \{0, \dots, T-1\}.$$

where W_t is as before, $W_t = \bar{\xi}_t \cdot \bar{S}_t$. It is called *admissible* if $C_T \leq W_T$ a.s.

Remark 9.12 The self-financing condition above is equivalent to the following two relations.

$$\begin{aligned} \Delta W_t &= \bar{\xi}_t \cdot \Delta \bar{S}_t - C_{t-1}, \quad t \in \{1, \dots, T\} \\ \Delta V_t &= \xi_t \cdot \Delta X_t - \gamma_{t-1}, \quad t \in \{1, \dots, T\}, \end{aligned}$$

where γ_{t-1} is discounted consumption, $\gamma_{t-1} = \frac{C_{t-1}}{S_{t-1}^0}$.

The problem we are going to address is the maximization of

$$\mathbb{E} \sum_{t=1}^T \alpha^t u(C_t),$$

where $\alpha \in (0, 1]$ (a discount factor) and u a utility function, subject to the constraints that C forms together with a trading strategy $\bar{\xi}$ an admissible consumption-investment plan. Note that, for simplicity, we use a single utility function at all times t , although it can be made time dependent too.

It is immediately clear that $C_T = W_T$, since u is increasing. This makes the first step in the dynamic programming approach easy, $v_T = u$. In order to motivate the resulting backward recursion, we now consider the problem at time $T - 1$, assuming that we have (optimally) invested and consumed up to that time. This can be viewed as a one period problem. We have to maximize

$$(9.7) \quad u(C_{T-1}) + \alpha \mathbb{E}[v_T(W_T) | \mathcal{F}_{T-1}]$$

subject to constraints that we now derive. Let w denote the wealth at time $T - 1$. By the self-financing condition we have

$$w = C_{T-1} + \bar{\xi}_T \cdot \bar{S}_{T-1},$$

whereas

$$W_T = \bar{\xi}_T \cdot \bar{S}_T.$$

To assure that the (optimal) consumption process is non-negative, we assume *without loss of generality* that $u(x) = -\infty$ for $x < 0$. Using the self-financing characterization as in Remark 9.12, we can rewrite (9.7) with $w' = \frac{w}{\bar{S}_{T-1}^0}$ as

$$u(C_{T-1}) + \alpha \mathbb{E}[v_T(S_T^0(w' - \gamma_{T-1} + \xi_T \cdot \Delta X_T)) | \mathcal{F}_{T-1}],$$

which we have to maximize over $\xi_T \in \mathcal{F}_{T-1}$. Of course we can iterate this procedure to get the dynamic programming equation for every $t \in \{1, \dots, T\}$

$$(9.8) \quad v_{t-1}(w') = \text{ess sup}_{\xi_t, \gamma_{t-1} \in \mathcal{F}_{t-1}} \{u(C_{t-1}) + \alpha \mathbb{E}[v_t(S_t^0(w' - \gamma_{t-1} + \xi_t \cdot \Delta X_t)) | \mathcal{F}_{t-1}]\} :$$

At the final step, one obtains $v_0(w') = v_0(w)$, which will then yields the optimal value $\mathbb{E} v_0(w)$. As before, the optimal processes C^* and ξ^* –assuming that they exist– have to be complemented by the process ξ^0 to obey the self-financing restriction.

9.3 Consumption-investment and the martingale method

We start out with the standing assumption that *the market is complete*. For some of the problems that we introduce later on, this assumption is not always needed. This will be explained, when those are treated.

To illustrate the underlying principles, we first recall a static problem, which is the maximization of $\mathbb{E} u(W_1)$. Here W_1 represents the value of a portfolio $\bar{\xi}$ at $t = 1$. Hence the maximization takes place over all $\bar{\xi} \in \mathbb{R}^{d+1}$, that of course have to satisfy the budget constraint $\bar{\xi} \cdot \bar{\pi} = w$, where w is the initially available capital.

The central idea behind the martingale method is to break down the optimization problem into two subproblems. The first one is to identify the optimal (random) pay-off W_1^* , given the budget constraint. The second one is then to identify the optimal portfolio ξ , i.e. the portfolio whose terminal wealth is equal to W_1^* . For simplicity, we assume for the time being that *the price of the riskless asset is constant and equal to one*. Let \mathbb{P}^* be the unique risk-neutral measure (of course we assume that the market is arbitrage-free, see Theorem 6.5, so it exists). Then the budget constraint is given by $\mathbb{E}^*W_1 = w$. Let \mathcal{B} be the set of random variables that are integrable w.r.t. \mathbb{P}^* with corresponding expectation equal to w . Hence, the first problem becomes the maximization of $\mathbb{E}u(W)$, subject to $W \in \mathcal{B}$, where we write W instead of W_1 . Let $\phi = \frac{d\mathbb{P}^*}{d\mathbb{P}}$.

One way to solve this problem is to employ a Lagrange multiplier (similar, but alternative to what we did in Section 7.1). So one likes to maximize

$$L(W, \lambda) = \mathbb{E}u(W) - \lambda(\mathbb{E}(\phi W) - w).$$

Differentiation of L w.r.t. λ yields the budget constraint. Next, one would like to differentiate w.r.t. W , which is a priori an infinite dimensional variable. This would lead to some *variational problem*. To circumvent this we think for a while that the underlying Ω is finite with positive probabilities of all singletons. Then we can represent W by a finite dimensional vector with a generic element $w_j = W(\omega_j)$. Assume that u is differentiable. Differentiation w.r.t. w_j of L can now be carried out under the expectation and yields

$$u'(W(\omega_j)) - \lambda\phi(\omega_j) = 0.$$

Since this equation has to hold for every ω_j , it follows that

$$(9.9) \quad \lambda = \mathbb{E}u'(W).$$

Compare this and the rest of this paragraph with Section 7.1. Let I denote the inverse of u' , which is assumed to exist (otherwise, one should work with the function I^+ as in Section 7.1). Then

$$W(\omega_j) = I(\lambda\phi(\omega_j)),$$

or, in short,

$$(9.10) \quad W^* = I(\lambda\phi).$$

Dropping the assumption that Ω is finite, we conjecture that Equations (9.9) with $W = W^*$ and (9.10) are needed to obtain the optimal claim. Compare to Theorem 7.2. The budget restriction tells us that also for the optimal W^* it must hold that $\mathbb{E}^*W^* = w$, so

$$\mathbb{E}^*I(\lambda\phi) = w,$$

which in principle yields λ , see Corollary 7.5.

Knowing the optimal contingent claim, one then has to find a corresponding hedge strategy (which exists, since we assumed that the market is complete), that is find $\bar{\xi}^*$ such that $W^* = \bar{\xi}^* \cdot \bar{S}$ a.s. The resulting $\bar{\xi}^*$ should of course coincide with the solution of Theorem 6.6.

The above approach extends to the analysis of a consumption-investment problem with a time horizon $T > 1$. As we shall see, here we don't need market completeness, but of course we can't dispense with the requirement that the market is free of arbitrage. For a $T = 1$ horizon this problem reduces to the following. First we pin down the admissible consumption-investment plan, see Definition 9.11 and recall that W_t is the notation for the undiscounted value of a portfolio at time t , which, under the assumption that $S_t^0 \equiv 1$, is equal to V_t . It is such that $C_0 + W_0 = w$ and $C_1 = W_1$. Notice that it follows that C_1 is an attainable claim! Therefore, for any risk-neutral measure \mathbb{P}^* it holds that $\mathbb{E}^* C_1 = W_0$ and we get $\mathbb{E}^* C_1 + C_0 = w$. But there is also a converse reasoning. If a consumption plan C is fixed, as well as an initial capital w , then a consumption-investment plan $(C, \bar{\xi})$ is admissible, if $\mathbb{E}^* C_1 + C_0 = w$ for any risk-neutral measure. Indeed, it now follows from Proposition 1.18 that C_1 is attainable, hence $C_1 = \bar{\xi} \cdot \bar{S}_1$, for some $\bar{\xi}$ and hence also $C_1 = W_1$.

A consumption-investment optimization problem is usually formulated as the maximization of

$$u(C_0) + \alpha \mathbb{E} u(C_1),$$

subject to the constraints that $C_0, C_1 \geq 0$ a.s. and $C_0 + \mathbb{E}^* C_1 = w$. Notice that the utility of C_0 and C_1 is represented by the same utility function u , but of course different choices for each of them are equally conceivable.

We adopt again the Lagrange multiplier approach to solve this problem. So we want to maximize

$$L(C_0, C_1, \lambda) = u(C_0) + \alpha \mathbb{E} u(C_1) - \lambda(C_0 + \mathbb{E}(\phi C_1) - w).$$

For the optimal consumption pair C_0^*, C_1^* we then get, by the same token as we used before and assuming that $I = (u')^{-1}$ is well defined on $(0, \infty)$,

$$(9.11) \quad C_0^* = I(\lambda)$$

$$(9.12) \quad C_1^* = I(\lambda \phi / \alpha),$$

whereas the optimal λ^* has to solve the equation $I(\lambda) + \mathbb{E}^* I(\lambda \phi / \alpha) = w$.

After this exposition of the problem in a static setting, we turn to the dynamic case and also take discounting into account. First we consider the problem of determining the optimal final wealth, resulting from investments only. As in the static case, we first determine an \mathcal{F}_T -measurable random variable W^* that is such that $\mathbb{E} u(W)$ is maximal, subject to the constraint that $\mathbb{E}^* V = w$, where $V = W/S_T^0$ and w is an initially available capital. Mimicking the static case, but taking care of discounting, we get the following results

$$W^* = I(\lambda \phi / S_T^0)$$

$$w = \mathbb{E}^* I(\lambda\phi/S_T^0)/S_T^0,$$

with ϕ equal to the Radon-Nikodym derivative $\frac{d\mathbb{P}^*}{d\mathbb{P}}$, where the probability measures are now defined on \mathcal{F}_T .

Since the market is assumed to be complete, we can in principle find a replicating strategy that yields W^* as its terminal value. Of course, this is a non-trivial exercise. But, as we shall see below, when we move to consumption-investment problems this problem is tractable. What one in principle has to do is to find $\bar{\xi}_T$ such that $\bar{\xi}_T \in \mathcal{F}_{T-1}$ and $\bar{\xi}_T \cdot \bar{S}_T = W^*$. As soon as this has happened, we know by the fact that $\bar{\xi}$ is a self-financing strategy, that $\bar{\xi}_T \cdot \bar{S}_{T-1} = \bar{\xi}_{T-1} \cdot \bar{S}_{T-1}$, from which one has to determine $\bar{\xi}_{T-1} \in \mathcal{F}_{T-2}$. Notice that $\mathbb{E}^* \bar{\xi}_{T-1} \cdot \bar{S}_{T-1} / S_{T-1}^0 = \mathbb{E}^* W^* / S_T^0$, for any underlying \mathbb{P}^* . See Exercise 9.7 for an example that shows how to carry out this programme in a concrete situation.

We turn to consumption-investment problems. We will see that completeness of the market is not always necessary, but of course the market has to be arbitrage-free. So risk-neutral measures \mathbb{P}^* exist.

Definition 9.13 A consumption process is called *attainable*, if there exists a trading strategy $\bar{\xi}$ such that $(C, \bar{\xi})$ is admissible and satisfies $C_T = W_T$. It is then said that $\bar{\xi}$ *replicates* C (at time T).

Let us first characterize attainable consumption processes. Recall Remark 9.12. Since the gains process G , $G_t = \sum_{s=1}^t \xi_s \cdot \Delta X_s$, is a martingale with expectation zero under any risk-neutral measure, it follows that for each $t \leq T$ one has

$$\mathbb{E}^* V_t + \sum_{s=0}^{t-1} \mathbb{E}^* \gamma_s = w.$$

If the consumption process is attainable, $\gamma_T = V_T$, we get

$$(9.13) \quad \sum_{t=0}^T \mathbb{E}^* \gamma_t = w, \text{ for any } \mathbb{P}^* \in \mathcal{P}.$$

Proposition 9.14 Assume a complete market. Given any initial wealth w , a consumption process is attainable iff (9.13) holds. In this case the initial value of the replicating portfolio is nonnegative.

Proof Necessity has already been proved. We turn to sufficiency. Any of the γ_t in (9.13) can be considered as a discounted contingent claim, since they are all nonnegative and \mathcal{F}_t -measurable; the payoff of these claims take place at the final time T . So, for each of them, there exists a replicating self-financing strategy $\bar{\xi}^{(t)}$. Take $\bar{\xi} = \sum_{t=0}^T \bar{\xi}^{(t)}$. Using Theorem 8.30, one can verify that $(C, \bar{\xi})$ is an admissible self-financing consumption-investment plan with $C_T = W_T$.

The martingale property of the gains process yields

$$\mathbb{E}^* [\Delta V_s | \mathcal{F}_{s-1}] + \gamma_{s-1} = 0.$$

Summing this equation for $s = t + 1, \dots, T$ and taking conditional expectation given \mathcal{F}_t yields

$$\mathbb{E}^*[V_T - V_t + \sum_{s=t}^{T-1} \gamma_s | \mathcal{F}_t] = 0.$$

Together with $V_T = \gamma_T$ this gives

$$V_t = \mathbb{E}^*[\sum_{s=t}^T \gamma_s | \mathcal{F}_t] \geq 0, \text{ a.s.}$$

In particular, $V_0 \geq 0$. □

After this intermediate result we are now in the position to state the optimization problem properly. It is the maximization of

$$\mathbb{E}[\sum_{t=0}^T \alpha^t u(C_t)],$$

subject to the constraint that C is a nonnegative adapted process and the budget constraint (9.13). We will also assume that $u(x) = -\infty$ for $x < 0$ and $u'(0) = \infty$. These conditions are sufficient to obtain an a.s. strictly positive optimal consumption process. In view of Proposition 9.14, the resulting optimal consumption process C^* will be attainable. Once we have found this, we have to find the replicating strategy.

Let us focus on the finding of C^* . We will use the following

Lemma 9.15 *Let $\mathbb{P}^* \in \mathcal{P}$ and $\phi = \frac{d\mathbb{P}^*}{d\mathbb{P}}$, the Radon-Nikodym derivative on \mathcal{F}_T , and $M_t = \mathbb{E}[\phi | \mathcal{F}_t]$, $t = 0, \dots, T$. Then it holds that*

$$\mathbb{E}^* \sum_{t=0}^T \gamma_t = \mathbb{E} \sum_{t=0}^T \gamma_t M_t.$$

Proof This follows from

$$\begin{aligned} \mathbb{E}^* \gamma_t &= \mathbb{E} \phi \gamma_t \\ &= \mathbb{E} (\gamma_t \mathbb{E}[\phi | \mathcal{F}_t]) \\ &= \mathbb{E} \gamma_t M_t, \end{aligned}$$

valid for all $t \in \{0, \dots, T\}$. □

From this lemma it follows that we can replace the constraint (9.13) with

$$\mathbb{E} \sum_{t=0}^T \gamma_t M_t = w.$$

We will solve the optimization problem by again using a Lagrange multiplier. So, we want to maximize

$$(9.14) \quad L(C_0, \dots, C_T, \lambda) = \mathbb{E} \sum_{t=0}^T \alpha^t u(C_t) - \lambda \mathbb{E} \left(\sum_{t=0}^T \gamma_t M_t - w \right).$$

The necessary conditions for a maximum become

$$\alpha^t u'(C_t) - \lambda N_t = 0, \quad t \in \{0, \dots, T\},$$

where $N_t = \frac{M_t}{S_t}$. Then, if $I = (u')^{-1}$ is well-defined everywhere, we obtain the optimal

$$C_t^* = I(\lambda \alpha^{-t} N_t), \quad t \in \{0, \dots, T\}.$$

Of course the optimal λ^* has to satisfy

$$\mathbb{E} \sum_{t=0}^T I(\lambda^* \alpha^{-t} N_t) N_t = w.$$

Here is an example that illustrates the above procedure.

Example 9.16 Let $u(x) = \log x$, $x > 0$. Then we have $I(x) = \frac{1}{x}$. We obtain

$$C_t^* = \frac{\alpha^t}{\lambda N_t},$$

and

$$w = \mathbb{E} \sum_{t=0}^T \frac{\alpha^t}{\lambda N_t} N_t,$$

from which it follows that

$$\lambda = \begin{cases} \frac{T+1}{w} & \text{if } \alpha = 1 \\ \frac{1-\alpha^{T+1}}{w(1-\alpha)} & \text{if } \alpha < 1. \end{cases}$$

In the first of these two cases, one can compute that $C_t^* = \frac{w}{(T+1)N_t}$ and $\gamma_t = \frac{w}{(T+1)M_t}$. The maximal value of the objective function becomes $(T+1) \log \frac{w}{T+1} - \sum_{t=0}^T \mathbb{E} \log N_t$.

We continue to study an optimization problem that combines the previous two, we want to maximize expected utility derived from both consumption and terminal wealth. The chief difference with the previous problem is that we don't require $W_T = C_T$ anymore. Therefore, we have to assume again that the market is complete. We denote by \mathcal{A}_w the set of all admissible consumption-investment plans that have w as initial wealth and that satisfy the terminal condition $C_T \leq W_T$. We will assume that two utility functions u_1 and u_2 (with $u_i \in C^1(0, \infty)$) are involved. The function u_1 describes the utility directly derived from consumption and u_2 the utility derived from terminal wealth as

well. Again, we assume that the u_i can be extended to the whole of \mathbb{R} by setting $u_i(x) = -\infty$ for $x < 0$, u right-continuous at $x = 0$ and moreover $\lim_{x \downarrow 0} u'_i(x) = \infty$. The aim is to maximize for $(C, \xi) \in \mathcal{A}_w$ the cumulative expected utility

$$(9.15) \quad \mathbb{E} \left[\sum_{t=0}^T \alpha^t u_1(C_t) + \alpha^T u_2(W_T - C_T) \right].$$

There is a variation on this problem conceivable, for instance by replacing the last utility term by $u_2(W_T)$ and/or having in the summation an upper limit equal to $T - 1$. We leave this possibility aside.

The dynamic programming approach to this problem results in a recursion formula that bears some obvious resemblance to (9.8),

$$v_{t-1}(w) = \text{ess sup} \{ u_1(C) + \alpha \mathbb{E} [u_2 \left((w - C) \frac{S_t^0}{S_{t-1}^0} + S_t^0 \xi \cdot \Delta X_t \right) | \mathcal{F}_{t-1}] : C, \xi \in \mathcal{F}_{t-1} \}.$$

There is a crucial difference however with the initialization of the dynamic programming algorithm. At time T the final utility is $\alpha^T (u_1(C_T) + u_2(W_T - C_T))$ and so one has to divide terminal wealth into what is kept (for future investments for instance) and what is consumed. Therefore the proper initialization becomes

$$v_T(w) = \max \{ u_1(c) + u_2(w - c) : 0 \leq c \leq w \}.$$

In the remainder of this section we focus on the risk-neutral approach. Paralleling the reasoning that led us to Proposition 9.14, we obtain

Proposition 9.17 *Given an initial wealth $w \geq 0$ and an admissible consumption-investment plan $(C, \bar{\xi})$, it holds that*

$$(9.16) \quad \mathbb{E}^* \left[\sum_{t=0}^{T-1} \gamma_t + V_T \right] = w.$$

Conversely, given $w \geq 0$ and a consumption process C with $C_T \leq W_T$, there exists a trading strategy $\bar{\xi}$ such that $(C, \bar{\xi})$ is admissible if relation (9.16) holds.

Proof Similar to the proof of Proposition 9.14. \square

It follows that we can recast the optimization problem as the maximization of (9.15) subject the constraints $W_T \in \mathcal{F}_T$, C a (nonnegative) adapted process such that $C_T \leq W_T$ and (9.16). Actually the assumptions that $u'_i(x) \rightarrow \infty$ as $x \downarrow 0$ will guarantee that the optimal consumption process C^* is such that $C_t^* > 0$ a.s. for all t and that the optimal terminal wealth is such that $W_T^* > C_T^*$ a.s. Hence the constraint $C_T \leq W_T$ will be automatically satisfied in the optimum and is therefore redundant.

Recall the definition of the martingale M of Lemma 9.15, its ‘discounted’ analogue N and Equation (9.14). In the present situation we have to maximize the Lagrangian

$$L(C_0, \dots, C_T, W_T, \lambda) =$$

$$\mathbb{E} \left[\sum_{t=0}^T \alpha^t u_1(C_t) + \alpha^T u_2(W_T - C_T) - \lambda \left(\sum_{t=0}^{T-1} C_t N_t + W_T N_T \right) \right].$$

As before, one can write down the first order necessary conditions, by computing partial derivatives. Solving these equations and assuming that I_1 and I_2 are properly defined inverse functions of u'_1 and u'_2 respectively, we obtain

$$\begin{aligned} C_t^* &= I_1(\alpha^{-t} \lambda N_t), \quad t = 0, \dots, T \\ W_T^* &= I_1(\alpha^{-T} \lambda N_T) + I_2(\alpha^{-T} \lambda N_T). \end{aligned}$$

The optimal value λ^* follows by inserting the optimal solution into Equation (9.16), provided that the resulting equation has a unique solution. One can show that this is for instance the case if $u_1 = u_2$ and u_1 of HARA type.

9.4 Exercises

9.1 Prove Lemma 9.1.

9.2 Prove Proposition 9.8.

9.3 Consider the optimization problem at beginning of Section 9.2. Show that

$$\frac{d\mathbb{P}^*}{d\mathbb{P}} = \frac{S_T^0 \tilde{u}'(W_T)}{\mathbb{E} S_T^0 \tilde{u}'(W_T)}$$

defines a risk-neutral measure.

9.4 Consider a market with underlying $\Omega = \{1, 2, 3, 4\}$. Assume that $T = 2$ and that $S_t^0 = 1$ for $t = 0, 1, 2$, the price of the riskless asset is constant and equal to one. Let the evolution for the price S_t of the *single* risky asset be as given in the table. We further assume that all singletons have probability $\frac{1}{4}$.

ω	$S_0(\omega)$	$S_1(\omega)$	$S_2(\omega)$
1	5	8	9
2	5	8	6
3	5	4	6
4	5	4	3

The aim is to maximize $\mathbb{E} u(W_T)$, as in Section 9.2, for $u(x) = 1 - \exp(-x)$.

- Let $t = 2$. Show that the optimal ξ_2^* satisfies $\xi_2^*(\omega) = -\frac{1}{3} \log 2$, if $\omega \in \{1, 2\}$ and that $\xi_2^*(\omega) = \frac{1}{3} \log 2$, if $\omega \in \{3, 4\}$.
- Compute the optimal ξ_1^* (a constant!).
- What is the resulting optimal expected utility?

9.5 Show that the recursions (9.5) and (9.6) lead to the same optimum at $t = 0$.

9.6 Consider a CRR model, in which the returns are *iid* with $\mathbb{P}(R_t = b) = p$ (where p is not necessarily equal to the risk neutral value p^*). Consider the maximization of $\mathbb{E}u(W_T)$, with $u(x) = \log x$. Compute the optimal trading strategy ξ^* via dynamic programming. *Hint*: show that $\tilde{v}_t(x) = \log x + k_t$ for certain constants k_t .

9.7 Consider a CRR model as in Exercise 9.6, so with a parameter p that determines the probability measure \mathbb{P} . Let p^* be as in Proposition 8.35. Assume that $u(x) = \log x$ and that the initial capital is w .

(a) Show that the optimal attainable terminal wealth is given by

$$W_T^* = w(1+r)^T \left(\frac{p}{p^*}\right)^{B_T} \left(\frac{1-p}{1-p^*}\right)^{T-B_T},$$

where B_T is the number of ‘up-movements’ of the stock.

(b) Assume that at time $T-1$ and that B_{T-1} ‘up-movements’ have been observed. Show, using the risk-neutral approach, that for the optimal replicating strategy one has

$$\begin{aligned} \xi_T^1 &= w(1+r)^T \left(\frac{p}{p^*}\right)^{B_{T-1}} \left(\frac{1-p}{1-p^*}\right)^{T-1-B_{T-1}} \frac{p-p^*}{S_{T-1}(b-a)p^*(1-p^*)} \\ \xi_T^0 &= w \left(\frac{p}{p^*}\right)^{B_{T-1}} \left(\frac{1-p}{1-p^*}\right)^{T-1-B_{T-1}} \left(\frac{p^*-p+bp^*(1-p)-ap(1-p^*)}{(b-a)p^*(1-p^*)}\right). \end{aligned}$$

(c) Show that $W_{T-1}^* = w(1+r)^{T-1} \left(\frac{p}{p^*}\right)^{B_{T-1}} \left(\frac{1-p}{1-p^*}\right)^{T-1-B_{T-1}}$ and that the fraction of the wealth W_{T-1}^* that is invested in the risky asset is equal to

$$\frac{(1+r)(p-p^*)}{(b-a)p^*(1-p^*)}.$$

(d) Conjecture what the fraction of the capital W_t^* is, that is invested in the risky asset at $t < T-1$.

9.8 Suppose that at each time $t \in \{0, \dots, T\}$ an investor has a certain capital W_t at his disposal. He consumes part of this, C_t say, and invests the remaining $W_t - C_t$. He does this partly, a deterministic fraction π_t in a riskless asset with fixed return r and the remaining money in a risky asset with random yield S_t . Let u be a utility function and $\rho \in (0, 1)$ a discount factor. The aim is to maximize $\sum_{t=0}^T \rho^t \mathbb{E}u(C_t)$. Assume $C_T = W_T$. Characterize the optimal consumption pattern for $u(x) = x^\gamma$, where $x \geq 0$ and $\gamma \in (0, 1)$. Show in particular that π_t is the same for all $t \leq T-1$ and that there exist constants $\alpha_t \in (0, 1)$ such that $C_t = \alpha_t W_t$.

9.9 Show that Equation (9.16) has a unique solution if $u_1 = u_2$ and of HARA type.

9.10 Verify the validity of Equations (9.11) and (9.12) by the variational arguments of Section 7.1. You are allowed to interchange expectation and differentiation when needed.

A Appendix

A.1 Separating hyperplanes

Theorem A.1 *Let C be a non-empty convex subset in \mathbb{R}^n such that $0 \in \mathbb{R}^n \setminus C$. Then there exists a linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}$ such that $Tx \geq 0$ for all x in C and $Tx_0 > 0$ for at least one $x_0 \in C$.*

Proof Assume first that $0 \notin \text{Cl } C$. Consider the continuous map $x \rightarrow \|x\|$ and let B_n be the closed ball around zero with radius n , $n \in \mathbb{N}$. Restricted to $C_n := B_n \cap \text{Cl } C$, this map attains a minimum d_n at some x_n if the intersection is not empty. Obviously the d_n are decreasing, since $C_n \subset C_{n+1}$. But, if $x \in B_{n+1} \setminus B_n$, then $\|x\| > \|x'\|$ for all $x' \in B_n$, hence the d_n are all equal to some d . Let then $x_0 \in \text{Cl } C$ such that $\|x_0\| = d$, note that $x_0 \neq 0$ and $d > 0$. Define $Tx = x_0 \cdot X$. Then $Tx_0 = d^2 > 0$. Let $x \in C$ and let y be the projection on the subspace spanned by x_0 . Then $y = \lambda x_0$ and one easily shows, exploiting the convexity of C , that $\lambda \geq 1$, see Exercise A.1. But then $Tx = Ty = \lambda Tx_0 \geq \lambda d^2$. This shows the assertion under the extra assumption $0 \notin \text{Cl } C$.

To show the assertion for the general case, we may now assume that $0 \in \partial C$. We show that $\mathbb{R}^n \setminus \text{Cl } C \neq \emptyset$. If C is contained in a linear subspace of \mathbb{R}^n with dimension less than n , the assertion is obvious. So we assume that the linear span of C is equal to \mathbb{R}^n and therefore there exists a basis of \mathbb{R}^n consisting of n linear independent vectors v_k in C . Let $z = -\sum_k v_k$ and suppose that $z \in \text{Cl } C$. Then there $z_m \in C$ such that $z_m \rightarrow z$ and in particular all their coordinates c_k^m w.r.t. this basis converge to -1 . Hence there is an certain index m_0 such that all $c_k^{m_0}$ are negative. Let $\alpha_k = \frac{-c_k^{m_0}}{1 - \sum_k c_k^{m_0}}$ for $k = 1, \dots, n$ and $\alpha_0 = \frac{1}{1 - \sum_k c_k^{m_0}}$. Then 0 is the convex combination $0 = \alpha_0 z_{m_0} + \sum_{k=1}^n \alpha_k v_k$ and thus in C , which contradicts the hypothesis. We conclude that $\text{Cl } C$ is not all of \mathbb{R}^n .

We can now choose a sequence of z_n that all have strictly positive Euclidean distance to C and $z_n \rightarrow 0$. Application of the first part of the proof yields the existence of linear functionals T_n on \mathbb{R}^n such that $\inf\{T_n(x - z_n) : x \in C\} > 0$. We can represent T_n by unit vectors η_n , $T_n x = \eta_n \cdot x$. By compactness of the unit sphere, there exists a subsequence (η_{n_k}) converging to some limit vector η , with $\|\eta\| = 1$. But then for all $x \in C$ one has

$$\eta \cdot x = \lim \eta_{n_k} \cdot (x - z_{n_k}) \geq 0.$$

If $\eta \cdot x$ were zero for all $x \in C$, then $\eta \cdot x = 0$ for all $x \in \mathbb{R}^n$, since \mathbb{R}^n is the linear span of C . But this cannot happen, because $\eta \neq 0$. Hence there must be $x_0 \in C$ such that $\eta \cdot x_0 > 0$. \square

Remark A.2 Notice that the first part of the proof shows that $\inf\{Tx : x \in C\} > 0$ if $x \notin \text{Cl } C$.

A.2 Hahn-Banach theorem and some ramifications

Let \mathcal{X} be a (real) vector space with a norm $\|\cdot\|$. Let $T : \mathcal{X} \rightarrow \mathbb{R}$ be a linear operator. The operator norm of T is defined by $\|T\| := \sup\{|Tx| : \|x\| = 1\}$. We call T bounded if $\|T\| < \infty$. Recall that T is continuous iff it is bounded.

Definition A.3 Let \mathcal{X} be a (real) linear space. Call a function $p : \mathcal{X} \rightarrow \mathbb{R}$ a quasinorm, if p is sub-additive, $p(x + y) \leq p(x) + p(y)$ for $x, y \in \mathcal{X}$ and homogeneous, $p(tx) = tp(x)$, for $t \geq 0, x \in \mathcal{X}$.

A simple and useful example is given by $p(x) = c\|x\|$, if \mathcal{X} is endowed with a norm $\|\cdot\|$ and $c > 0$.

Theorem A.4 (Hahn-Banach) *Let T_0 be a linear operator defined on a linear subspace \mathcal{Y} of a real vector space \mathcal{X} and p a quasinorm on \mathcal{X} . Suppose that $|T_0y| \leq p(y)$ for all $y \in \mathcal{Y}$. Then T_0 admits an extension $T : \mathcal{X} \rightarrow \mathbb{R}$ such that $|Tx| \leq p(x)$ for all $x \in \mathcal{X}$. In particular, if T_0 is a bounded linear operator on a subspace of a normed space, then it extends to a bounded linear operator on \mathcal{X} such that $\|T\| = \|T_0\|$.*

Proof Assume that $\mathcal{Y} \neq \mathcal{X}$. Then there exists $x_1 \in \mathcal{X} \setminus \mathcal{Y}$. Let \mathcal{Y}_1 be the linear hull of $\{x_1\} \cup \mathcal{Y}$. Suppose that T_1 is an extension of T_0 to \mathcal{Y}_1 . Since every $x \in \mathcal{Y}_1$ can uniquely be written as $x = \alpha x_1 + y$, for some $y \in \mathcal{Y}$ and $\alpha \in \mathbb{R}$, it must hold that $T_1x = \alpha T_1x_1 + T_0y$. Write $\xi_1 := T_1x_1$, then we have

$$(A.1) \quad T_1x = \alpha\xi_1 + T_0y.$$

Conversely, every T_1 defined by (A.1) for some $\xi_1 \in \mathbb{R}$ is a continuation of T_0 to \mathcal{Y}_1 . We will show that it is possible to choose $\xi_1 \in \mathbb{R}$ such that $|T_1x| \leq p(x)$ for all $x \in \mathcal{Y}_1$. Thereto it is sufficient to prove that

$$(A.2) \quad T_1x \leq p(x), \forall x \in \mathcal{Y}_1.$$

From the proposed definition of T_1 with $\alpha = \pm 1$, it then follows from (A.2) that one necessarily has

$$(A.3) \quad \xi_1 \leq p(x_1 + y) - T_0y, \forall y \in \mathcal{Y}$$

$$(A.4) \quad \xi_1 \geq -p(x_1 + y) + T_0y, \forall y \in \mathcal{Y}.$$

But these two inequalities are also sufficient for (A.2) to hold. We now show that one can choose ξ_1 such that (A.3) and (A.4) hold true. Let y, y' be arbitrary elements of \mathcal{Y} . Then

$$T_0(y) - T_0(y') = T_0(y - y') \leq p(y + x_1 - (y' + x_1)) \leq p(y + x_1) + p(y' + x_1),$$

which yields

$$-p(y' + x_1) - T_0y' \leq p(y + x_1) - T_0y.$$

Taking the infimum over y and the supremum over y' , we get

$$\sup\{-p(y + x_1) - T_0y : y \in \mathcal{Y}\} \leq \inf\{p(x_1 + y) - T_0y : y \in \mathcal{Y}\}.$$

Hence a ξ_1 satisfying (A.3) and (A.4) exists.

We finish the proof by invoking Zorn's lemma. Consider the family of all extensions T of T_0 to linear subspaces of \mathcal{X} that contain \mathcal{Y} and that satisfy $|Tx| \leq p(x)$ for all x in the domain of T . This family can be endowed with the partial ordering defined by $T_1 \preceq T_2$ iff T_2 is an extension of T_1 . Then there exists a maximal element, T say, in this family w.r.t. this partial ordering. By the preceding part of the proof, the domain of T is all of \mathcal{X} . Indeed, if this were not the case, then we could take \mathcal{Y} in the previous part \mathcal{Y} as the domain of T , whereas we have shown that T then admits an extension to a linear subspace of \mathcal{X} that strictly contains \mathcal{Y} , which contradicts maximality of T . \square

In the proof of the next theorem we need the concept of Minkowski functional. Let E be a subset of \mathcal{X} . Then one defines $\mu_E(x) = \inf\{t > 0 : t^{-1}x \in E\}$. If E is absorbing (for all $x \in \mathcal{X}$ there exists $t > 0$ such that $tx \in E$), then μ_E is finite.

Lemma A.5 *Let E be an absorbing convex subset of a linear space \mathcal{X} . Then μ_E is a quasinorm and $\{\mu_E < 1\} \subset E$. If E is open, then $\{\mu_E < 1\} = E$.*

Proof Exercise. \square

Theorem A.6 *Let \mathcal{Y} and \mathcal{Z} be nonempty disjoint convex sets of a real normed linear space \mathcal{X} .*

- (i) *Assume that \mathcal{Y} is open. Then there exists a continuous linear functional T on \mathcal{X} and a number $\gamma \in \mathbb{R}$ such that*

$$Tx < \gamma \leq Ty, \forall x \in \mathcal{Y}, y \in \mathcal{Z}.$$

- (ii) *If \mathcal{Y} is compact and \mathcal{Z} is closed, then*

$$\sup\{Tx : x \in \mathcal{Y}\} < \inf\{Tx : x \in \mathcal{Z}\}.$$

Proof (i) Fix $y_0 \in \mathcal{Y}$ and $z_0 \in \mathcal{Z}$. Put $x_0 = z_0 - y_0$ and $\mathcal{C} = \mathcal{Y} - \mathcal{Z} + x_0$. Then \mathcal{C} is a convex neighbourhood of zero and $x_0 \notin \mathcal{C}$, since \mathcal{Y} and \mathcal{Z} are disjoint. Let p be the Minkowski functional of \mathcal{C} . By Lemma A.5 we know that p is a quasinorm and that $p(x_0) \geq 1$. Let \mathcal{X}_0 be the 1-dimensional subspace generated by x_0 . Define T_0 on \mathcal{X}_0 by $T_0(tx_0) = t$. Then T_0 is bounded and linear on \mathcal{X}_0 and $T_0(tx_0) \leq tp(x_0) = p(tx_0)$ for $t \geq 0$, whereas for $t < 0$ we have $T_0(tx_0) = t < 0 \leq p(tx_0)$. By Theorem A.4, T_0 can be extended to a linear map T on \mathcal{X} such that $T \leq p$. In particular, for $x \in \mathcal{C}$, we have $Tx \leq p(x) \leq 1$ and $T(-x) = -Tx \geq -1$. Hence $|Tx| \leq 1$ on $\mathcal{C} \cap (-\mathcal{C})$, so T is bounded on a neighbourhood of zero. But then T is continuous.

Let $y \in \mathcal{Y}$ and $z \in \mathcal{Z}$ be arbitrary. Since $Tx_0 = T_0x_0 = 1$, we have $Ty - Tz + 1 = T(y - z + x_0) \leq p(y - z + x_0)$. In view of Lemma A.5, $p(y - z + x_0) < 1$ since \mathcal{C} is open. It follows that $Ty < Tz$ for all $y \in \mathcal{Y}$ and $z \in \mathcal{Z}$. Define $\gamma = \sup\{Ty : y \in \mathcal{Y}\}$. Then $\gamma \leq Tz$ for all $z \in \mathcal{Z}$. But since \mathcal{Y} is open, also $\{Ty : y \in \mathcal{Y}\}$ is open and the supremum is not attained. This proves part (i).

To prove part (ii), we first notice that $d(\mathcal{Y}, \mathcal{Z}) > 0$, where d is the metric induced by the norm, since \mathcal{Y} is compact and \mathcal{Z} is closed. Hence, there exists $\delta > 0$ such that also the open δ -neighbourhood \mathcal{Y}^δ of \mathcal{Y} has disjoint intersection with \mathcal{Z} . Application of part (i) yields that $Ty < \gamma \leq Tz$ for all $y \in \mathcal{Y}^\delta$ and $z \in \mathcal{Z}$. But, since \mathcal{Y} is compact, we also have $\sup\{Ty : y \in \mathcal{Y}\} < \gamma$, which proves the second part. \square

We apply Theorem A.6 to the case where $\mathcal{X} = L^1(\Omega, \mathcal{F}, \mathbb{P})$. Recall that the dual space of a normed space is the linear space of all bounded linear functionals. It is well known that for $p \in [1, \infty)$ the dual space of $L^p(\Omega, \mathcal{F}, \mathbb{P})$ is isomorphic to $L^q(\Omega, \mathcal{F}, \mathbb{P})$ with $q = \frac{p}{p-1}$ for all $p \geq 1$, but we only need this fact for $p = 1$.

Lemma A.7 *The dual space of $L^1(\Omega, \mathcal{F}, \mathbb{P})$ is $L^\infty(\Omega, \mathcal{F}, \mathbb{P})$.*

Proof Let $T : L^1(\Omega, \mathcal{F}, \mathbb{P}) \rightarrow \mathbb{R}$ be a bounded linear operator. Define on \mathcal{F} the map ν by

$$\nu(F) = T(\mathbf{1}_F).$$

Obviously, ν is by linearity of T an additive map and by continuity of T even σ -additive. Indeed, if $F_n \downarrow \emptyset$, then $|\nu(F_n)| \leq \|T\| \mathbb{P}(F_n) \downarrow 0$. Hence ν is a finite signed measure on \mathcal{F} , that is absolute continuous w.r.t. \mathbb{P} . It follows from the Radon-Nikodym theorem that there is $Y \in L^1(\Omega, \mathcal{F}, \mathbb{P})$ such that

$$(A.5) \quad \nu(F) = \mathbb{E}[\mathbf{1}_F Y], \quad \forall F \in \mathcal{F}.$$

Next we show that Y is a.s. bounded. Let $F = \{Y > c\}$, for some $c > 0$. By continuity of T , we have

$$cP(Y > c) \leq \mathbb{E}[\mathbf{1}_F Y] = |T(\mathbf{1}_F)| \leq \|T\| \|\mathbf{1}_F\|_1 = \|T\| \mathbb{P}(Y > c).$$

Hence, if $\mathbb{P}(Y > c) > 0$ it follows that $\|T\| \geq c$. Stated otherwise $\|T\| \geq \sup\{c > 0 : \mathbb{P}(Y > c) > 0\}$. A similar argument yields $\|T\| \geq \sup\{c > 0 : \mathbb{P}(Y < -c) > 0\}$. It follows that $\|Y\|_\infty \leq \|T\| < \infty$.

We finally show that for every $X \in L^1(\Omega, \mathcal{F}, \mathbb{P})$, it holds that $T(X) = \mathbb{E}[XY]$. By the above construction this is true for X of the form $X = \mathbf{1}_F$. Hence also for (nonnegative) simple functions. Let $X \in L^1(\Omega, \mathcal{F}, \mathbb{P})$ be arbitrary. Choose a sequence (X_n) of simple functions such that $X_n \rightarrow X$ a.s. and $|X_n| \leq |X|$. Then, by dominated convergence, $X_n \rightarrow X$ in $L^1(\Omega, \mathcal{F}, \mathbb{P})$ as well and, since Y is a.s. bounded, we also have $X_n Y \rightarrow XY$ in $L^1(\Omega, \mathcal{F}, \mathbb{P})$. But then $T(X) = \mathbb{E}[XY]$. \square

Remark A.8 In the proof of Lemma A.7 one actually has that $\|Y\|_\infty = \|T\|$. The analogous version for $p \in (1, \infty)$ can be proven as follows. Start from Equation (A.5). It follows that for every $X \in L^\infty(\Omega, \mathcal{F}, \mathbb{P})$ one has $T(X) = \mathbb{E}XY$. For every $n \in \mathbb{N}$, put $X_n = \text{sgn}(Y)|Y|^{q-1}\mathbf{1}_{E_n}$, where $E_n = \{|Y| \leq n\}$. Note that every X_n is bounded, $X_n Y = |Y|^q \mathbf{1}_{E_n}$ and $|X_n|^p = |Y|^q \mathbf{1}_{E_n}$. We obtain

$$\mathbb{E}|Y|^q \mathbf{1}_{E_n} = T(X_n) \leq \|T\| \cdot \|X_n\|_p = \|T\| \cdot (\mathbb{E}|Y|^q \mathbf{1}_{E_n})^{1/p},$$

from which it follows that $(\mathbb{E}|Y|^q \mathbf{1}_{E_n})^{1/q} \leq \|T\|$ (here it is used that $p > 1$). By letting $n \rightarrow \infty$, we obtain $\|Y\|_q \leq \|T\| < \infty$, so $Y \in L^q(\Omega, \mathcal{F}, \mathbb{P})$.

Finally, for $X \in L^p(\Omega, \mathcal{F}, \mathbb{P})$ we put $X_n = X \mathbf{1}_{\{|X| \leq n\}}$, so that $X_n \in L^\infty(\Omega, \mathcal{F}, \mathbb{P})$ and $\|X - X_n\|_p \rightarrow 0$. It follows by Hölder's inequality (used in the fourth step) that

$$\begin{aligned} T(X) &= T(X - X_n) + T(X_n) \\ &= T(X - X_n) + \mathbb{E} X_n Y \\ &= T(X - X_n) + \mathbb{E} (X_n - X) Y + \mathbb{E} X Y \\ &\rightarrow \mathbb{E} X Y. \end{aligned}$$

Moreover, one also easily shows that actually $\|T\| = \|Y\|_q$.

Corollary A.9 *Let \mathcal{Y} and \mathcal{Z} be nonempty disjoint convex sets in $L^1(\Omega, \mathcal{F}, \mathbb{P})$. If \mathcal{Y} is compact and \mathcal{Z} is closed, there exists an almost surely bounded random variable B such that $\sup\{\mathbb{E}[YB] : Y \in \mathcal{Y}\} < \inf\{\mathbb{E}[ZB] : Z \in \mathcal{Z}\}$.*

Proof Combine Theorem A.6 and Lemma A.7. □

A.3 Existence of essential supremum

If (f_n) is a sequence of measurable functions on some measurable space (Ω, \mathcal{F}) , the function f defined by $f(\omega) = \sup_n f_n(\omega)$ is measurable as well. If instead of a sequence we take an arbitrary collection $\{f_i\}_{i \in I}$ measurability of the supremum is no longer guaranteed. But even if this happens, the pointwise supremum may have undesirable properties. Take for instance $(\Omega, \mathcal{F}, \mathbb{P}) = ([0, 1], \mathcal{B}, \lambda)$, where λ is Lebesgue measure. Let $f_x = \mathbf{1}_{\{x\}}$, $x \in [0, 1]$. Then $f = 1$, whereas each f_x is almost surely zero. Both observations trigger the following definition and theorem.

Theorem A.10 *Let $\{f_i\}_{i \in I}$ be an arbitrary collection of measurable functions on $(\Omega, \mathcal{F}, \mathbb{P})$. Then there exists a measurable function $f : \Omega \rightarrow \mathbb{R}$ such that*

- (i) $f \geq f_i$ a.s. for all $i \in I$
- (ii) If f' is any measurable function with the property that $f' \geq f_i$ a.s. for all $i \in I$, then $f' \geq f$ a.s.

The function f is thus a.s. unique and called the essential supremum of $\{f_i\}_{i \in I}$, notation: $f = \text{ess sup}\{f_i\}_{i \in I}$.

Proof Without loss of generality we may take the collection $F := \{f_i\}_{i \in I}$ bounded, apply an arctan-transformation if needed. Let F' be a countable sub-collection of F , then $f^{F'} := \sup\{f : f \in F'\}$ is measurable. Let

$$s := \sup\{\mathbb{E} f^{F'} : F' \subset F, F' \text{ countable}\}.$$

Choose a sequence of countable $F'_n \subset F$ such that $\mathbb{E} f^{F'_n} \rightarrow s$ and let $F'_\infty = \cup_n F'_n$. Then F'_∞ is countable and $\mathbb{E} f^{F'_\infty} = s$. Let $f = f^{F'_\infty}$. If (i) is not true for

this f , then there exists $i^* \in I$ such that $\mathbb{P}(f < f_{i^*}) > 0$. Put $F^* = F'_\infty \cup \{f_{i^*}\}$. Then $\mathbb{E} f^{F^*} > \mathbb{E} f = s$, contradicting the definition of s . Hence we conclude that (i) holds. Let now f' be measurable such that $f' \geq f_i$ a.s. for all $i \in I$, then obviously $f' \geq f_i$ for $i \in F'_\infty$ a.s. and hence $f' \geq f$, which proves (ii). \square

A.4 Results on the weak topology for measures

Let S be a complete separable metric space with metric d and let \mathcal{S} be its Borel σ -field. Denote by $\mathcal{M}_1(\mathcal{S})$ the set of all probability measures on \mathcal{S} . By $\mathcal{D}(\mathcal{S})$ we denote the subset of $\mathcal{M}_1(\mathcal{S})$ that consists of all (finite) convex combinations of Dirac measures. Recall that a subset E of a metric space is *totally bounded*, if for every $\varepsilon > 0$, there exist finitely many open balls of radius ε whose union contains E . Recall also that a subset of a metric space is compact iff it is totally bounded and complete.

Lemma A.11 *Let μ be a probability measure on (S, \mathcal{S}) . Then μ is tight, i.e. for each $\varepsilon > 0$ there exists a compact set K such that $\mu(K) > 1 - \varepsilon$.*

Proof For each $n \in \mathbb{N}$ there is a countable family of open balls B_{nj} , each with radius $\frac{1}{n}$, that covers S . Let $U_n = \bigcup_{j=1}^n B_{nj}$, then $U_n \uparrow S$ and hence for all $\varepsilon > 0$, there is m_n such that $\mathbb{P}(U_{nm_n}) > 1 - 2^{-n}\varepsilon$. Let $D = \bigcap_{n=1}^{\infty} U_{nm_n}$, then D is obviously totally bounded, and so is its closure K . But since K , being a closed subset of S , is complete, it is then also compact. Moreover, $\mu(K^c) \leq \mu(D^c) \leq \sum_{n \geq 1} \mu(U_{nm_n}^c) < \varepsilon$. \square

Proposition A.12 *Let $\mu \in \mathcal{M}_1(\mathcal{S})$. Then there exists a sequence $(\mu_n) \subset \mathcal{D}(\mathcal{S})$ such that $\mu_n \rightarrow \mu$ weakly.*

Proof We first show that the assertion holds, if $\mu(K^c) = 0$ for some compact set K . For every n we can cover K by a finite set of open balls $B_{n,j}$, $j = 1, \dots, k_n$ each having radius $\frac{1}{n}$. Put $A_{n,1} = B_{n,1} \cap K$ and for $j > 1$, recursively, $A_{n,j} = B_{n,j} \cap K \setminus (\bigcup_{i=1}^{j-1} A_{n,i})$. Then all $A_{n,j}$ are contained in $B_{n,j}$ and $\bigcup_{j=1}^{k_n} A_{n,j} = K$. Ignoring the j for which $A_{n,j}$ is empty, we select $x_{n,j} \in A_{n,j}$. Put

$$\mu_n = \sum_j \mu(A_{n,j}) \delta_{x_{n,j}},$$

where $\delta_{x_{n,j}}$ is the Dirac measure concentrated at $x_{n,j}$. Note that also μ_n is concentrated on K . Let $f \in C_b(S)$. Since μ is concentrated on K , we have

$$\int f \, d\mu = \sum_j \int_{A_{n,j}} f \, d\mu.$$

Since f is uniformly continuous on K , we have that

$$\eta_n := \sup\{|f(x) - f(y)| : d(x, y) < \frac{1}{n}\} \rightarrow 0.$$

Hence

$$\begin{aligned}
\left| \int_{A_{n,j}} f \, d\mu - \mu(A_{n,j})f(x_{n,j}) \right| &= \left| \int_{A_{n,j}} (f - f(x_{n,j})) \, d\mu \right| \\
&\leq \int_{A_{n,j}} |f - f(x_{n,j})| \, d\mu \\
&\leq \eta_n \mu(A_{n,j}).
\end{aligned}$$

By summing over j we obtain

$$\left| \int f \, d\mu - \int f \, d\mu_n \right| \leq \eta_n \mu(K) = \eta_n,$$

which yields the result.

For an arbitrary probability measure $\mu \in \mathcal{M}_1(S)$ we argue as follows. Let $\varepsilon > 0$. In view of Lemma A.11, there exists a compact set K such that $\mu(K) > 1 - \varepsilon$. Define the (conditional) probability measure μ' by $\mu'(B) = \mu(B|K)$. Let $f \in C_b(S)$ and let μ_n be the measures as in the first part of the proof (with μ' replacing μ). Then

$$\begin{aligned}
\left| \int f \, d\mu - \int f \, d\mu_n \right| &\leq \varepsilon \|f\| + |\mu(K) \int_K f \, d\mu' - \int_K f \, d\mu_n| \\
&\leq \varepsilon \|f\| + |(\mu(K) - 1) \int_K f \, d\mu'| + \left| \int_K f \, d\mu' - \int_K f \, d\mu_n \right| \\
&\leq 2\varepsilon \|f\| + \left| \int_K f \, d\mu' - \int_K f \, d\mu_n \right|.
\end{aligned}$$

Since the last term vanishes in view of the first part of the proof, the conclusion of the theorem follows. \square

Corollary A.13 *If S_0 is a countable dense subset, then we can choose the $x_{n,j}$ in the proof of Proposition A.12 in S_0 . Moreover, we can approximate μ with a convex mixture of Dirac distributions, where the mixing coefficients are rational.*

Proof Obvious. \square

A.5 Proofs of results of Section 9.1

This section contains proofs of results in Section 9.1. We shall use the following result.

Lemma A.14 *If \mathcal{G} is a σ -algebra, X is a \mathcal{G} -measurable random variable (or vector), Y is independent of \mathcal{G} and f is a measurable function such that the expectations below exist, then $\mathbb{E}[f(X, Y)|\mathcal{G}] = \hat{f}(X)$, where $\hat{f}(x) = \mathbb{E}f(x, Y)$.*

So, the conditional expectation is obtained by taking expectation w.r.t. Y , i.e. integrating out the variable Y only, and leaving X untouched. We continue with the proofs of the technical results.

Proof See measure theory for the general case and Exercise A.2 for a simple special case. \square

Lemma A.15 Suppose v_0, \dots, v_T are functions on \mathbb{R}^d satisfying

$$\begin{aligned} v_T(x) &\geq g_T(x), \forall x \in \mathbb{R}^d, \\ v_t(x) &\geq g_t(x, y) + \hat{v}_{t+1}(x, y), \forall (x, y) \in \mathbb{R}^d \times \mathbb{R}^m, \forall t \in \{0, \dots, T-1\}. \end{aligned}$$

Then for all sequences u it holds that

$$(A.6) \quad v_t(X_t^u) \geq J_t(u) \text{ a.s., } t = 0, \dots, T,$$

and $\mathbb{E} v_0(X_0) \geq J(u)$.

Proof Let $t = T$. Then $v_T(X_T^u) \geq g_T(X_T^u) = \mathbb{E}[g_T(X_T^u)|\mathcal{F}_T] = J_T(u)$ and (A.6) holds for $t = T$. We proceed by backward induction, so assume that (A.6) holds true at times $t+1, \dots, T$. Notice that by independence of ε_t and \mathcal{F}_t and Lemma A.14 it holds that

$$(A.7) \quad \hat{v}_{t+1}(X_t^u, U_t) = \mathbb{E}[v_{t+1}(f_t(X_t^u, U_t, \varepsilon_t))|\mathcal{F}_t] = \mathbb{E}[v_{t+1}(X_{t+1}^u)|\mathcal{F}_t].$$

Then, by the assumption on v_t , (A.7), and the induction assumption,

$$\begin{aligned} v_t(X_t^u) &\geq g_t(X_t^u, U_t) + \mathbb{E}[v_{t+1}(X_{t+1}^u)|\mathcal{F}_t] \\ &\geq \mathbb{E}[g_t(X_t^u, U_t) + J_{t+1}(u)|\mathcal{F}_t] \\ &= \mathbb{E}[g_t(X_t^u, U_t) + \mathbb{E}[\sum_{s=t+1}^{T-1} g_s(X_s^u, U_s) + g_T(X_T^u)|\mathcal{F}_{t+1}]|\mathcal{F}_t] \\ &= \mathbb{E}[\sum_{s=t}^{T-1} g_s(X_s^u, U_s) + g_T(X_T^u)|\mathcal{F}_t] \\ &= J_t(u). \end{aligned}$$

This shows (A.6). Applying this inequality for $t = 0$ and taking expectations yields the final assertion. \square

Lemma A.16 Suppose that the U_t are such that $U_t = u_t(X_t)$, for some measurable functions $u_t : \mathbb{R}^d \rightarrow \mathbb{R}^m$, and let \mathcal{M} be the class of sequences $u = (u_0, \dots, u_{T-1})$ of such functions. Suppose that for all $u \in \mathcal{M}$ one defines functions v_0^u, \dots, v_T^u by

$$\begin{aligned} v_T^u(x) &= g_T(x) \\ v_t^u(x) &= g_t(x, u_t(x)) + \mathbb{E} v_{t+1}^u(f_t(x, u_t(x), \varepsilon_t)), t = 0, \dots, T-1. \end{aligned}$$

Then it holds that X^u is Markov w.r.t. to the given filtration, $v_t(X_t^u) = J_t(u)$ for $t = 0, \dots, T$ and

$$J_t(u) = \mathbb{E}[\sum_{s=t}^{T-1} g_s(X_s^u, u_s(X_s^u)) + g_T(X_T^u)|X_t^u].$$

Proof Similar to the proof of Lemma A.15 upon noting that the conditional expectations w.r.t. \mathcal{F}_t reduce to conditional expectations w.r.t. X_t^u in view of Lemma 9.1. \square

Proof of Theorem 9.6 (i) This assertion directly follows from Lemma A.15.

(ii) Suppose that the supremum in (9.4) is attained at $y = u_t^*(x)$. Then, for all $t \in \{0, \dots, T-1\}$,

$$v_t(x) = g_t(x, u_t^*(x)) + \hat{v}_{t+1}(x, u_t^*(x)).$$

We now apply Lemma A.16 to obtain $\mathbb{E} v_0(X_0) = \mathbb{E} J_0(u^*) = J(u^*)$. Since for every other $u \in \mathcal{M}$, it holds that $J(u) \leq \mathbb{E} v_0(X_0)$, it follows that u^* is optimal. Likewise one shows that $v_t(X_t^{u^*}) = J_t(u^*)$.

Conversely, assume that $u^* \in \mathcal{M}$ is optimal. Let $t = T-1$ and $X_{T-1}^{u^*}$ be the current state. Suppose that the supremum in (9.4) is not attained for $u_{T-1}^*(X_{T-1}^{u^*})$ with positive probability. From the definition of v_{T-1} it follows that there exists some $\tilde{u}_{T-1}(X_{T-1}^{u^*})$ such that a.s.

$$\begin{aligned} & g_{T-1}(X_{T-1}^{u^*}, \tilde{u}_{T-1}(X_{T-1}^{u^*})) + \hat{v}_T(X_{T-1}^{u^*}, \tilde{u}_{T-1}(X_{T-1}^{u^*})) \\ & \geq g_{T-1}(X_{T-1}^{u^*}, u_{T-1}^*(X_{T-1}^{u^*})) + \hat{v}_T(X_{T-1}^{u^*}, u_{T-1}^*(X_{T-1}^{u^*})), \end{aligned}$$

where the inequality is strict with positive probability. After taking expectations, a strict inequality emerges. We claim that $\tilde{u} = (u_0^*, \dots, u_{T-2}^*, \tilde{u}_{T-1})$ is a sequence that yields a higher performance than u^* . Indeed $X_t^{\tilde{u}} = X_t^{u^*}$ for all $t \leq T-1$ and hence $\mathbb{E} g_t(X_t^{\tilde{u}}, \tilde{u}_t(X_t^{\tilde{u}})) = \mathbb{E} g_t(X_t^{u^*}, u_t^*(X_t^{u^*}))$ for $t \leq T-2$, whereas

$$\begin{aligned} & \mathbb{E} g_{T-1}(X_{T-1}^{\tilde{u}}, \tilde{u}_{T-1}(X_{T-1}^{\tilde{u}})) + \mathbb{E} \hat{v}_T(X_{T-1}^{\tilde{u}}, \tilde{u}_{T-1}(X_{T-1}^{\tilde{u}})) \\ & > \mathbb{E} g_{T-1}(X_{T-1}^{u^*}, u_{T-1}^*(X_{T-1}^{u^*})) + \mathbb{E} \hat{v}_T(X_{T-1}^{u^*}, u_{T-1}^*(X_{T-1}^{u^*})). \end{aligned}$$

But then $J(\tilde{u}) > J(u^*)$, which contradicts optimality of u^* . Hence we must have $J_{T-1}(u^*) = v_{T-1}(X_{T-1}^{u^*})$ by virtue of Lemma A.16.

One proceeds by induction. Suppose that the supremum in (9.4) is attained for $u_s^*(X_s^{u^*})$ with probability one, for all $t+1 \leq s \leq T-1$, but that, with positive probability, this is not true for $s = t$. Then one argues as above that there exists a \tilde{u}_t such that with $\tilde{u} = (u_0^*, \dots, u_{t-1}^*, \tilde{u}_t, u_{t+1}^*, \dots, u_{T-1}^*)$ it holds that $J(\tilde{u}) > J(u^*)$, which again contradicts optimality of u^* .

In passing, we have also shown that $v_t(X_t^{u^*}) = J_t(u^*)$ and $\sup_u J(u) = J(u^*) = \mathbb{E} v_0(X_0)$. \square

A.6 Exercises

A.1 This concerns the statement $\lambda \geq 1$ in the proof of Theorem A.1. Define $x_t = tx + (1-t)x_0$, $t \in [0, 1]$. For which t is $\|x_t\|$ minimal? Exploit this to prove the assertion.

A.2 Prove Lemma A.14 for the special case in which X and Y are discrete random variables.

Index

- Arrow-Pratt coefficient, 29
- atom, 8
- attainable pay-off, 3
- axiom
 - Archimedean, 20
 - independence, 20
- Bellman's optimality principle, 85
- budget constraint, 40
- certainty equivalent, 27
- consumption process, 90
 - attainable, 94
- consumption-investment plan, 90
 - admissible, 90
- contingent claim, 72
 - attainable, 72
- Cox-Ross-Rubinstein model, 79
- discounted net gains, 2
- dynamic programming, 85
- equivalent martingale measure, 2, 62
- essential supremum, 88, 104
- Fundamental Theorem of Asset Pricing
 - first, 2, 64
 - second, 8, 77
- Hahn-Banach theorem, 71, 101
- indifference relation, 12
- Kullback-Leibler information, 46
- market
 - arbitrage free, 1, 62
 - complete, 8, 77
 - extended, 4, 74
 - non-redundant, 5, 42
- martingale measure, 2, 62
- numéraire, 65
- numerical representation, 13
 - expected utility, 27
 - Von Neumann-Morgenstern, 20
- order dense, 13
- pay-off
 - admissible, 50
 - attainable, 3
- portfolio, 1
 - arbitrage opportunity, 1
 - hedge, 7
 - replicating, 7
 - value, 1
- preference intervals, 15
- preference order
 - monotone, 26
 - risk averse, 26
 - uniform, 33
- preference relation
 - continuous, 15
 - strict, 12
 - weak, 12
- pricing rule, 50
- relative entropy, 46
- risk premium, 27
- risk-neutral measure, 2, 62
- separating hyperplane, 75, 100
- stochastic dominance
 - first order, 37
 - second order, 33
- topology
 - ψ -weak, 24
 - weak, 23
- trading strategy, 60
 - arbitrage opportunity, 62
 - discounted gains process, 61
 - discounted value process, 61
 - hedge, 72
 - replicating, 72
 - self-financing, 60
- utility function, 26
 - HARA, 30
 - Inada conditions, 30
- value function, 88